Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2021-eular.550

POS1337 **BEHÇET'S DISEASE AND COURSE OF PREGNANCY - RESULTS FROM THE NATIONWIDE PREGNANCY REGISTER REVNATUS**

H. S. Koksvik¹, B. Jakobsen¹, H. Bjørngaard¹, C. Skorpen^{2,3}, M. Wallenius^{1,2}.
¹St. Olavs Hospital, Trondheim University Hospital, Norwegian National Advisory Unit on Pregnancy and Rheumatic Diseases, Trondheim, Norway; ²NTNU, Department of Neuromedicine and Movement Science, Trondheim, Norway; ³Alesund Hospital, Department of Rheumatology, Alesund, Norway

Background: Behçet's disease (BD) is a complex and multisystemic disorder of unknown etiology. It is characterized by recurrent oral and genital ulcers and inflammatory eye involvement. It can also affect the cardiovascular, pulmonary and skeletal system. BD is commonly diagnosed in women in reproductive age. Pregnancy outcomes are reported in previous studies with discrepancies in the rate of pregnancy-related complications.

Objectives: To describe the course of pregnancy in patients with BD.

Methods: We prospectively collected data on pregnancy outcome in women with BD through the nationwide Norwegian registry on pregnancy and rheumatic diseases (RevNatus) from 2016-2020. All women were diagnosed with ICD-10 code M35.2. 13 pregnancies in 11 women were recorded. One woman had two pregnancies including one twin birth.

Results: Median age at conception was 30 years (range 25-32), with median disease duration 7 years (range 1-15). There were no serious organ manifestations. Tnf-alpha-inhibitors were used in 2nd trimester in 3 pregnancies. All pregnancies resulted in live births with median gestational age at delivery of 38 weeks (range 36-41). No preeclampsia or HELLP were recorded.

Table 1. Pregnancy outcome in 13 BD pregnancies

Patient #	Pregnancy #	Medication at 2.nd trimester	Oral or genital ulcers during pregnancy	Disease activity during pregnancy	CRP value at 2nd trimester	Gestational age at delivery, week	Mode of delivery	Birth weight, gram
A	1	Azathioprine Colchicine LMWH	no	persistent*	13	38	SVD	3475
B	2	Adalimumab Colchicine LDPrednisolone	yes	persistent *	4	40	SVD	3215
C	3	no	yes	persistent*	5	41	SVD	3355
D	4	Adalimumab LDPrednisolone	yes	persistent*	23	36	SVD	1940
E	5	Etanercept ASA	no		2	41	SVD	3410
	6a (twin)	Etanercept ASA	no		5	37	Elective Sectio	2400
	6b (twin)							2410
F	7	LDPrednisolone	yes	flare**	15	38	SVD	3180
	8	Adalimumab LDPrednisolone	yes	flare**	4	37	Acute Sectio	3049
G	9	no	no	remission***	1	41	SVD	3150
H	10	LDPrednisolone	yes	persistent *	10	38	SVD	2480
I	11	ASA	no	persistent*	3	39	SVD	3410
J	12	no	yes	persistent*	1	37	SVD	3115
K	13	Ciklosporine	no	remission***	3	37	SVD	3130

SVD = spontaneous vaginal delivery, ASA = acetylsalicylic acid, LDPrednisolone = Low dose Prednisolone <10 mg/day, * = persistent throughout pregnancy, ** = flare once during pregnancy, *** = remission throughout pregnancy

Conclusion: Pregnancies in women with BD in this cohort were uncomplicated. We did not reveal any serious events, and they all resulted in live births. However, BD is a very heterogeneous disease, and the women in our cohort did not have any serious organ manifestations. Larger cohorts representing the heterogeneity are needed to study the course and outcome of pregnancies in BD.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.666

POS1338 **INCIDENCE, PREVALENCE, AND MORTALITY OF CHRONIC PERIAORTITIS: A POPULATION-BASED STUDY**

U. Ghaffar¹, K. J. Warrington¹, S. Duong², C. S. Crowson^{1,2}, M. Burke¹, B. Viers³, A. Potretzke³, H. Bjarnason⁴, M. Koster¹. ¹Mayo Clinic College of Medicine and Science, Rheumatology, Rochester, United States of America; ²Mayo Clinic School of Health Sciences, Biomedical Statistics and Informatics,

Rochester, United States of America; ³Mayo Clinic College of Medicine and Science, Urology, Rochester, United States of America; ⁴Mayo Clinic College of Medicine and Science, Vascular and Interventional Radiology, Rochester, United States of America

Background: Chronic periaortitis is an inflammatory condition that typically involves the infrarenal portion of the abdominal aorta. Few studies are available to outline the epidemiology of this rare condition. To date, no epidemiologic studies on periaortitis have been performed in North America.

Objectives: To evaluate the epidemiology, presentation and outcomes of patients with chronic periaortitis from 1998 through 2018.

Methods: An inception cohort of patients with incident chronic periaortitis from January 1, 1998 through December 31, 2018, in Olmsted County, Minnesota, USA, was identified based on comprehensive individual medical record review utilizing the Rochester Epidemiology Project medical record linkage system. Inclusion required radiographic and/or histologic confirmation of periaortic soft tissue thickening around at least part of the infra-renal abdominal aorta or the common iliac arteries. Data were collected on demographic characteristics, clinical presentation, renal and radiographic outcomes, and mortality. Incidence rates were age and sex adjusted to the 2010 United States white population.

Results: Eleven incident cases of chronic periaortitis were identified during the study period. Mean±SD age at diagnosis was 61.8±13.4 years. The cohort included 9 men (82%) and 2 women (18%). The most common presenting symptom was pain with 55% (6/11) reporting abdominopelvic-pelvic pain, 36% (4/11) back pain, and 18% (2/11) flank pain. Obstructive uropathy was present in 73% (8/11) subjects: 3 (27%) unilateral left, 1 (9%) unilateral right, and 4 (36%) bilateral. Mean creatinine at presentation was 2.7±3.4 mg/dL. Ureteral stenting was required at diagnosis in seven patients: unilateral left in 2, unilateral right in 1 and bilateral in 4. All 11 patients received glucocorticoids with a median (IQR) dose of 40 (30, 60) mg/day. Additional non-glucocorticoid therapeutics were used in 10 patients.

Renal function stage at last follow up declined in 2 patients, remained the same in 3 patients and improved in 6 patients. Mean creatinine at last follow-up was 1.2±0.2 mg/dL. Among the seven patients requiring baseline indwelling ureteral stent placement only two required ongoing ureteral stenting at last follow up. None of the four patients without ureteral stenting at diagnosis progressed to require stenting during the follow-up period. No patient underwent ureterolysis surgery in this cohort. Periaortic soft tissue thickening at last follow up had increased in thickness in 1 (9%), was unchanged in 2 (18%), decreased in size but did not resolve in 6 (55%), and fully resolved in 18%.

Age- and sex-adjusted incidence rates per 100,000 population were 0.26 for females, 1.56 for males and 0.87 overall. Overall prevalence on January 1, 2015 was 8.98 per 100,000 population. Median (IQR) length of follow-up was 10.1 (2.5, 13.8) years. Overall mortality was similar to the expected age, sex, and calendar estimates of the Minnesota population with standardized mortality ratio (95% CI) for the entire cohort 2.07 (0.67, 4.84).

Conclusion: This study reports the first epidemiologic data on chronic periaortitis in the United States. In this cohort of patients with chronic periaortitis, men were approximately 4 times more commonly affected than women. Mortality was not increased compared to the general population.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.698

POS1339 **MORE FREQUENT AND EARLIER HIP INVOLVEMENT IN SPONDYLOARTHRITIS ASSOCIATED WITH FAMILIAL MEDITERRANEAN FEVER**

M. Bektaş¹, S. San¹, C. Cetin¹, B. Ç. Yalçın Dulundu¹, N. Koca¹, B. Ince¹, Y. Yalçınkaya¹, B. Artım-Esen¹, M. Inanc¹, M. L. Ocal¹, A. Gül¹. ¹Istanbul Faculty of Medicine, Internal Medicine, Rheumatology Division, Istanbul, Turkey

Background: Familial Mediterranean fever (FMF) is a hereditary autoinflammatory disorder caused by the MEFV gene variants. Although association between FMF and spondyloarthritis (SpA) has previously been reported, clinical and laboratory features of patients with FMF and SpA have not been defined in detail.

Objectives: We aimed to evaluate clinical and laboratory characteristics, disease outcome and biologic responses of patients with FMF+SpA compared to patients with only SpA who were followed-up in our tertiary referral center.

Methods: Database of FMF Clinic was screened for FMF patients with coexistent SpA and 113 patients were identified fulfilling Tel Hashomer and ASAS criteria for FMF and SpA, respectively. A group of patients with SpA without FMF matched for age, gender and disease duration were selected as the control group.

Results: Thirteen patients were excluded because of missing data, and 100 patients (F/M: 52/48) were included into the analysis. Mean follow-up time was 93.6 ± 77 (range [r]: 3-324) months and mean patient age was 43.3 ± 12 (r: 20-87) years. Mean age of onset for FMF was 12.5 ± 8 (r: 1-36) and for SpA was 25 ± 11 (r: 7-72) years. SpA findings was classified as axial in 35.4%, axial and peripheral in 47.9% and only peripheral in 16.7% in FMF+SpA group. Half (49%) of the