González¹, Universidad Autónoma de Nuevo León, Rheumatology Service, Monterrey, Mexico; ²Hospital Universitario "Dr. José Eleuterio González", Universidad Autónoma de Nuevo León, Cardiology Service, Monterrey, Mexico

Background: Systemic lupus erythematosus (SLE) is a chronic and autoimmune disease characterized by systemic involvement. Patients with SLE have accelerated atherosclerosis, resulting in an up to nine-fold increased risk of cardiovascular disease, compared to the general population (1), being the leading cause of death for these patients. Speckle tracking echocardiography (STE) is an accurate technique to estimate myocardial function and deformation.

Objectives: This study aims to determine the association between echocardiographic findings and the presence of antibodies in SLE patients.

Methods: This was a cross-sectional and observational study. A total of forty-three patients ≥ 18 years with a diagnosis of SLE according to EULAR/ACR 2019 criteria were included for this study. Those with a history of cardiovascular disease (myocardial infarction, cerebrovascular accident, or peripheral arterial disease) and pregnancy were excluded. Transthoracic echocardiogram was performed and reviewed by 2 board-certified cardiologists, in all study subjects. Blood samples obtained from all patients were analyzed for the following: anti-nuclear antibodies (ANA), anti-SSA/Ro, SSB/La antibodies, anti-cardiolipin antibodies (IgA, IgM, IgG), and complement levels. Distribution was evaluated with the Kolmogorov-Smirnov test. Correlations between numerical variables were done using Spearman's rho, considering two-tailed p-values < 0.05 as statistically significant.

Results: The 39 female patients (90.7%) and 4 male patients (9.3%) had a mean age of 35.5 ± 12.0 years and a median disease duration of 72 months (14-132). At the time of inclusion, 90.7% of the patients were being treated with glucocorticoids and anti-malarials. Concerning traditional cardiovascular risk factors; 20.9% of the patients had hypertension, 70% had dyslipidemia, 2.3% had diabetes mellitus and 18.6% were active smokers. Correlations between echocardiographic findings and antibodies are shown in Table 1. We found a moderate positive correlation between global circumferential strain and IgA anticardiolipin antibody ($r=0.507$, $p=0.002$), a low positive correlation in left ventricular ejection fraction with anti-Ro ($r=0.397$, $p=0.012$) and anti-La ($r=0.397$, $p=0.012$) and a low positive correlation between TAPSE and C3 levels ($r=0.396$, $p=0.013$).

Conclusion: There is an association between anticardiolipin antibody titers, anti-Ro, and anti-La with echocardiographic alterations. All SLE patients especially those who had positive antibodies should be screened for the presence of structural cardiac abnormalities. STE can be helpful as a noninvasive diagnostic tool, that could result in earlier treatment and prognosis.

REFERENCES:

- [1] Hesselvig JH, Ahlehoff O, Dreyer L, et al. Cutaneous lupus erythematosus and systemic lupus erythematosus are associated with clinically significant cardiovascular risk: a Danish nationwide cohort study. *Lupus* 2017;26(1):48-53. doi: 10.1177/0961203316651739

Table 1. Spearman rho correlations between antibody titers and echocardiographic findings.

Variables	GLS, mean $\pm$ SD -19.11 $\pm$ 3.33	LVEF, mean $\pm$ SD 57.43 $\pm$ 7.17	TAPSE, mean $\pm$ SD 22.23 $\pm$ 3.24
ANA, median (p25-p75) 640 (160-2550)	NS	NS	NS
IgA Anti-Cardiolipin, median (p25-p75) 2 (2-3)	0.507**	NS	NS
IgM Anti-Cardiolipin, median (p25-p75) 2 (2-4)	NS	NS	NS
IgG Anti-Cardiolipin, median (p25-p75) 4 (3-6)	NS	NS	NS
Anti-Ro, median (p25-p75) 17 (2-80)	NS	0.326*	NS
Anti-La, median (p25-p75) 3 (2-5.5)	NS	0.397*	NS
C3, mean $\pm$ SD 91.41 $\pm$ 37.38	NS	NS	0.396*

**Correlation is significant at the 0.01 level (2-tailed). *Correlation is significant at the 0.05 level. NS, not significant; GLS, global circumferential strain; LVEF, left ventricular ejection fraction; TAPSE, tricuspid annular plane systolic excursion.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3399

POS1405

OPTICAL SPECTRAL TRANSMISSION (HANDSCAN) IN PATIENTS WITH PSORIATIC ARTHRITIS: FIRST DATA ON DIAGNOSTIC VALUE AND ASSOCIATIONS WITH CLINICAL AND ULTRASOUND DISEASE ACTIVITY MARKERS

K. Triantafyllias¹, S. Liverakos², C. Noack³, A. Schwarting^{1,2}. ¹ACURA Clinics, Rheumatology, Bad Kreuznach, Germany; ²Johannes Gutenberg University Medical Center, Internal Medicine I, Department of Rheumatology and Clinical

Immunology, Mainz, Germany; ³University of Mannheim, Graduate School of Economics and Social Sciences, Mannheim, Germany

Background: Valid assessment of disease activity leads to improvement of long-term outcomes in patients with inflammatory arthritis (1). Optical spectral transmission (OST) is a modern diagnostic tool able to assess the blood-specific absorption of light transmitted through a tissue, promising quantification of inflammation in the finger and wrist joints of patients with rheumatoid arthritis (RA) (commercial device: HandScan – Demcon/Hemics, The Netherlands) (2). Even though an increasing number of studies have evaluated diagnostic value of this new technology in RA patients (2,3), no data exist regarding psoriatic arthritis (PsA).

Objectives: To examine for the first time the diagnostic value of OST in detecting inflammation in patients with PsA and to evaluate its relationship with disease activity markers and various epidemiological and anthropometric patient characteristics.

Methods: OST-Measurements were performed in a group of PsA patients and a group of healthy controls. The difference between OST in the two groups was statistically examined and relationships of OST with clinical (tender / swollen joint counts, disease activity on a visual analogue scale) and serological disease activity markers were evaluated. Moreover, joint ultrasound (US) examinations were performed in a subgroup of PsA patients and OST associations with a Power Doppler- and a Grey Scale-US score were examined. Finally, relationships of OST with various anthropometric and epidemiologic parameters (BMI, hand-size, gender, age) were assessed.

Results: We recruited 49 PsA patients [65.3% female; mean age 53.3 years (± 11.8 SD)] and 114 control subjects [77.2% female; mean age 46 years (± 12.8 SD)]. OST was statistically significantly higher in the patient group, compared to the control group [14.95 (12.04 - 17.18, IQR) vs. 10.31 (7.84 - 13.79, IQR); $p < 0.001$]. OST correlated moderately-strongly with both examined US scores (Power Doppler-score: $r = 0.5$; $p = 0.026$ and Grey Scale-score: $r = 0.52$; $p = 0.028$). Moreover, OST showed a moderate, statistically significant association with C-reactive protein (CRP) ($r = 0.298$; $p = 0.037$). Finally, males had significantly higher OST values than females and OST associated moderately-weakly with body mass index (BMI) in the control group ($\rho = 0.24$; $p < 0.001$).

Conclusion: This is the first report of a possible diagnostic value of OST in patients with PsA. OST correlated with ultrasound and serological activity markers and may thus prove to be a useful tool of disease activity assessment, next to well established diagnostic modalities, such as the joint US. Correlations of OST with patient characteristics implicate the need to take also anthropometric and epidemiological patient characteristics into account when interpreting OST results in order to avoid confounding.

REFERENCES:

- [1] Katchamart W, et al. Systematic monitoring of disease activity using an outcome measure improves outcomes in rheumatoid arthritis. *J Rheumatol* 2010;37:1411-1415.
- [2] Triantafyllias, et al. Diagnostic value of optical spectral transmission in rheumatoid arthritis: associations with clinical characteristics and comparison with joint ultrasonography. *J Rheumatol* 2020 1;47(9):1314-1322.
- [3] Onna M Van, et al. Assessment of disease activity in patients with rheumatoid arthritis using optical spectral transmission measurements, a non-invasive imaging technique. *Ann Rheum Dis* 2016;75:511-518.

Disclosure of Interests: Konstantinos Triantafyllias Speakers bureau: Pfizer, Novartis, Janssen, Chugai, Stefanie Liverakos: None declared, Claudia Noack: None declared, Andreas Schwarting: None declared

DOI: 10.1136/annrheumdis-2021-eular.3670

POS1406

DEVELOPMENT OF A DIAGNOSTIC ALGORITHM FOR THE DIFFERENTIAL DIAGNOSIS OF INTERSTITIAL LUNG DISEASE: PRELIMINARY DATA FROM A MULTICENTER RETROSPECTIVE CASE-CONTROL STUDY

B. Maranini¹, T. Chiodin¹, C. A. Scirè¹, M. Govoni¹, E. Lucioni², S. Chiarello², F. Scabbia², I. Marchi³, G. Zanframundo⁴, L. Cavagna⁴, E. Bellis⁴, M. Silva⁵, G. Tringali⁵, A. Carnevale². ¹Rheumatology Unit, University of Ferrara and Azienda Ospedaliero-Universitaria S. Anna, Department of Medical Sciences, Ferrara, Italy; ²Radiology Unit, Azienda Ospedaliero-Universitaria S. Anna, Department of Radiology, Ferrara, Italy; ³Pneumology Unit, University of Ferrara and Azienda Ospedaliero-Universitaria S. Anna, Department of Medical Sciences, Ferrara, Italy; ⁴University and IRCCS Policlinico S. Matteo Foundation of Pavia, Department of Rheumatology, Pavia, Italy; ⁵Department of Medicine and Surgery (DiMeC), University of Parma, Scienze Radiologiche, Parma, Italy

Background: Interstitial lung diseases (ILDs) represent a heterogeneous group of disorders with different treatment and prognosis. ILD may be the presenting or