of Southern Denmark, Aabenraa, Denmark; ⁵Section of Rheumatology, Department of Clinical Sciences Lund, Lund University, Lund, Denmark; ⁶Spenshult, Research and Development Centre, Halmstad, Sweden; ⁷Research Initiative for Activity Studies and Occupational Therapy, Research Unit of General Practice, Department of Public Health, University of Southern Denmark, Odense, Denmark; ⁸Research Unit of User Perspectives, Department of Public Health, University of Southern Denmark, Odense, Denmark

Background: In people with rheumatoid arthritis (RA), modifiable lifestyle factors such as smoking, being overweight/obese, alcohol overuse and physical inactivity may not only affect treatment response and quality of life, but can also increase the risk for cardio-vascular diseases and other comorbidities (1,2). Evidence and EULAR guidelines (3) support lifestyle changes in patients with RA. If a patient need to change several habits, the challenge may seem overwhelming and substantial support will be needed. There is little information concerning the prevalence of a combined number of unhealthy lifestyle (UL) factors in people with RA. **Objectives:** I) To study the prevalence of unhealthy lifestyle factors in two Scandinavian RA cohorts. II) To study the association between disease impact and two or more unhealthy lifestyle factors.

Methods: Patients diagnosed with RA who participated in a cardiovascular screening consultation at a specialist clinic during 2016-2018 and responded to four lifestyle questions, constituted the Danish cohort (data retrieved from the national registry DANBIO). Patients with RA belonging to the BARFOT cohort, and who in a 2017 survey responded to four lifestyle questions, constituted the Swedish cohort. Lifestyle information was dichotomized as present tobacco use or not, BMI <25 kg/m² vs. ≥25 kg/m², alcohol overuse or not, and health enhancing physical activity (≥ 150 minutes/week) or less. The combined number of UL factors (0, 1, 2, 3, 4) were calculated. Crude logistic regression analyses were performed to determine the association between disease impact and two or more UL factors (controlled for age, gender and disease duration). Independent factors (disease impact) were pain (NRS 0-10, best to worst), fatigue (NRS 0-10, best to worst), function (HAQ, 0-3, best to worst) and quality of life (EQ-5D-3L 0-1, worst to best).

Results: The 566 included Danish patients had a mean age of 61.82 (SD 11.13) years, a disease duration of mean 12.40 (SD 10.95) years, and 72% were women. The 995 Swedish patients had a mean age of 66.38 (SD 12.90) years, a disease duration of mean 15.55 (SD 3.85) years, and 72% were women. 95% of the Danish patients and 82% of the Swedish patients reported at least one UL factor, while 66% and 47% respectively reported two or more (Figure 1). The most common ones were overweight/obesity and physical inactivity in both cohorts. Male gender OR 1.86 95% CI [1.21-2.85] and shorter disease duration OR 0.97 95% CI [0.95-0.99] were associated with two or more UL factors in the Danish cohort. In the Swedish cohort, male gender OR 1.42 95% CI [1.07 – 1.89], worse pain OR 1.10 95% CI [1.04 – 1.15], fatigue OR 1.09 95% CI [1.04 – 1.15], function OR 1.64 95% CI [1.28 – 2.10], and worse quality of life OR 0.35 95% CI [0.20 – 0.60] were associated with two or more UL factors.

Conclusion: Every other patient with RA had two or more UL factors in both the Danish and Swedish cohort, and more often they were men. The combined number of UL factors was not necessarily associated with disease impact. The findings are important for health professionals working with lifestyle interventions in patients with RA.

Figure 1. The combined number of unhealthy lifestyle (UL) factors in two Scandinavian RA cohorts.

REFERENCES:

- [1] Smolen JS, Landewé RBM, Bijlsma JWW, Burmester GR, Dougados M, Kerschbaumer A, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying anti-rheumatic drugs: 2019 update. *Ann Rheum Dis.* 2020;79(6):685-99.
- [2] Lindhardt J, Ahlehoj O, Gislason GH, Madsen OR, Olesen JB, Torp-Pedersen C, et al. The risk of myocardial infarction in rheumatoid arthritis and diabetes mellitus: a Danish nationwide cohort study. *Ann Rheum Dis.* 2011;70(6):929-34.

- [3] Agca R, Heslinga SC, Rollefstad S, Heslinga M, McInnes IB, Peters MJ, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis.* 2017;76(1):17-28.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1378

POS1424

PREGNANCY COURSE IN A COHORT OF WOMEN WITH MYOSITIS - RESULTS FROM REVNATUS

B. Jakobsen¹, H. Bjørngaard¹, H. S. Koksvik¹, T. T. Pedersen¹, M. Wallenius^{1,2}.

¹St. Olavs Hospital, Trondheim University Hospital, Norwegian National Advisory Unit on Pregnancy and Rheumatic Diseases, Trondheim, Norway; ²Norwegian University of Science and Technology, Department of Neuromedicine and Movement Science, Trondheim, Norway

Background: Myositis is a rare autoimmune disease characterized by proximal and symmetrical muscle weakness and inflammatory infiltrates on muscle biopsy, and may affect women in reproductive age. Only few reports exist on pregnancy outcomes in women with myositis. Well-controlled disease at time of conception seems to give better outcome for the mother and the fetus (1). Inflammatory active disease before and during pregnancy has been associated with preterm birth and infant small for gestational age (1).

Objectives: To describe the course of pregnancy in patients with myositis.

Methods: We prospectively collected data on pregnancy in women with myositis in the nationwide Norwegian quality registry - RevNatus. ICD-10 codes: ICD-10 M33.0 (5), M33.1 (2), M33.2 (1), M33.9 (3). Eleven pregnancies in ten women with a known pregnancy outcome from 2016 – 2020 were included.

Results: Median disease duration was 10 years (range 3-26), and median age at conception was 29 years (range 25-43). Five women had juvenile myositis. Two pregnancies ended in spontaneous abortions, one in 1st trimester and one in 2nd trimester. Nine pregnancies ended with live birth at median pregnancy week 39 (range 35-41) and a median birth weight of 3355 gram (range 2580-4750). None of the women experienced preeclampsia or HELLP. Three women were positive for ANA. Median CRP in second trimester was three (range 0-7). Among nine women with live births, seven were in remission, and two women had persistent disease in second trimester, defined by rheumatologist. Seven women used DMARDs in pregnancy, and two women used prednisolone 5mg throughout pregnancy.

Patient #	Pregnancy #	Maternal age at conception	Disease duration years	Medication during pregnancy	Disease activity 2 nd trimester	Gestational age at delivery	Infant birth weight (gram)	Delivery mode
A	1	29	25	none	Remission	38	3355	Standard vaginal delivery
B	2	34	7	azathioprine	Persistent	39	3725	Standard vaginal delivery
C	3	29	15	azathioprine				Spontaneous abortion week 7
D	4	35	3	ciclosporin prednisolone 5mg	Persistent	40	3000	Standard vaginal delivery
E	5	28	13	none	Remission	37	2765	Operational vaginal delivery
F	6	25	10	azathioprine hydroxy-chloroquine	Remission	41	4085	Standard vaginal delivery
F	7	27	13	azathioprine hydroxy-chloroquine prednisolone 5mg	Remission	41	3815	Standard vaginal delivery
G	8	36	13	prednisolone 5mg				Spontaneous abortion > week 12
H	9	29	9	azathioprine	Remission	35	2580	Standard vaginal delivery
I	10	43	5	azathioprine prednisolone 5mg	Remission	37	3140	Standard vaginal delivery
J	11	36	28	none	Remission	41	4750	Acute sectio

Conclusion: We did not observe disease flare during pregnancy and no serious adverse outcomes in the pregnancies ending with live births. However, data are sparse, and larger cohorts are necessary to evaluate possible risk factors.

References: 1. Munira S, Christopher-Stine L. Pregnancy in myositis and scleroderma. *Best Pract Res Clin Obstet Gynaecol*. 2020 Apr;64:59-67. doi: 10.1016/j.bpobgyn.2019.10.004. Epub 2019 Oct 18. PMID: 31928915.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1539

POS1425

ANTIBODIES TO PORPHYROMONAS GINGIVALIS ASSOCIATE WITH THE PRESENCE OF RHEUMATOID ARTHRITIS-RELATED AUTOANTIBODIES IN PATIENTS WITH PERIODONTITIS

C. De Vries¹, G. Ruacho¹, N. Sippl¹, B. A. Potempa², L. Ryden³, J. S. Potempa², B. Klinge⁴, E. Svenungsson¹, K. Lundberg¹. ¹Karolinska Institute, Department of Medicine, Division of Rheumatology, Stockholm, Sweden; ²University of Louisville, Department of Oral Immunology & Infectious Diseases, Louisville, United States of America; ³Nya Karolinska, Department of Medicine, Stockholm, Sweden; ⁴Huddinge Hospital, Department of Dental Medicine, Division of Oral Diseases, Huddinge, Sweden

Background: Epidemiologic studies have demonstrated a link between periodontitis (PD) and rheumatoid arthritis (RA), specifically RA characterized by anti-citrullinated protein antibodies (ACPA). The keystone pathogen driving PD, *Porphyromonas gingivalis* (Pg), is the only pathogen known to express peptidylarginine deiminase (PAD), a citrullinating enzyme. Hence, Pg has been proposed to be involved in triggering the ACPA response, by generating citrullinated antigens in an inflammatory milieu(1). Another major virulence factor of Pg is arginine gingipain B (RgpB), a proteinase which cleaves proteins so that PAD can access the site where citrullination takes place. We have previously shown elevated anti-RgpB IgG levels in ACPA+ RA patients, even before clinical onset(2, 3), and we hypothesize that anti-RgpB IgG could serve as a serological marker to identify PD patients with increased risk of developing ACPA+ RA.

Objectives: Based on this hypothesis, we set out to investigate whether anti-RgpB IgG was associated with PD, PD severity, autoimmunity in general, and the ACPA response in particular.

Methods: Anti-RgpB IgG, as well as RA- and systemic lupus erythematosus (SLE)-related autoantibodies targeting cyclic citrullinated peptide(s) (CCP2), rheumatoid factor (RF), dsDNA, cardiolipin, and β 2 glycoprotein, were measured by ELISA in serum samples from the ParoKrank study, which is a well-characterized cohort of 805 patients with a first myocardial infarction and 805 matched controls, where periodontal status has been determined by dentists(4). In this study, individuals with PD (n=941) were compared to individuals without PD (n=557).

Results: We detected significantly elevated ($p < 0.0001$) anti-RgpB IgG levels in PD compared to non-PD individuals, with highest levels recorded in severe PD. Anti-RgpB IgG levels were significantly increased in PD patients positive for CCP2 and/or RF (n=50), when compared to PD patients negative for CCP2 and RF (n=507), $p < 0.05$, and when compared to non-PD individuals positive for CCP2 and/or RF (n=62), $p < 0.05$. Notably, these differences were not seen for SLE-related autoantibodies. In addition, anti-RgpB IgG levels were significantly elevated amongst MI patients versus controls ($p < 0.05$), supporting the previous finding that PD is more common among MI patients(4).

Conclusion: Our data demonstrates a specific association between severe PD, elevated anti-RgpB IgG levels and RA-related autoantibodies, supporting a role for Pg in linking PD to ACPA+ RA. Further investigation will be needed to confirm whether anti-RgpB IgG can be used as a serological marker to identify PD patients with increased risk of developing ACPA+ RA.

REFERENCES:

- Rosenstein ED, Greenwald RA, Kushner LJ, Weissmann G. Hypothesis: the humoral immune response to oral bacteria provides a stimulus for the development of rheumatoid arthritis. *Inflammation*. 2004;28(6):311-8.
- Khramova N, Jiang X, Sherina N, Potempa B, Israelsson L, Quirke AM, et al. Antibodies to *Porphyromonas gingivalis* Indicate Interaction Between Oral Infection, Smoking, and Risk Genes in Rheumatoid Arthritis Etiology. *Arthritis Rheumatol*. 2016;68(3):604-13.
- Johansson L, Sherina N, Khramova N, Potempa B, Larsson B, Israelsson L, et al. Concentration of antibodies against *Porphyromonas gingivalis* is increased before the onset of symptoms of rheumatoid arthritis. *Arthritis Res Ther*. 2016;18(1):201.
- Rydén L, Buhlin K, Ekstrand E, Faire Ud, Gustafsson A, Holmer J, et al. Periodontitis Increases the Risk of a First Myocardial Infarction. *Circulation*. 2016;133(6):576-83.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1580

POS1426

PATIENTS WITH IMMUNE MEDIATED INFLAMMATORY DISEASES ARE OVERREPRESENTED IN LOW-FREQUENCY VIRAL SYMPTOM CLUSTERS

K. Tascilar^{1,2}, D. Simon^{1,2}, G. Krönke^{1,2}, A. Kleyer^{1,2}, A. Ramming^{1,2}, R. Atreya^{2,3}, M. Tenbusch⁴, K. Überla⁴, C. Berking^{2,5}, M. Sticherling^{2,5}, M. F. Neurath^{2,3}, G. Schett^{1,2}. ¹Friedrich Alexander University of Erlangen Nuremberg and Universitätsklinikum Erlangen, Department of Medicine-3, Rheumatology and Immunology, Erlangen, Germany; ²Friedrich Alexander University of Erlangen Nuremberg and Universitätsklinikum Erlangen, Deutsches Zentrum für Immuntherapie, Erlangen, Germany; ³Friedrich Alexander University of Erlangen Nuremberg and Universitätsklinikum Erlangen, Department of Medicine-1, Gastroenterology, Pneumology and Endocrinology, Erlangen, Germany; ⁴Friedrich Alexander University of Erlangen Nuremberg and Universitätsklinikum Erlangen, Institute of Clinical and Molecular Virology, Erlangen, Germany; ⁵Friedrich Alexander University of Erlangen Nuremberg and Universitätsklinikum Erlangen, Department of Dermatology, Erlangen, Germany

Background: Viral respiratory infections are common in the general population and result in a spectrum of outcomes ranging from effective viral clearance with no symptoms, to a maladaptive immune response that can result in severe symptomatic disease and death. Although patients with immune-mediated inflammatory diseases (IMID) are considered susceptible to poor outcomes from infectious syndromes, it is not known whether IMID patients are overall more prone to manifest common viral infection symptoms.

Objectives: To explore frequency patterns of common viral infection symptoms in IMID patients.

Methods: We previously recruited patients with IMIDs and individuals with no IMIDs for a seroprevalence study between February 1st and April 30th 2020 (1). Participants were questioned for the presence of eleven common viral disease symptoms. We clustered these data using an unsupervised binary data clustering algorithm (2) into 6 symptom clusters based on symptom frequency. Three major clusters (broadly symptomatic, intermediately symptomatic and oligo-/asymptomatic) and 2 sub-clusters (higher and lower frequency) for each major cluster. In addition, qualitative symptom clustering was done. We estimated standardized residuals to quantify the over/underrepresentation of IMID diagnosis frequencies in each subject cluster. We used Poisson regression to compare symptom counts by diagnosis.

Results: We analyzed 1909 participants (757 with IMIDs; 1152 non-IMID controls; Table 1). Within each major subject cluster (Figure 1A), IMID patients showed the highest positive deviation from the expected frequencies in lower frequency sub-clusters while non IMID controls showed the highest positive deviations in the higher frequency sub-clusters (Figure 1B). Inflammatory bowel disease and psoriasis were remarkably overrepresented in the lower frequency sub-cluster of the broadly-symptomatic cluster while RA was overrepresented in the lower frequency sub-clusters of intermediate and oligo-/asymptomatic clusters. X axis of Figure 1A presents qualitative symptom clusters. Regression analysis shows that RA patients among other IMIDs reported overall less symptoms (RR= 0.69, 95%CI, 0.58 - 0.80) compared to non-IMID controls.

Table 1. Participant characteristics and distribution of IMID diagnoses across subject clusters.

	Clusters						
	Broad Symptomatic		Intermediate Symptomatic		Oligo-Asymptomatic		
	Overall	Higher	Lower	Higher	Lower	Higher	Lower
N	1909	101	85	412	259	283	769
Age, years, mean (SD)	45.4 (15.2)	42.4 (13.3)	47.3 (15.2)	42.4 (12.9)	50.4 (15.5)	41.8 (14.9)	46.8 (15.9)
Male	1080 (56.6)	42 (41.6)	38 (44.7)	196 (47.6)	137 (52.9)	178 (62.9)	489 (63.6)
Diagnosis, n(%)							
No-IMID	1152 (60.3)	72 (71.3)	44 (51.8)	280 (68.0)	112 (43.2)	207 (73.1)	437 (56.8)
RA	226 (11.8)	7 (6.9)	5 (5.9)	29 (7.0)	56 (21.6)	17 (6.0)	112 (14.6)
IBD	178 (9.3)	5 (5.0)	15 (17.6)	46 (11.2)	29 (11.2)	19 (6.7)	64 (8.3)
SpA	142 (7.4)	7 (6.9)	5 (5.9)	23 (5.6)	25 (9.7)	14 (4.9)	68 (8.8)
Psoriasis	89 (4.7)	4 (4.0)	9 (10.6)	14 (3.4)	8 (3.1)	13 (4.6)	41 (5.3)
Other	122 (6.4)	6 (5.9)	7 (8.2)	20 (4.9)	29 (11.2)	13 (4.6)	47 (6.1)
Symptom count/ patient, mean (SD)	1.2 (1.7)	6.0 (1.3)	3.9 (1.1)	2.2 (1.0)	1.5 (0.6)	0.5 (0.5)	0.0 (0.0)

IBD, inflammatory bowel disease.