



OPEN ACCESS

CLINICAL SCIENCE

Efficacy and safety of emapalumab in macrophage activation syndrome

Fabrizio De Benedetti ,¹ Alexei A Grom ,^{2,3} Paul A Brogan ,⁴ Claudia Bracaglia ,¹ Manuela Pardeo,¹ Giulia Marucci,¹ Despina Eleftheriou,⁴ Charalampia Papadopoulou ,⁴ Grant S Schulert ,^{2,3} Pierre Quartier,^{5,6} Jordi Antón ,^{7,8} Christian Laveille,⁹ Rikke Frederiksen,¹⁰ Veronica Asnaghi,¹⁰ Maria Ballabio,¹⁰ Philippe Jacqmin,¹¹ Cristina de Min¹⁰

Handling editor Josef S Smolen

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ard-2022-223739>).

For numbered affiliations see end of article.

Correspondence to

Dr Fabrizio De Benedetti, Division of Rheumatology, Ospedale Pediatrico Bambino Gesù, IRCCS, Rome 00165, Italy; fabrizio.debenedetti@opbg.net

Part of the content in this manuscript has previously been presented at the American College of Rheumatology Convergence 2021, the 37th Annual Meeting of the Histiocyte Society, the 63rd American Society of Hematology Annual Meeting and Exposition, the European Alliance for Associations of Rheumatology 2022 Congress, the American Society of Pediatric Hematology and Oncology Conference 2022 and the Paediatric Rheumatology European Society 2022 Congress.

Received 18 December 2022

Accepted 2 February 2023

Published Online First

31 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: De Benedetti F, Grom AA, Brogan PA, et al. *Ann Rheum Dis* 2023;**82**:857–865.

ABSTRACT

Objectives Macrophage activation syndrome (MAS) is a severe, life-threatening complication of systemic juvenile idiopathic arthritis (sJIA) and adult-onset Still's disease (AOSD). The objective of this study was to confirm the adequacy of an emapalumab dosing regimen in relation to interferon- γ (IFN γ) activity by assessing efficacy and safety. The efficacy outcome was MAS remission by week 8, based on clinical and laboratory criteria.

Methods We studied emapalumab, a human anti-IFN γ antibody, administered with background glucocorticoids, in a prospective single-arm trial involving patients who had MAS secondary to sJIA or AOSD and had previously failed high-dose glucocorticoids, with or without anakinra and/or ciclosporin. The study foresaw 4-week treatment that could be shortened or prolonged based on investigator's assessment of response. Patients entered a long-term (12 months) follow-up study.

Results Fourteen patients received emapalumab. All patients completed the trial, entered the long-term follow-up and were alive at the end of follow-up. The investigated dosing regimen, based on an initial loading dose followed by maintenance doses, was appropriate, as shown by rapid neutralisation of IFN γ activity, demonstrated by a prompt decrease in serum C-X-C motif chemokine ligand 9 (CXCL9) levels. By week 8, MAS remission was achieved in 13 of the 14 patients at a median time of 25 days. Viral infections and positive viral tests were observed.

Conclusions Neutralisation of IFN γ with emapalumab was efficacious in inducing remission of MAS secondary to sJIA or AOSD in patients who had failed high-dose glucocorticoids. Screening for viral infections should be performed, particularly for cytomegalovirus.

Trial registration number NCT02069899 and NCT03311854.

INTRODUCTION

Macrophage activation syndrome (MAS) is a form of secondary haemophagocytic lymphohistiocytosis (HLH) occurring as a life-threatening complication of rheumatic diseases.¹ It is most frequent in systemic juvenile idiopathic arthritis (sJIA) and adult-onset Still's disease (AOSD), affecting about 10%–20% of patients.^{1–4} sJIA and AOSD are considered the same disease named differently depending only on age at

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Macrophage activation syndrome (MAS) is a severe and potentially life-threatening complication of systemic juvenile idiopathic arthritis and adult-onset Still's disease.
- ⇒ There are no therapeutic options that have been prospectively investigated in MAS.
- ⇒ Data in animal models and ex vivo data from humans with MAS led to the hypothesis that interferon- γ (IFN γ) has a pathogenic role in MAS.

WHAT THIS STUDY ADDS

- ⇒ This open-label multicentre trial using emapalumab, an anti-IFN γ antibody, in patients who have failed to respond to high-dose glucocorticoids, demonstrates that IFN γ has a pathogenic role in MAS and that its neutralisation markedly improved MAS.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ The results of this study show that neutralisation of IFN γ with emapalumab is a therapeutic option for patients with severe MAS who have failed standard of care with high-dose glucocorticoids.

onset below or above 16 years, respectively.^{2,3,5} The incidence and features of MAS are similar in the two age groups.^{2,5} We use the term sJIA/AOSD to collectively identify these patients.

Similar to other forms of HLH, MAS is caused by excessive activation and expansion of T lymphocytes and macrophages resulting in hyperinflammation.^{1,6,7} MAS is characterised by fever, hepatosplenomegaly, cytopenias, liver dysfunction, coagulation abnormalities and hyperferritinaemia, and may progress to multiple organ failure, with mortality rates of 10%–20%.⁴

MAS is treated with high-dose glucocorticoids with satisfactory response in two-thirds of the patients.^{4,8} In patients unresponsive to glucocorticoids, ciclosporin is usually added. Treatment with cyclophosphamide, etoposide, intravenous immunoglobulin, etanercept, anakinra, tocilizumab, JAK inhibitors and plasmapheresis, have been described

Treatment

in case reports or small series.^{4 8–12} None of these regimens have been prospectively investigated.

Overproduction of interferon- γ (IFN γ) is present and pathogenic in animal models of MAS.^{13–16} In sJIA/AOSD, high IFN γ activity demonstrated by high serum levels of C-X-C motif chemokine ligand 9 (CXCL9), a chemokine selectively induced by IFN γ ,¹⁷ is associated with MAS onset and severity.^{18–22}

We studied emapalumab, a fully human anti-IFN γ antibody, in patients with MAS secondary to sJIA/AOSD who failed high-dose glucocorticoids.

METHODS

This phase II, open-label, single-arm trial (NI-0501-06; ClinicalTrials.gov Identifier NCT03311854), conducted at five sites in Italy, France, Spain, the UK and the USA, comprised screening, a treatment period of 28 days and a short-term 4-week follow-up. Data were collected up to 12 months in a long-term follow-up study (NI-0501-05; ClinicalTrials.gov Identifier NCT02069899) (online supplemental figure S1 in the online supplemental appendix). Results from both studies are reported.

FDB, AG, MB and CdM designed the study. The sponsor was also responsible for data collection, management and analysis. All authors vouch for the accuracy and completeness of the data and analyses, and for the fidelity of the study to the protocol.

Patient involvement

The study was presented and discussed at the 2016 and 2017 meetings of the sJIA Foundation in Washington, DC, USA. Information on the study for patients and families was published on the sJIA Foundation website. Dissemination of the data gathered from the study is planned to be discussed with the sJIA Foundation.

Patients

Patients had sJIA based on International League Against Rheumatism criteria,²³ or AOSD based on Yamaguchi criteria.²⁴ For patients presenting with MAS at sJIA onset, presumption of sJIA was based on Childhood Arthritis and Rheumatology Research Alliance criteria.²⁵ Patients had MAS, according to American College of Rheumatology/EULAR criteria,²⁶ and an inadequate response to high-dose intravenous glucocorticoids according to the treating physicians. High-dose glucocorticoids were defined as ≥ 2 mg/kg/day of prednisone equivalent in two divided doses, or at least 60 mg/day in patients weighing 30 kg or more, including but not limited to pulses up to 30 mg/kg/day for at least 3 consecutive days. Patients with active infections potentially favoured by IFN γ neutralisation (typical and atypical mycobacteria, *Histoplasma capsulatum*, *Shigella*, *Salmonella*, *Campylobacter* and *Leishmania*) were excluded (see online supplemental appendix 1). Eligibility criteria are listed in online supplemental table S1.

Treatment

Emapalumab infusions were administered on a background of glucocorticoids. The initial dose of emapalumab was 6 mg/kg on day 0, followed by 3 mg/kg every 3 days until day 15 and twice weekly until day 28. Treatment with emapalumab could be stopped on investigator's assessment of remission, but not before three emapalumab doses have been administered. Frequency between infusions could be shortened, dose could be increased or treatment prolonged on investigator's assessment of unsatisfactory response. Ciclosporin could be continued, if started at least 3 days before initiating emapalumab. Interleukin (IL)-1

and IL-6 inhibitors were not allowed. An amendment allowed continuation of anakinra, if started at least 3 days before initiating emapalumab, and its introduction during the study, at a maximum dose of 4 mg/kg/day to treat the underlying sJIA/AOSD. Tocilizumab and canakinumab were not allowed during the trial. Prophylaxis against herpes zoster was administered as per local standards. Glucocorticoid tapering could be initiated as soon as the patients' conditions allowed based on investigator's assessment.

Outcomes

The objective was to confirm the adequacy of the emapalumab dosing regimen in relation to IFN γ activity by assessing efficacy and safety. Serum levels of emapalumab, total (free and emapalumab-bound) IFN γ , CXCL9 and soluble IL-2 receptor were measured (online supplemental appendices 2 and 3).

The efficacy outcome was MAS remission by week 8, defined as resolution of clinical signs and symptoms (online supplemental table S2) according to the physician global assessment (visual analogue scale $\leq 1/10$) and white blood cell and platelet count above lower limit of normal, lactate dehydrogenase (LDH), alanine aminotransferase and aspartate aminotransferase below 1.5 times upper limit of normal (ULN), fibrinogen > 100 mg/dL and ferritin levels²⁷ decreased by at least 80% or below 2000 ng/mL, whichever was lower. Other efficacy evaluations included glucocorticoid dose (expressed as mg/kg/day of prednisone-equivalent) and survival. Adverse events (AEs) were assessed. During the long-term follow-up, evaluations included MAS episodes, AEs, pharmacokinetics and pharmacodynamics.

Statistical analysis

The analysis population included all patients. Categorical variables are presented with the number and percentage within each category. Continuous variables are reported as median (range). Laboratory parameters of MAS were measured locally.

RESULTS

Study population

Fourteen patients received emapalumab. All completed the study and entered long-term follow-up. Thirteen patients had sJIA onset before 16 years, and one AOSD with onset at 16 years and 9 months. Four patients had presumption of sJIA, which was later confirmed. Six patients had a total of 19 previous MAS episodes in their history, treated with high-dose glucocorticoids, with the addition of anakinra and/or ciclosporin in approximately half.

During the week preceding emapalumab, all patients were receiving high-dose intravenous glucocorticoids, as per protocol. In addition, eight were receiving ciclosporin and seven anakinra (table 1). Of the patients treated with anakinra, three were receiving standard doses for sJIA (≤ 4.0 mg/kg/day), and four high doses ranging from 7.5 to 15 mg/kg/day. Patients had severe MAS at baseline, shown by high physician assessment of MAS activity and marked abnormalities of laboratory parameters (table 2). Notably, worsening or no improvement of laboratory parameters was observed between screening and baseline, despite ongoing treatment.

Of the 14 patients, 6 received emapalumab up to day 28. In seven patients who had MAS remission per investigator's assessment, it was possible to discontinue emapalumab earlier. One patient continued emapalumab up to day 39. The median treatment duration was 27 days (range, 7–39) with the number of infusions ranging from 3 to 17 per patient.

Table 1 Baseline demographics and clinical characteristics of the patients

Demographic	(n=14)
Age, years, median (range)	11.0 (2–25)
Sex, female, n (%)	10 (71.4)
Weight, kg, median (range)	45.9 (12.0–68.8)
Age at diagnosis of sJIA/AOSD, years, median (range)	10.5 (1–17)
Previous MAS episodes	
Patients with previous MAS episodes, n (%)	6 (43%)
Total number of previous MAS episodes	19
Number of MAS episodes per patient*	3 (1–6)
Treatment of the current MAS episode prior to emapalumab	
High-dose intravenous glucocorticoids, n (%)	14 (100)
Average daily dose during week –1,† mg/kg prednisone-equivalent, median (range)	15.7 (0.8–36.4)
Ciclosporin,‡ n (%)	8 (57.1%)
Anakinra,‡ n, (%)	7§ (50)
Average daily dose during week –1,† mg/kg, median (range)	4.2 (1.6–13.9)
IVIg, n (%)	3 (21.4)

*Patients with no previous MAS episode(s) are excluded from the calculation of the median (range).
†Week prior to emapalumab administration.
‡Five patients received anakinra and ciclosporin, concomitantly.
§One additional patient received anakinra until 11 days prior to start of emapalumab administration.
AOSD, adult-onset Still's disease; IVIg, intravenous immunoglobulin; MAS, macrophage activation syndrome; sJIA, systemic juvenile idiopathic arthritis.

Pharmacokinetics and pharmacodynamics

All patients received an initial dose of emapalumab of 6 mg/kg followed by doses of 3 mg/kg. The initial dose allowed to rapidly reach serum concentrations of emapalumab close to the steady-state concentrations obtained with the doses of 3 mg/kg (figure 1A). The dosing interval was transiently shortened from 3 to 2 days in three patients based on clinical and laboratory parameters suggestive of incomplete response. In two of these patients the dosing interval was shortened for several infusions. In these two patients, measurement

of emapalumab concentrations showed rapid drug clearance consistent with target-mediated drug disposition associated with high levels of total IFN γ (online supplemental figure S2).²⁸

At baseline, CXCL9 concentrations were markedly elevated in all patients, indicating high IFN γ activity (figure 1B). Total IFN γ concentrations were variable across patients (figure 1C). Total IFN γ concentrations at day 3 (at equilibrium between free and emapalumab-bound IFN γ) reflect IFN γ production at baseline.^{27 28} There was no correlation between total IFN γ at day 3 and CXCL9 levels at baseline (not shown). Soluble IL-2 receptor levels at baseline were markedly elevated in all patients (figure 1D).

Emapalumab administration led to a rapid decrease in serum CXCL9 levels, indicating neutralisation of IFN γ activity and therefore the appropriateness of the selected dosing regimen. Patients with elevated total IFN γ at day 3 also showed a decrease in total IFN γ concentrations over time indicating decreasing IFN γ production over time. Soluble IL-2 receptor concentrations also decreased markedly (figure 1D), indicating decreasing T cell activation. During long-term follow-up, in the absence of high IFN γ production, as shown by low levels of total IFN γ , emapalumab showed a slow linear terminal elimination phase with a half-life of 24 days, like that in healthy subjects (online supplemental figure S3).²⁸ Total IFN γ , CXCL9 and soluble IL-2 receptor levels were close to, or within, the normal range (online supplemental table S3).

Efficacy

During emapalumab administration, physician global assessment of MAS activity and MAS laboratory parameters rapidly improved in all patients while glucocorticoids were tapered.

By week 8, 13 patients (93%) achieved MAS remission at a median time of 25 days after emapalumab initiation, the earliest at day 9 (figure 2). At week 8, 2 of the 13 patients who had previously achieved MAS remission did not meet the criteria of MAS remission because of a single laboratory abnormality (one with LDH 1.7-fold above the ULN; one with white blood cells at $4.8 \times 10^9/L$ with lower limit of normal at $5.5 \times 10^9/L$). One

Table 2 MAS activity, as defined by physician's assessment, and laboratory features of MAS before, during and after emapalumab treatment

	Screening (n=14)	Baseline (n=14)	Week 2 (n=14)	Week 4 (n=14)	Week 8 (n=14)
Physician assessment of MAS activity*, cm, median (range)	8.3 (2.0–10.0)	8.0 (2.0–10.0)	2.3 (0.0–9.5)	0.3 (0.0–6.5)	0 (0.0–1.0)
Ferritin, ng/mL, median (range)	19 865 (367–192 584)	25 709 (716–192 584)	1410 (65–29 099)	180 (22–18 430)	55 (4–561)
Increased, n (%)†	13 (93)	14 (100)	9 (64)	2 (14)	0 (0)
WBCs, $10^9/L$, median (range)	5.8 (1.0–25.7)	5.1 (0.9–25.7)	8.4 (0.9–32.6)	12.3 (2.4–34.7)	9.5 (4.8–22.6)
Decreased, n (%)‡	6 (43)	7 (50)	4 (29)	1 (7)	1 (7)
Platelets, $10^9/L$, median (range)	152 (48–546)	112 (52–558)	284 (55–590)	336 (104–591)	383 (214–915)
Decreased, n (%)‡	6 (43)	9 (64)	4 (29)	2 (14)	0 (0)
Aspartate aminotransferase, U/L, median (range)	244 (26–3814)	190 (19–2039)	34 (5–146)	29 (16–100)	28 (15–52)
Increased, n (%)§	12 (86)	10 (71)	3 (21)	1 (7)	0 (0)
Alanine aminotransferase, U/L, median (range)	190 (139–3179)	302 (70–1492)	61 (18–327)	45 (23–195)	28 (7–59)
Increased, n (%)§	14 (100)	14 (100)	6 (43)	5 (35.7)	0 (0)
LDH, U/L, median (range)	1699 (859–12 734)	1502 (588–12 734)	518 (283–1572)	592 (277–810)	497 (306–741)
Increased, n (%)§	13 (100)	13 (93)	6 (43)	4 (29)	2 (14)
Fibrinogen, mg/dL, median (range)	230 (137–341)	174 (70–465)	244 (145–515)	279 (200–460)	337 (190–460)
Decreased, n (%)¶	0 (0)	1 (7)	0 (0)	0 (0)	0 (0)

*MAS activity was based on a 10 cm visual analogue scale, with scores ranging from 0 to 10 and higher score indicating higher activity.

†Increased ferritin was defined as ferritin >684 ng/mL as per 2016 Classification Criteria for MAS complicating sJIA.²³

‡Decreased white blood cell and platelet were defined as counts below the lower limit of normal for age of local laboratory.

§Increased alanine aminotransferase, aspartate aminotransferase and LDH were defined as values above 1.5 the upper limit of normal range of local laboratory.

¶Decreased fibrinogen was defined as fibrinogen <100 mg/dL.

LDH, lactate dehydrogenase; MAS, macrophage activation syndrome; WBC, white blood cells.




Figure 1 Pharmacokinetic (emapalumab* (A)) and pharmacodynamic (CXCL9† (B), total IFN γ ‡ (C) and sIL-2R§ (D)) parameters in patients with MAS treated with emapalumab. *Serum emapalumab concentrations in patients treated with emapalumab (initial dose of 6 mg/kg, followed by intended maintenance dosing of 3 mg/kg every 3 days until day 15, and twice weekly until day 28). †Serum concentrations of CXCL9, a biomarker of IFN γ activity, being, at baseline, 4–362 times the 95th percentile of healthy volunteers (271 pg/mL),²⁴ indicated by the horizontal dotted line. ‡Total IFN γ (free and emapalumab-bound) that reflects IFN γ production, ranging, at day 3, from normal to 12 times the 95th percentile of healthy volunteers receiving emapalumab (2084 pg/mL) indicated by the horizontal dotted line. §Serum sIL-2R, a biomarker of T cell activation, being, at baseline, 1.6–20 times the 95th percentile of healthy volunteers (1071 pg/mL, unpublished data) indicated by the horizontal dotted line. CXCL9, C-X-C motif chemokine ligand 9; IFN γ , interferon- γ ; MAS, macrophage activation syndrome; sIL-2R, soluble interleukin-2 receptor.

patient who stopped emapalumab after three doses on investigator's assessment of remission never met the criteria of MAS remission only because of LDH levels 1.5-fold above the ULN.

All laboratory parameters of MAS rapidly improved on initiation of emapalumab with evident improvement at week 1 and with all parameters being normal in the majority of patients by week 4 (figure 3, table 2 and online supplemental figure S4).

During long-term follow-up, 13 patients did not have MAS episodes. One patient had a MAS episode 11 months after stopping emapalumab, when emapalumab was undetectable in serum. At the end of the long-term follow-up, 10 patients met the MAS remission criteria. Four did not meet these criteria because of either mild abnormality in one laboratory value ($n=1$), MAS activity at 1.5 cm ($n=1$), missing data ($n=1$) or absence of data as the patient missed the 12-month visit (online supplemental table S4).

Tapering of glucocorticoids occurred rapidly after emapalumab initiation. During the week preceding emapalumab, the median average daily dose was 15.7 mg/kg/day prednisone-equivalent, 2.3 mg/kg during week 2 and 0.56 mg/kg during week 8 (figure 4). At the end of the long-term follow-up, five patients were not receiving glucocorticoids and six were receiving <0.3 mg/kg/day prednisone-equivalent. Two patients were receiving between 1 and 2 mg/kg; the above-mentioned patient with a MAS episode and one patient with lung disease associated with sJIA. Data on glucocorticoid dose for one patient at the last follow-up visit are not available, as the patient missed the visit.

Eight patients were receiving ciclosporin at baseline. Ciclosporin was discontinued in two patients early after emapalumab initiation (day 4 and 10) and in four additional patients during long-term follow-up.

During the trial, anakinra was continued in four patients at a dose of ≤ 4 mg/kg, the standard dose for sJIA/AOSD treatment and in one patient at a dose of 7.5 mg/kg. These patients did not present with flares of the underlying sJIA/AOSD. During the trial, while MAS was improving, six flares of sJIA/AOSD were observed: three in the three patients who discontinued anakinra before emapalumab initiation and three in three patients who had not previously received and were not receiving anakinra. During long-term follow-up, four sJIA/AOSD flares occurred in four patients. sJIA/AOSD flares and the background treatment at time of the flares are described in online supplemental table S5.

Safety

No deaths were reported during the trial and the long-term follow-up. During the trial, after initiation of emapalumab, 88 AEs were reported in 13 patients (table 3 and online supplemental table S6). All events were mild or moderate in intensity except two (one cardiopulmonary failure, and one neutropenia; neither related to emapalumab). The most frequently reported AEs were infections and positive tests for infectious agents in the absence of clinical symptoms. All infectious events were of viral origin. No bacterial or opportunistic infections were reported. Six viral events in three patients (two infections and four positive




Figure 2 Time to MAS remission*. *Time to MAS remission defined as resolution of clinical signs and symptoms according to the investigator (visual analogue scale $\leq 1/10$) (online supplemental table S2) and resolution of the abnormalities of MAS laboratory parameters: white blood cell and platelet count above lower limit of normal, LDH below 1.5 times ULN, alanine aminotransferase/aspartate aminotransferase below 1.5 times ULN, fibrinogen >100 mg/dL and ferritin levels decreased by at least 80% or below 2000 ng/mL, whichever was lower. The continuous line represents the proportion of patients with MAS remission, the dotted line represents the 95% CI. The cross indicates the censored patient. LDH, lactate dehydrogenase; MAS, macrophage activation syndrome; ULN, upper limit of normal.

tests) were reported as related to emapalumab. One cytomegalovirus (CMV) reactivation was reported as serious. In total, there were five CMV events (three reactivations, one infection and one positive test with no symptoms). All viral events resolved spontaneously or with standard treatment. Two infusion-related reactions (pruritic rash), not reported as severe, occurred during a total of 128 infusions. The rate of AEs and of infectious events was not increased during concomitant treatment with anakinra and emapalumab compared with treatment with emapalumab alone (table 4).

During long-term follow-up, 37 AEs were reported in 12 patients. Three were serious: one flare of sJIA (see online supplemental table S5), one oedema of the ankle and one MAS episode (that, as mentioned above, occurred 11 months after stopping emapalumab, when emapalumab was undetectable). Ten infectious events occurred in seven patients; four when emapalumab levels were measurable, and six after emapalumab levels became undetectable. All reported infections were viral, except for one cestode infection.

DISCUSSION

We show that neutralisation of IFN γ with emapalumab was efficacious in inducing MAS remission in patients with MAS secondary to

sJIA/AOSD who failed standard of care including high-dose glucocorticoids with or without anakinra and/or ciclosporin.

The dosing regimen of emapalumab was chosen based on the data gathered in primary HLH patients on (a) the production rate of IFN γ (through the assessment of total IFN γ), (b) the rapid clearance of emapalumab consequent to target-mediated drug disposition in the presence of high IFN γ production and (c) the concentration of emapalumab required to neutralise high IFN γ levels. Through modelling and simulation, on the basis of the high levels of CXCL9 present in patients with MAS, it was possible to predict the dose of emapalumab required to achieve IFN γ neutralisation. A dosing regimen with an initial dose of 6 mg/kg, followed by 3 mg/kg every 3 days maintenance doses, was selected to achieve rapid efficacy as MAS usually has acute onset and may worsen rapidly, becoming life-threatening. In the majority of patients, the chosen regimen rapidly achieved IFN γ neutralisation, shown by prompt decrease in serum CXCL9 levels and prompt clinical and laboratory response. Shortening of the dosing interval might be considered in patients with unsatisfactory response. Indeed, in two patients with initial unsatisfactory response, shortening of the dosing interval led to achievement of response. In these patients, we found high IFN γ production and low emapalumab concentrations due




Figure 3 Changes in ferritin* (A), aspartate aminotransferase† (B), platelets (C) and LDH‡ (D) levels over time. *Panel A insert shows in detail changes from baseline to week 4 for patients with baseline levels of ferritin <50 000 ng/mL. †Panel B insert shows in detail changes from baseline to week 4 for patients with baseline levels of aspartate aminotransferase below 300 U/L. ‡Panel D insert shows in detail changes from baseline to week 4 for patients with baseline levels of LDH below 2500 U/L. LDH, lactate dehydrogenase.

to rapid drug clearance consistent with target-mediated drug disposition.

The patients recruited in this study had markedly elevated CXCL9 levels, reflecting high IFN γ activity. This finding is consistent with previous observations in patients with MAS.^{16 18} In contrast with primary HLH patients treated with emapalumab,²⁷ we did not find a correlation of CXCL9 levels at baseline with total IFN γ levels at day 3, that, as mentioned, reflect IFN γ production. Therefore, in MAS, the pathogenic role of IFN γ may be due to increased IFN γ production and increased sensitivity to IFN γ . Indeed, monocytes from patients with MAS have increased responsiveness to IFN γ , possibly through increased expression of tripartite motif containing protein 8 (TRIM-8), which potentiates response to IFN γ .^{29 30}

In this trial, the first prospectively investigating a treatment for MAS, we used a novel clinically meaningful efficacy measure, namely MAS remission, that combines resolution of clinical signs and symptoms and of the abnormalities in MAS laboratory parameters. Incidentally, the ferritin threshold (decrease by at least 80% or be below 2000 ng/mL, whichever was lower), previously used in the primary HLH trial with emapalumab,²⁷ is justified by the often-needed multiple transfusions and the frequent fluctuations of ferritin >1000 ng/mL in patients with HLH/MAS with good response to treatment. Notably, the highest value for ferritin at week 8 in our trial was 561 ng/mL.

After emapalumab initiation, MAS improved rapidly with median time to MAS remission of 25 days. This improvement occurred while background glucocorticoids were rapidly tapered: during the second week of emapalumab treatment, the median glucocorticoid dose was already 85% lower compared with that administered during the week preceding initiation of emapalumab. In general, patients seemed to benefit from emapalumab even when the MAS remission criteria were partially achieved. Indeed, in the few patients who did not reach MAS remission, the physician assessment of disease activity was ≤ 1 and MAS remission was not achieved because of minor abnormalities in only one of the laboratory parameters.

Elevated production of IFN γ is a feature of animal models of primary HLH and of secondary HLH, including infection-associated HLH and MAS.¹³ In these models, when the effect of therapeutic IFN γ neutralisation was tested, benefit was always observed with prevention of death and/or disease improvement. The results of this trial, together with the efficacy of emapalumab demonstrated in primary HLH,²⁷ and the anecdotal cases of patients with different forms of secondary HLH successfully treated with emapalumab,^{31–37} suggest that, in humans, IFN γ is an important driver of MAS/HLH, independently of the trigger or the underlying predisposing condition.

Notably, some inflammatory flares of the underlying sJIA/AOSD were observed, suggesting that, in the context of sJIA/




Figure 4 Changes in glucocorticoid dose* over time†. *The dose of glucocorticoids is shown as average daily dose in each week of the study, starting from the week preceding emapalumab treatment (week -1). The dose is shown in mg/kg of prednisone equivalent. †Weeks are defined starting from week -1 (from day -6 to day 0), followed by week 1 (from day 1 to day 7); week 2 (from day 8 to day 14); week 3 (from day 15 to day 21); up to week 8 (from day 50 to day 56).

AOSD, increased IFN γ activity is a feature of the hyperinflammation of MAS, but not of the autoinflammation that maintains sJIA. From a clinical perspective, we observed six flares of the underlying sJIA/AOSD in six out of the nine patients who were not receiving anakinra for the treatment of sJIA while receiving emapalumab. In contrast, no sJIA flares were observed in the five patients who continued anakinra after emapalumab initiation. Importantly, during the trial, no increase in AEs and/or infectious AEs was observed in patients concomitantly exposed to emapalumab and anakinra. Therefore, continuation of anakinra for the underlying sJIA/AOSD, at the doses conventionally used and allowed in this trial, should be considered.^{38–40}

The contribution of the concomitant anakinra to the achievement of MAS remission with emapalumab is to be considered negligible. Only one patient continued anakinra at a dose >4 mg/kg/day, a so-called higher dose that has been reported as potentially efficacious in MAS.^{10 14 41} Of note, several patients recruited to our study received anakinra before treatment with emapalumab and, in these patients, anakinra, even at doses >4 mg/kg/day, did not prevent or improve MAS. Furthermore, of the three patients in the trial who did not achieve MAS remission at week 8, two were receiving concomitant anakinra.

IFN γ neutralisation might increase predisposition to selected infections, as inferred based on data in humans with defective IFN γ activity, that is, subjects carrying autoantibodies to IFN γ or with IFN γ receptor deficiency.⁴² While herpes zoster infections are common in individuals with defective IFN γ activity, no cases were reported in this trial. Notably, per-protocol, patients received prophylaxis with acyclovir. We did not observe any infections from typical or atypical mycobacteria, *H. capsulatum*, *Shigella*, *Salmonella*, *Campylobacter* or *Leishmania*, known to occur more frequently in subjects with defective IFN γ activity. Noteworthy, patients with infections from these agents were excluded from the trial. No bacterial or opportunistic

Table 3 AEs during the treatment with emapalumab and during the long-term follow-up

Event category	NI-0501-06 study (up to week 8) (n=14)	Long-term follow-up (up to week 52) (n=14)
AEs		
Events, n	88	37
Patients with at least one event, n (%)	13 (93)	12 (86)
	Emapalumab levels at or above LLOQ	Emapalumab levels below LLOQ
Serious AEs		
Events, n	9*	2
Patients with at least one event, n (%)	6 (43)	2 (14)
Cardiopulmonary failure†	1	0
sJIA flare‡	1	1
CMV infection reactivation§	1	0
Intracardiac thrombus	1	0
Pericarditis	1	0
Thrombocytopenia	1	0
Pneumotosis intestinalis	1	0
Juvenile myoclonic epilepsy	1	0
Rash	1	0
Oedema of the ankle region	0	1
MAS¶	0	0
Events related to study drug		
Events, n	9	0
Patients with at least one event, n (%)	4 (28)	0
CMV infection reactivation§	1	0
Viral infection	1	0
Viral URTI	0	0
Viral test positive	4	0
Infusion-related reaction	1	0
Rash pruritic	1	0
Diarrhoea	1	0
Infections		
Events, n	10	4
Patients with at least one event, n (%)	6 (43)	2 (14)
CMV infection reactivation	3	0
CMV infection	1	0
Epstein-Barr infection	0	0
Rhinovirus infection	0	1
Enterovirus infection	0	0
Viral infection	1	0
Viral URTI	0	3
Nasopharyngitis	1	0
Viral test positive**	4	0
Cestode infection	0	0
Systemic juvenile idiopathic arthritis flares		
Events, n	1	3
Patients with at least one event, n (%)	1 (7)	3 (21)
Infusion-related reactions to emapalumab		
Events, n	2	Not applicable
Patients with at least one event, n (%)	2 (14)	Not applicable
*Of the nine serious AEs reported during the trial, one was severe in intensity (cardiopulmonary failure) and the others were of moderate intensity. One of these events was reported as related to emapalumab. All nine events resolved.		
†In the context of rapidly worsening MAS, the patient experienced worsening of pre-existing cardiocirculatory instability a few hours after the first emapalumab infusion. Due to this deterioration, the patient was transferred to the ICU and required intensification of inotropic support (dopamine dose increase). Of note, emapalumab dosing interval was shortened (see online supplemental figure S2) until MAS gradually improved and the patient recovered from the event.		
‡Table 3 lists only the episodes of sJIA flare reported as a serious AE. Of note, sJIA flares were identified based on AE reporting as well as based on the rationale for change in medications or medication dose provided by the investigator and these are described in online supplemental table S5.		
§This event is the same event reported in the Safety section of the Results and is reported in table 3 under serious AEs, under related AEs and is one of the three CMV reactivations reported under infections.		
¶This event is the same event reported in the Efficacy section of the Results.		
**One each for CMV, adenovirus, BK polyoma virus, reactivations.		
AE, adverse event; CMV, cytomegalovirus; ICU, intensive care unit; LLOQ, lower limit of quantification; MAS, macrophage activation syndrome; sJIA, systemic juvenile idiopathic arthritis; URTI, upper respiratory tract infection.		

Table 4 Rate of AEs* and infectious AEs during the trial while exposed to emapalumab alone or emapalumab and anakinra

	Emapalumab	Emapalumab and anakinra
Exposure, days at risk	303	506
AEs		
Events, n	45	43
Rate per 100 patient-days	14.9	8.5
Infectious AEs		
Events, n	5	5
Rate of per 100 patient-days	1.7	1

*One event of flare of sJIA and two infusion-related reactions to emapalumab were excluded from this analysis.
AE, adverse event; sJIA, systemic juvenile idiopathic arthritis.

infections were reported. Viral infections or positive viral tests were reported in six patients during the trial and in two during follow-up while emapalumab was detectable. All events resolved spontaneously or with standard treatment. Of note, screening for Epstein-Barr virus, CMV and adenovirus every 2 weeks was required by protocol during the trial. Despite the fact that CMV is not reported at increased frequency in humans with defective IFN γ activity,⁴² it should be noted that five events were related to CMV, with four reported as reactivations or infections and one as test positive. Whether concomitant prolonged immunosuppression with high-dose glucocorticoids in critically ill patients might have contributed to this observation remains to be established. Despite CMV infections not being reported in humans with defective IFN γ activity,⁴² based on our data, patients with MAS receiving emapalumab should be screened for the presence of CMV. Although the entire age spectrum of AOSD is not covered, with only one patient above 20 years of age, it is highly unlikely that the infection risk changes with age.

The main limitation of this study is the single-arm nature of the design. However, a randomised controlled study in this patient population does not appear to be possible and, notably, ethical. This is based, first, on the absence of a standardised and/or validated treatment that has been prospectively investigated in patients with MAS who have failed high-dose glucocorticoids. Therefore, no drug is available as comparator for a head-to-head design. Second, the risk of death associated with this condition makes it unethical to continue, with no additional therapy, an ineffective treatment (ie, high-dose glucocorticoids) in patients who have already failed this treatment. Therefore, the option of continuing high-dose glucocorticoids plus placebo as a comparator arm is also not acceptable.

In conclusion, we demonstrate that IFN γ is an important driver of MAS secondary to sJIA/AOSD and that its neutralisation with emapalumab leads to remission of MAS in patients who failed high-dose glucocorticoids. Attention should be paid to viral infections, particularly to CMV; periodic screening for viral infections should be performed.

Author affiliations

¹Division of Rheumatology, Ospedale Pediatrico Bambino Gesù, IRCCS, Rome, Italy

²Division of Rheumatology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

³Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

⁴University College London Great Ormond Street Institute of Child Health, London, UK

⁵Pediatric Immuno-Hematology and Rheumatology Unit, RAISE Rare Disease Reference Centre, Hôpital Universitaire Necker-Enfants Malades, Assistance Publique-Hôpitaux de Paris, Paris, France

⁶Université Paris-Cité, Paris, France

⁷Pediatric Rheumatology, Hospital Sant Joan de Deu, Barcelona, Spain

⁸Faculty of Medicine, Universitat de Barcelona, Barcelona, Spain

⁹Calvagone Sarl, Liègues, France

¹⁰Swedish Orphan Biovitrum AG (Sobi), Basel, Switzerland

¹¹MnS Modelling and Simulation, Dinant, Belgium

Twitter Claudia Bracaglia @claudiabrac, Grant S Schultert @GrantSchultert and Jordi Antón @AntonJordi68

Acknowledgements The authors wish to acknowledge the contribution of the study participants and their families. Medical writing assistance was provided by Blair Hesp PhD CMPP of Kainic Medical Communications (Dunedin, New Zealand), which was funded by Sobi (Basel, Switzerland). PQ thanks Ms Thi An Vu (Clinical Research Unit, Necker Hospital, Paris, France) for her support. PAB acknowledges the conduct of the trial at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health being supported by the NIHR Great Ormond Street Hospital Biomedical Research Centre. AAG acknowledges PRCSG Clinical Research Associates at Cincinnati Children's Hospital.

Contributors FDB, AAG, CdM, MB and PJ provided substantial contributions to the conception of the work. All authors substantially contributed to the acquisition, analysis or interpretation of data for the manuscript. All authors participated in drafting, revising and critically reviewing the manuscript for important intellectual content. All authors approved final version of this manuscript to be published and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. FDB is responsible for the overall content as guarantor.

Funding NI-0501-05 and NI-0501-06 (ClinicalTrials.gov numbers, NCT02069899 and NCT03311854) were funded by Sobi.

Competing interests FDB: consultant and research grants from AbbVie, Sobi, Pfizer, Roche, Sanofi, Novartis, Novimmune; AAG: consultant for Novartis, AB2 Bio, Novimmune, Sobi; PAB: consultant for Sobi, Novartis, Roche, UCB; CB, MP, GM: none declared; DE: speaker bureau for Sobi; CP: speaker bureau for Sobi; GS: consultant for Novartis, Sobi, Novimmune, AB2 Bio; PQ: consultant for AbbVie, Chugai-Roche, Lilly, Novartis, Pfizer, Sobi and speaker bureau for AbbVie, Chugai-Roche, Lilly, Novartis, Pfizer, Sobi; JA: consultant for Sobi, Novartis, Roche, Pfizer, AbbVie, GSK; CL: consultant for Sobi; RF: previously employed by Sobi; VA: previously employed by Sobi; MB: previously employed by Sobi; PJ: consultant for Sobi; CdM: consultant for Sobi, previously employed by Sobi.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the 'Methods' section for further details.

Patient consent for publication Not applicable.

Ethics approval Approval was obtained from institutional review boards or independent ethics committees at each participating centre. The initial approval was obtained from the Institutional Review Board at Ospedale Pediatrico Bambino Gesù, IRCCS, Rome, Italy (reference number 1245/2016). Participants gave written informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The data sharing policy of Sobi is available at the following site: <https://www.sobi.com/en/policies>. As noted on that site, requests for access to the study data can be submitted to medical.info@sobi.com.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Fabrizio De Benedetti <http://orcid.org/0000-0001-8749-8232>

Alexei A Grom <http://orcid.org/0000-0001-5717-136X>

Paul A Brogan <http://orcid.org/0000-0001-6178-6893>

Claudia Bracaglia <http://orcid.org/0000-0002-9834-9619>

Charalampia Papadopoulou <http://orcid.org/0000-0002-1237-0557>

Grant S Schultert <http://orcid.org/0000-0001-5923-7051>
 Jordi Antón <http://orcid.org/0000-0002-8792-4219>

REFERENCES

- Grom AA, Horne A, De Benedetti F. Macrophage activation syndrome in the era of biologic therapy. *Nat Rev Rheumatol* 2016;12:259–68.
- Efthimiou P, Kontzias A, Hur P, et al. Adult-onset still's disease in focus: clinical manifestations, diagnosis, treatment, and unmet needs in the era of targeted therapies. *Semin Arthritis Rheum* 2021;51:858–74.
- Jamilloux Y, Gerfaud-Valentin M, Martinon F, et al. Pathogenesis of adult-onset still's disease: new insights from the juvenile counterpart. *Immunol Res* 2015;61:53–62.
- Minoia F, Davi S, Horne A, et al. Clinical features, treatment, and outcome of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis: a multinational, multicenter study of 362 patients. *Arthritis Rheumatol* 2014;66:3160–9.
- Nigrovic PA, De Benedetti F, Kimura Y, et al. The elephant in the room: diagnostic/ classification criteria for systemic juvenile idiopathic arthritis and adult-onset still's disease [In press]. *Pediatr Rheumatol* 2022.
- De Matteis A, Colucci M, Rossi MN, et al. Expansion of cd4dimcd8+ T cells characterizes macrophage activation syndrome and other secondary HLH. *Blood* 2022;140:262–73.
- Chaturvedi V, Marsh RA, Zoreff-Lorenz A, et al. T-cell activation profiles distinguish hemophagocytic lymphohistiocytosis and early sepsis. *Blood* 2021;137:2337–46.
- Erkens R, Esteban Y, Towe C, et al. Pathogenesis and treatment of refractory disease courses in systemic juvenile idiopathic arthritis: refractory arthritis, recurrent macrophage activation syndrome and chronic lung disease. *Rheum Dis Clin North Am* 2021;47:585–606.
- Horne A, von Bahr Greenwood T, Chiang SCC, et al. Efficacy of moderately dosed etoposide in macrophage activation syndrome-hemophagocytic lymphohistiocytosis. *J Rheumatol* 2021;48:1596–602.
- Eloseily EM, Weiser P, Crayne CB, et al. Benefit of anakinra in treating pediatric secondary hemophagocytic lymphohistiocytosis. *Arthritis Rheumatol* 2020;72:326–34.
- Boom V, Anton J, Lahdenne P, et al. Evidence-based diagnosis and treatment of macrophage activation syndrome in systemic juvenile idiopathic arthritis. *Pediatr Rheumatol Online J* 2015;13:55.
- Verwey EL, Schultert GS. Interfering with interferons: targeting the JAK-STAT pathway in complications of systemic juvenile idiopathic arthritis (SJIA). *Rheumatology (Oxford)* 2022;61:926–35.
- De Benedetti F, Prencipe G, Bracaglia C, et al. Targeting interferon- γ in hyperinflammation: opportunities and challenges. *Nat Rev Rheumatol* 2021;17:678–91.
- Girard-Guyonvarc'h C, Palomo J, Martin P, et al. Unopposed IL-18 signaling leads to severe TLR9-induced macrophage activation syndrome in mice. *Blood* 2018;131:1430–41.
- Prencipe G, Caiello I, Pascarella A, et al. Neutralization of IFN- γ reverts clinical and laboratory features in a mouse model of macrophage activation syndrome. *J Allergy Clin Immunol* 2018;141:1439–49.
- Weiss ES, Girard-Guyonvarc'h C, Holzinger D, et al. Interleukin-18 diagnostically distinguishes and pathogenically promotes human and murine macrophage activation syndrome. *Blood* 2018;131:1442–55.
- Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 2011;89:207–15.
- Bracaglia C, de Graaf K, Pires Marafon D, et al. Elevated circulating levels of interferon- γ and interferon- γ -induced chemokines characterise patients with macrophage activation syndrome complicating systemic juvenile idiopathic arthritis. *Ann Rheum Dis* 2017;76:166–72.
- Mizuta M, Shimizu M, Inoue N, et al. Clinical significance of serum CXCL9 levels as a biomarker for systemic juvenile idiopathic arthritis associated macrophage activation syndrome. *Cytokine* 2019;119:182–7.
- Rodriguez-Smith JJ, Verwey EL, Clay GM, et al. Inflammatory biomarkers in COVID-19-associated multisystem inflammatory syndrome in children, Kawasaki disease, and macrophage activation syndrome: a cohort study. *Lancet Rheumatol* 2021;3:e574–84.
- Takakura M, Shimizu M, Irabu H, et al. Comparison of serum biomarkers for the diagnosis of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis. *Clin Immunol* 2019;208:108252.
- Han JH, Suh C-H, Jung J-Y, et al. Elevated circulating levels of the interferon- γ -induced chemokines are associated with disease activity and cutaneous manifestations in adult-onset still's disease. *Sci Rep* 2017;7:46652.
- Petty RE, Southwood TR, Manners P, et al. International League of Associations for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001. *J Rheumatol* 2004;31:390–2.
- Yamaguchi M, Ohta A, Tsunematsu T, et al. Preliminary criteria for classification of adult still's disease. *J Rheumatol* 1992;19:424–30.
- DeWitt EM, Kimura Y, Beukelman T, et al. Consensus treatment plans for new-onset systemic juvenile idiopathic arthritis. *Arthritis Care Res (Hoboken)* 2012;64:1001–10.
- Ravelli A, Minoia F, Davi S, et al. 2016 classification criteria for macrophage activation syndrome complicating systemic juvenile idiopathic arthritis: a European League Against Rheumatism/American College of Rheumatology/Paediatric Rheumatology International Trials Organisation collaborative initiative. *Arthritis Rheumatol* 2016;68:566–76.
- Locatelli F, Jordan MB, Allen C, et al. Emapalumab in children with primary hemophagocytic lymphohistiocytosis. *N Engl J Med* 2020;382:1811–22.
- Jacquin P, Laveille C, Snoeck E, et al. Emapalumab in primary haemophagocytic lymphohistiocytosis and the pathogenic role of interferon gamma: a pharmacometric model-based approach. *Br J Clin Pharmacol* 2022;88:2128–39.
- Pascarella A, Bracaglia C, Caiello I, et al. Monocytes from patients with macrophage activation syndrome and secondary hemophagocytic lymphohistiocytosis are hyperresponsive to interferon gamma. *Front Immunol* 2021;12:663329.
- Schultert GS, Pickering AV, Do T, et al. Monocyte and bone marrow macrophage transcriptional phenotypes in systemic juvenile idiopathic arthritis reveal TRIM8 as a mediator of IFN- γ hyper-responsiveness and risk for macrophage activation syndrome. *Ann Rheum Dis* 2021;80:617–25.
- Lounder DT, Bin Q, de Min C, et al. Treatment of refractory hemophagocytic lymphohistiocytosis with emapalumab despite severe concurrent infections. *Blood Adv* 2019;3:47–50.
- McNerney KO, DiNofia AM, Teachey DT, et al. Potential role of IFN γ inhibition in refractory cytokine release syndrome associated with CAR T-cell therapy. *Blood Cancer Discov* 2022;3:90–4.
- Tucci F, Gallo V, Barzaghi F, et al. Emapalumab treatment in an ADA-SCID patient with refractory hemophagocytic lymphohistiocytosis-related graft failure and disseminated Bacillus Calmette-Guérin infection. *Haematologica* 2021;106:641–6.
- Lam MT, Coppola S, Krumbach OHF, et al. A novel disorder involving dyshematopoiesis, inflammation, and HLH due to aberrant Cdc42 function. *J Exp Med* 2019;216:2778–99.
- Triebwasser MP, Barrett DM, Bassiri H, et al. Combined use of emapalumab and ruxolitinib in a patient with refractory hemophagocytic lymphohistiocytosis was safe and effective. *Pediatr Blood Cancer* 2021;68:e29026.
- Shamriz O, Kumar D, Shim J, et al. T cell-Epstein-Barr virus-associated hemophagocytic lymphohistiocytosis (HLH) occurs in non-Asians and is associated with a T cell activation state that is comparable to primary HLH. *J Clin Immunol* 2021;41:1582–96.
- Gabr JB, Liu E, Mian S, et al. Successful treatment of secondary macrophage activation syndrome with emapalumab in a patient with newly diagnosed adult-onset still's disease: case report and review of the literature. *Ann Transl Med* 2020;8:887.
- Urien S, Bardin C, Bader-Meunier B, et al. Anakinra pharmacokinetics in children and adolescents with systemic-onset juvenile idiopathic arthritis and autoinflammatory syndromes. *BMC Pharmacol Toxicol* 2013;14:40.
- Vastert SJ, Jamilloux Y, Quartier P, et al. Anakinra in children and adults with still's disease. *Rheumatology (Oxford)* 2019;58:vi9–22.
- KINERET (anakinra). Summary of product characteristics. 2022. Available: www.ema.europa.eu/en/documents/product-information/kineret-epar-product-information_en.pdf
- Mehta P, Cron RQ, Hartwell J, et al. Silencing the cytokine storm: the use of intravenous anakinra in haemophagocytic lymphohistiocytosis or macrophage activation syndrome. *Lancet Rheumatol* 2020;2:e358–67.
- Shih H-P, Ding J-Y, Yeh C-F, et al. Anti-interferon- γ autoantibody-associated immunodeficiency. *Curr Opin Immunol* 2021;72:206–14.