

Correspondence on 'Historically controlled comparison of glucocorticoids with or without tocilizumab versus supportive care only in patients with COVID-19-associated cytokine storm syndrome: results of the CHIC study'

We have read with great interest the study conducted by Ramiro *et al*¹ entitled 'Historically controlled comparison of glucocorticoids with or without tocilizumab versus supportive care only in patients with COVID-19-associated cytokine storm syndrome: results of the CHIC study'. This study was among the first publications showing benefit of tocilizumab administration in patients with severe COVID-19-associated cytokine storm syndrome, in July 2020.¹ In the same way, we observed a benefit of tocilizumab administration in a cohort of 206 patients during the first wave of the pandemic, in June 2020.²

We have now more 'evidence-based medicine' to discuss tocilizumab's place in COVID-19. Thus, nine randomised clinical trials (RCTs)^{3–11} have been published about tocilizumab administration in COVID-19; however, the effects on mortality remain heterogeneous in these trials.¹² We recently made a review of these RCTs¹³ and concluded that heterogeneity results seen in overall RCTs are due to the heterogeneous population and that tocilizumab is effective in severe COVID-19. Based on this conclusion, we performed a subgroup meta-analysis on tocilizumab RCTs (in severe COVID-19 and non-severe COVID-19). We selected by a systematic search on PubMed and the preprint server MedRxiv (until 27 April 2021) all RCTs that compared the outcome of patients with COVID-19 treated with tocilizumab versus standard of care or placebo (our endpoint was the mortality of 28–30 days). Due to the heterogeneous description about respiratory support at baseline (different scales or clinical descriptions were used in the RCTs), we cannot divide severe COVID-19 and non-severe COVID-19. However, in these RCTs, the mortality increases in correlation with the severity of respiratory support at baseline¹³; furthermore, clinical severity at baseline is among the main risk factors associated with mortality in COVID-19 pneumonia.¹⁴ So, we chose to use the mortality in control group to divide the RCTs in two groups: severe COVID-19 group (high mortality in control group) and non-severe COVID-19 group (low mortality in control group). Most of these RCTs included patients in the beginning of the pandemic in wealthy countries (especially North America and Europe), so we chose a mortality proportion of 17% which corresponds

to the in-hospital mortality in the beginning of the pandemic among hospitalised adults with COVID-19 in the USA.¹⁵ Our analysis RCTs were divided in two groups: severe COVID-19 group (mortality proportion in control group $\geq 17\%$) and non-severe COVID-19 group (mortality proportion in control group $< 17\%$). Overall, there were 24.5% (821 of 3357) deaths in the tocilizumab group and 29.1% (908 of 3125) in the control group at days 28–30 (pooled OR, 0.85; 95% CI, 0.76 to 0.96; $p=0.006$). Considering the subgroup analysis, this benefit was confirmed and amplified in the severe COVID-19 group (pooled OR, 0.82; 95% CI, 0.73 to 0.93; $p=0.001$) but not in the non-severe COVID-19 group (pooled OR, 1.46; 95% CI, 0.91 to 2.34; $p=0.12$) (figure 1).

Based on these results, we conclude that tocilizumab is an effective treatment in severe COVID-19 pneumonia by improving survival. Subgroup analyses are needed in tocilizumab RCTs to determine more accurately tocilizumab's place in COVID-19.

Timothée Klopstein¹,¹ Vincent Gendrin,¹ Thierry Conrozier,² Aurélie Gerazime,³ Marc Puyraveau,³ Souheil Zayet¹

¹Infectious Disease Department, North Franche-Comté Hospital, Montbéliard, France

²Rheumatology Department, North Franche-Comté Hospital, Montbéliard, France

³Methodology unit, Clinical Investigation Center INSERM 1431, CHU Besançon, Besançon, France

Correspondence to Dr Timothée Klopstein, Infectious Disease Department, North Franche-Comté Hospital, 25209 Montbéliard, France; timothee.klopstein@hnfc.fr

Contributors TK, SZ and VG drafted the manuscript. AG and MP made the statistical analysis. All authors revised the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Klopstein T, Gendrin V, Conrozier T, *et al.* *Ann Rheum Dis* 2023;**82**:e153.

Received 12 May 2021

Accepted 14 May 2021

Published Online First 23 June 2021



► <http://dx.doi.org/10.1136/annrheumdis-2021-220787>

Ann Rheum Dis 2023;**82**:e153. doi:10.1136/annrheumdis-2021-220771

ORCID iD

Timothée Klopstein <http://orcid.org/0000-0003-4334-9889>

REFERENCES

- 1 Ramiro S, Mostard RLM, Magro-Checa C, *et al.* Historically controlled comparison of glucocorticoids with or without tocilizumab versus supportive care only in patients with COVID-19-associated cytokine storm syndrome: results of the CHIC study. *Ann Rheum Dis* 2020;**79**:1143–51.




Figure 1 Forest plot for the effect of tocilizumab on mortality at days 28–30 in randomised trials in severity event subgroup.

- 2 Klopfenstein T, Zayet S, Lohse A, *et al*. Impact of tocilizumab on mortality and/or invasive mechanical ventilation requirement in a cohort of 206 COVID-19 patients. *Int J Infect Dis* 2020;99:491–5.
- 3 Salvarani C, Dolci G, Massari M, *et al*. Effect of tocilizumab vs standard care on clinical worsening in patients hospitalized with COVID-19 pneumonia: a randomized clinical trial. *JAMA Intern Med* 2021;181:24–31.
- 4 Stone JH, Frigault MJ, Serling-Boyd NJ, *et al*. Efficacy of tocilizumab in patients hospitalized with Covid-19. *N Engl J Med* 2020;383:2333–44.
- 5 Salama C, Han J, Yau L, *et al*. Tocilizumab in patients hospitalized with Covid-19 pneumonia. *N Engl J Med* 2021;384:20–30.
- 6 Hermine O, Mariette X, Tharaux P-L, *et al*. Effect of tocilizumab vs usual care in adults hospitalized with COVID-19 and moderate or severe pneumonia: a randomized clinical trial. *JAMA Intern Med* 2021;181:32–40.
- 7 Veiga VC, Prats JAGG, Farias DLC, *et al*. Effect of tocilizumab on clinical outcomes at 15 days in patients with severe or critical coronavirus disease 2019: randomised controlled trial. *BMJ* 2021;372:n84.
- 8 Rosas IO, Brău N, Waters M, *et al*. Tocilizumab in hospitalized patients with severe Covid-19 pneumonia. *N Engl J Med* 2021;384:1503–16.
- 9 Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial - The Lancet [Internet]. Available: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)00676-0/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)00676-0/fulltext)
- 10 Gordon AC, Al-Beidh F, Rowan KM. Interleukin-6 receptor antagonists in critically ill patients with Covid-19 – preliminary report. *medRxiv* 2021.
- 11 Soin AS, Kumar K, Choudhary NS, *et al*. Tocilizumab plus standard care versus standard care in patients in India with moderate to severe COVID-19-associated cytokine release syndrome (COVINTOC): an open-label, multicentre, randomised, controlled, phase 3 trial. *Lancet Respir Med* 2021;9:511–521.
- 12 Tleyjeh IM, Kashour Z, Riaz M, *et al*. Efficacy and safety of tocilizumab in COVID-19 patients: a living systematic review and meta-analysis: first update. *Clin Microbiol Infect* 2021. doi:10.1016/j.cmi.2021.04.019. [Epub ahead of print: 26 Apr 2021].
- 13 Klopfenstein T, Gendrin V, Kadiane-Oussou N'dri Juliette, *et al*. Tocilizumab in COVID-19 pneumonia: practical proposals based on a narrative review of randomised trials. *Rev Med Virol* 2021:e2239.
- 14 Weiss P, Murdoch DR. Clinical course and mortality risk of severe COVID-19. *Lancet* 2020;395:1014–5.
- 15 Kim L, Garg S, O'Halloran A, *et al*. Risk Factors for Intensive Care Unit Admission and In-hospital Mortality Among Hospitalized Adults Identified through the US Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET). *Clin Infect Dis* 2021;72:e206–14.