

Supplementary online material – Summary Tables

MAKING A DIAGNOSIS OF OSTEOARTHRITIS – KNEE

Q2: What is the accuracy of different imaging modalities for detecting OA features (soft tissues and bone involvement)?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome				
						Se	Sp	LR+	LR-
Akgul O 2014 [1]	110	Cross-sectional	US	Physical examination	Popliteal cyst	0.66	0.98	42.22	0.34
Esen S 2013 [2]	100	Cross-sectional	US	Physical examination		Se	Sp	LR+	LR-
					Effusion	0.76	0.51	1.63	0.42
					Pes anserinus bursitis	0.50	0.95	12.25	0.52
					Popliteal cyst	0.36	0.88	3.27	0.72
Ike RW 2010 [3]	14	Cross-sectional	US	Physical examination, synovial fluid aspiration		Physical examination		SF aspiration	
						Se	Sp	Se	Sp
					Effusion	1	0	1	0
Pendleton A 2008 [4]	86	Cross-sectional	US	Physical examination		Se	Sp	LR+	LR-
					Synovitis	0.73	0.15	0.87	1.74
					Effusion	0.67	0.50	1.34	0.66
Chatzopoulos D 2008 [5]	196	Cross-sectional	US	Radioisotope scan		Se	Sp	LR+	LR-
					Popliteal cyst	0.28	0.90	2.89	0.79
Lee LC 2008 [6]	95	Cross-sectional	US	Surgery	US cartilage grading correlated to histologic grading over anterior and middle areas (r=0.40 and 0.36, p<0.001)				
Yoon CH 2008 [7]	51	Cross-sectional	US	CR MRI	US vs CR (cartilage)	Minimal thickness medial condyle		r=0.417	
						Maximal thickness medial condyle		r=0.412	
					US vs MRI (cartilage)	Minimal thickness medial condyle		r=0.568	
						Maximal thickness medial condyle		r=0.844	
Song IH 2008 [8]	41	Cross-sectional	US	MRI	Effusion superior recess (Se)			0.72	
					Effusion lateral recess (Se)			0.81	

Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; CR: conventional radiography; MRI: magnetic resonance imaging; US: ultrasonography; sd: standard deviation; r: correlation coefficient

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome				
Bhattacharya R 2007 [9]	77	Cross-sectional	CR	Surgery		Se	Sp	LR+	LR-
					Skyline (PFJ)	0.81	0.60	2.08	0.30
					Lateral (PFJ)	0.83	0.39	0.37	0.43
Chang CB 2008 [10]	151	Cross-sectional	CR	Surgery	JSN/cartilage	Se 0.83		Sp 0.50	
Dervin G 2001 [11]	152	Cross-sectional	CR	Surgery		AP full-extension		45°flexion	
						Se	Sp	Se	Sp
					JSN/cartilage	0.73	0.99	0.78	0.96
Waldstein W 2013 [12]	84	Cross-sectional	CR	Surgery	poor correlation lateral JSW/Outerbridge grades: $r = -0.154$; $p = 0.146$ significant correlation Outerbridge grades/ lateral KLG: $r = 0.553$; $p = 0.001$				
Ciccattini F 2003 [13]	252	Cross-sectional	CR	MRI	Significant correlation medial and lateral cartilage volume/ JSN TFJ ($p < 0.001$); no significant correlation with lateral TFJ JSN , $p = 0.059$				
Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; CR: conventional radiography; MRI: magnetic resonance imaging; r: correlation coefficient; PFJ: patellofemoral joint; JSN: joint space narrowing; JSW: joint space width									

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome								
Boegard T 1999 [14]	58	Cross-sectional	MRI	Radioisotope scan	k	MT	MF	LT	LF	patella	Femoral trochlea		
					Subchondral lesions	0.72	0.79	0.59	0.64	0.49	0.64		
					Osteophytes	0.20	0.30	0.14	0.31	0.03	0.10		
					Cartilage defects	0.54	0.26	0.42	0.15	0.06	0.18		
Crema MD 2013 [15]	20	Cross-sectional	MRI (3D)	MRI (2D)				Se	Sp				
					Bone attrition			0.84			0.99		
					Cartilage morphology			0.96			0.98		
					BMLs			0.84			0.97		
					Effusion			0.88			1		
					Meniscal tear			0.96			0.97		
					Meniscal extrusion			1.0			0.95		
					Osteophytes			0.86			0.85		
					Subchondral cysts			0.80			0.98		
Popliteal cysts			0.99			0.99							
Gudbergesen H 2013 [16]	192	Cross-sectional	MRI	CR	KLG correlated to cartilage damage, BMLs , meniscal pathology (r=0.15-0.76) BLOKS assessed knee joint pathology cosegregated with compartment and grade specific KLG (P < 0.0001)								
De Lange Borkaar 2014 [17]	42	Cross-sectional	MRI	Surgery	Synovitis grade r=0.57 p<0.001								
Fernandez Madrid F 1995 [18]	9	Cross-sectional	MRI	Surgery	Synovitis: Se 1.0								
Bergman A 1994 [19]	9	Cross-sectional	MRI	Surgery	The location and extent of the subchondral signal abnormalities on MR corresponded to the focal areas of fibrous tissue replacement of fatty marrow								
Broderick L 1994 [20]	28	Cross-sectional	MRI	Surgery				Se	Sp	LR+	LR-		
					Cartilage score			0.76			0.98		

CR: conventional radiography; MRI: magnetic resonance imaging; r: correlation coefficient; MT: medial tibia; MF: medial femur; LT: lateral tibia; LF: lateral femur; k: kappa statistics; Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; KLG: Kellgren and Lawrence grade; BMLs: bone marrow lesions

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome
Kalunian K 2000 [21]	8	Cross-sectional	MRI	Surgery	Meniscal tears: Se 0.33, Sp 0.96
Loeuille D 2011 [22]	30	Cross-sectional	MRI	Surgery, physical examination (SF aspiration)	No significant correlation between SF volume and histologic edema and congestion with MRI synovitis score
Saadat E 2008 [23]	8	Cross-sectional	MRI	Surgery	Se
					Sp
					Cartilage lesions
					0.72
					0.69
Takayama Y 2013 [24]	16	Cross-sectional	MRI	Surgery	AUC= 0.881 (T1p) AUC=0.681 (T2)
Von Engelhardt L 2010 [25]	32	Cross-sectional	MRI	Surgery	Grading of cartilage lesions
					Se
					Sp
					1
					0.20
2					
0.52					
3					
0.36					
4					
0.70					
0.94					
Wong CS 2013 [26]	28	Cohort	MRI	Surgery	T2 inversely correlated with GAG concentration (r2 = 0.375, p = 0.001) T1p values inversely correlated with GAG concentration (r2 = 0.200, p = 0.025)
Yoshioka H 2004 [27]	16	Cross-sectional	MRI	Surgery	Se
					Sp
					Cartilage lesions
					1.0
					0.676
Zanetti M 2000 [28]	16	Cross-sectional	MRI	Surgery	Significant differences between the zone with the edema pattern (zone 1) and the control zone (zone 6) for bone marrow fibrosis (P = 0.014), bone marrow necrosis (p = 0.021), and trabecular abnormalities (p = 0.011), not for bone marrow edema (p=0.069), cysts (p=0.343), bone marrow bleeding (p=0.696).
Graichen H 2004 [29]	21	Cross-sectional	MRI	Surgery	Cartilage Volume
					r=0.98
					Cartilage thickness
					r=0.92

Se: sensitivity; Sp: specificity; SF: synovial fluid; AUC: area under the curve; T1p: T1 relaxation time; T2: T2 relaxation time; MRI: magnetic resonance imaging

MANAGEMENT OF OSTEOARTHRITIS – KNEE

Q4: In people with OA do imaging have an impact on management?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome	
Brealey SD 2007[30]	553	RCT	MRI	-	Change in management	Change in therapeutic confidence
					Treatment altered: MRI 261/279 Orthopedic referral 258/266; Treatment not altered MRI 18/279 , Orthopedic referral 8/266. $p=0.059$	Increased: MRI 163/259, orthopedic 119/241 No change: MRI 87/259, orthopedic 100/241 Decreased: MRI 9/259, orthopedic 22/241 $p = 0.002$
Ritchie JF 2014[31]	50	Cross-sectional	CR	-	change in management: 166/400 opinions	

MRI: magnetic resonance imaging; CR: conventional radiography

PROGNOSIS OF OA-KNEE

Q5: What is the prognostic (prediction of outcome) value of imaging in OA?

Cross-sectional studies, CR: symptoms

Reference	No of subjects	Study design	Imaging modality	Outcome					
Barker K 2004 [32]	123	Cross-sectional	CR	KLG/WOMAC					
Cubukcu D 2012 [33]	114	Cross-sectional	CR	KLG/WOMAC					
Bevers K 2014 [34]	180	Cross-sectional	CR	KLG/pain					
Neogi T 2010 [35]	2940	Cross-sectional	CR	KLG/inconsistent pain vs consistent pain	OR (95% CI)				
					KLG 0	1			
					KLG 1	1.01 (0.80-1.27)			
					KLG 2	0.78 (0.61-0.99)			
					KLG 3	0.51 (0.41,0.63)			
KLG 4	0.27 (0.18,0.36)								
Sanghi D 2011 [36]	180	Cross-sectional	CR	KLG/VAS pain and WOMAC	VAS pain	WOMAC pain	WOMAC stiffness	WOMAC function	
					KLG 2	5.86 (1.25)	9.86 (1.91)	2.14 (1.35)	21.36 (5.03)
					KLG 3	6.27 (0.81)	10.32 (2.04)	2.28 (1.25)	23.50 (5.73)
					KLG 4	6.31 (1.05)	10.98 (1.71)	2.56 (1.42)	24.82 (5.74)
					p	0.15	0.03	0.32	0.05
Madsen OR 1995 [37]	20	Cross-sectional	CR	KLG/ pain and function	Pain disability score	r = 0.50, p < 0.05			
					Muscle strength , walking and stair climbing	No significant correlation			

CR: conventional radiography; KL/KLG: Kellgren and Lawrence grade; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index; R: correlation coefficient; VAS: visual analogue scale; OR (95% CI): odds ratio and 95% confidence interval.

Reference	Study design	No of subjects	Imaging modality	Outcome					
Cicutti F 1996 [38]	Cross-sectional	250	CR	JSN/osteophytes and pain			OR (95%CI)		
					AP view	JSN	2.13 (0.78,5.87)		
						osteophytes	5.00 (2.40,10.43)		
					Lateral view	JSN	1.54 (0.57,4.14)		
						osteophytes	2.87 (1.41,5.82)		
					Skyline view	JSN	1.42 (0.53,3.82)		
osteophytes	7.56 (3.85,14.81)								
Hernandez Vaquero D 2012 [39]	Cross-sectional	1329	CR	Ahlback grade/Hospital for Special Surgery Knee-Rating Score	Mean (sd)	I 56.91 (10.75); II 55.64 (12.01) III 51.86 (12.96); IV 49.44 (12.69) V 47.35 (12.59) p<0.001			
Lim W 2008 [40]	Cross-sectional	107	CR	Varus alignment/WOMAC pain	WOMAC pain mean, sd (no significant differences)	Least varus	Moderate varus	Most varus	
						34.9 ± 16.3	35.7 ± 14.6	36.5 ± 14.6	
Illingworth K 2014 [41]	Cross-sectional		CR MRI	JSW/KOOS and WOMAC		Medial TF (r, p)	Lateral TF (r, p)		
					KOOS pain	0.03 p>0.05	0.06 p>0.05		
					KOOS symptoms	0.16 p<0.05	0.15 p<0.05		
					WOMAC pain	0.08 p>0.05	0.05 p>0.05		
					WOMAC stiffness	0.10 p>0.05	0.09 p>0.05		
Knoop J 2012 [42]	Cross-sectional	105	CR MRI	CR and MRI/bony tenderness		MTF (OR, p)	LTF (OR, p)	PF (OR, p)	
					JSW	0.90 (0.82)	0.96 (0.95)	1.80 (0.27)	
					CR osteophytes	0.77 (0.59)	0.85 (0.81)	1.34 (0.58)	
					Cartilage integrity	0.77 (0.59)	0.85 (0.81)	1.34 (0.58)	
					MRI osteophytes	0.92 (0.32)	1.06 (0.60)	1.09 (0.20)	
					BMLs	0.97 (0.76)	0.91 (0.53)	0.84 (0.21)	
					Effusion	0.75 (0.52)			
					Synovitis	0.84 (0.70)			
Link TM 2003 [43]	Cross-sectional	50	CR MRI	CR and MRI/WOMAC	Cartilage lesions	Significantly higher WOMAC pain and function			
					Central osteophytes	Significantly higher WOMAC stiffness			
					BMLs, effusion, popliteal cysts, subchondral cysts, meniscal and ligamentous lesions	No significant differences in WOMAC subscales			

CR: conventional radiography; MRI: magnetic resonance imaging; AP: anteroposterior; JSN: joint space narrowing; sd: standard deviation; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index; TF: tibiofemoral; KOOS: Knee injury and Osteoarthritis Outcome Score; JSW: joint space width; BMLs: bone marrow lesions

Cross-sectional studies, CR: structure

Reference	No of subjects	Study design	Imaging modality	Outcome				
Hernigou P 2002 [44]	40	Cross-sectional	CR CT	Intercondylar notch width/ACL damage				
				CT notch width (mean, sd)	Normal ACL	Damaged ACL	No ACL	
				CR notch width (mean, sd)	19.6 (1.7)	14.2 (2.5)	10.4 (2.5)	
Janakiramanan N 2008 [45]	202	Cross-sectional	CR	Alignment/MRI cartilage lesions				
Issa S 2007 [46]	146	Cross-sectional	CR	Alignment/MRI cartilage lesions				
Khan FA 2008 [47]	306	Cross-sectional	CR	Alignment/compartmental OA				
Wang L 2013 [48]	26	Cross-sectional	CR	Alignment/MRI T1p				
				LF (mean, sd)	Varus	Neutral	Valgus	p
				MF (mean, Sd)	64±14	58±10	53±2	0.0016, 0.047
				MT (mean, sd)	69±9	-	59±10	0.048
Friedrich K 2010 [49]	24	Cross-sectional	CR	Alignment/T2				
Teichtahl A 2006 [50]	121	Cross-sectional	CR	Alignment/JSN and osteophytes				
				OR (95%CI) for degree of valgus	JSN	Osteophytes		
				Medial	0.77 (0.70, 0.85)	0.90 (0.83, 0.97)		
von Eisenhart Rothe 2006 [51]	26	Cross-sectional	CR	Alignment/MRI cartilage thickness				
Boissonneault A 2014 [52]	664	Cross-sectional	CR	CR OA/alignment				
				Lateral cartilage thickness	r=-0.54; p=0.01			
Lateral compartment knee OA (no OA as reference): more valgus OR 1.03 (1.00, 1.07) Medial compartment knee OA (no OA as reference): more varus OR 0.97 (0.94, 1.00)								
CR: conventional radiography; MRI: magnetic resonance imaging; CT: computer tomography; ACL: anterior cruciate ligament; sd: standard deviation; OR (95% CI): odds ratio with and 95% confidence interval; T1p: T1 relaxation time; T2: T2 relaxation time; JSN: joint space narrowing.								

Reference	No of subjects	Study design	Imaging modality	Outcome				
Ozgul G 2013 [53]	50	Cross-sectional	CR	PF angle and sulcus angle/KLG		KLG 1	KLG 2-4	p
					PF angle (mean sd)	23.0±4.5	19.6±6.2	0.005
					Sulcus angle (mean sd)	128.4±6.9	130.0±7.6	0.29
van der Esch M 2014 [54]	298	Cross-sectional	CR	Association between compartments and within compartments of JSN, osteophytes, sclerosis, cysts	Association among features within the joint and association among features within each compartment			
Cotofana S 2012 [55]	61	Cross-sectional	CR	Osteophytes/MRI cartilage thickness	Significantly lower cartilage thickness in different compartment in knees with osteophytes			
Ferraro B 2010 [56]	483	Case-control	CR	2D:4D type 3 finger pattern/knee OA	OR (95% CI) 2.59 (1.54,4.37)			
CR: conventional radiography; MRI: magnetic resonance imaging; sd: standard deviation; OR (95% CI): odds ratio with and 95% confidence interval; KLG: Kellgren and Lawrence grade; 2D:4D : 2 nd finger/4 th finger ratio								

Cross-sectional studies, CR: function

Reference	No of subjects	Study Design	Imaging modality	Outcome		TF medial (p)	TF lateral (p)	PF (p)
Holla JF 2011 [57]	497	Cross-sectional	CR	CR/range of motion				
					JSN	0.001	0.83	0.96
					Osteophytes	0.0001	0.01	0.0001
					Subchondral sclerosis	0.02		
				Tibial tubercles spiking	0.07			
McDaniel C 2011 [58]	138	Cross-sectional	CR	CR/gait velocity and standing balance		Gait velocity	Standing balance	
					JSN	r=-0.19	r=-0.04	
					Osteophytes	r=-0.16	r=-0.01	
					KLG	r=-0.12	r=-0.02	

CR: conventional radiography; TF: tibiofemoral; PF: patellofemoral; JSN: joint space narrowing; KLG: Kellgren and Laerence grade; r: correlation coefficient

Cross-sectional studies, MRI: symptoms

Reference	No of subjects	Study design	Imaging modality	Outcome			
Ai F 2010 [59]	28	Cross-sectional	MRI	MRI/severe pain	OR (95%CI)		
					Cartilage defects	0.63 (0.34,1.82)	
					Medial meniscal lesion	1.43 (0.28,7.21)	
					Lateral meniscal lesion	0.87 (0.24,3.12)	
					BML	0.38 (0.15,2.98)	
					Effusion	0.043 (0.00235,0.83)	
De Lange Brokaar 2014[17]	42	Cross-sectional	MRI	MRI synovitis/pain	Spearman's correlation 0.32 p=0.041		
Ballegaard C 2014 [60]	95	Cross-sectional	MRI DCE	MRI/KOOS pain	KOOS pain		
					Perfusion	r= -0.42 (p<0.0001)	
					Hoffa synovitis	r= -0.21 (p=0.046)	
					Slope	r= -0.33 (p<0.0001)	
Lo GH 2009 [61]	160	Cross-sectional	MRI	MRI/pain	Weightbearing pain (OR 95%CI)	Non weightbearing pain (OR 95%CI)	
					Effusion	4.30 (1.71,10.83)	1.72 (0.60,4.89)
					Synovitis	3.04 (0.57,16.17)	0.94 (0.17,5.06)
					BMLs	3.20 (0.96,10.68)	1.47 (0.37,5.39)
					OR (95%CI)	Pain	Stiffness
Kornaat P 2006 [62]	71	Cross-sectional	MRI	MRI/pain and stiffness	Cartilage defects	1.12 (0.40, 3.14)	1.35 (0.48, 3.82)
					Osteophytes	1.05 (0.38, 2.91)	1.44 (0.52, 4.04)
					Subchondral cysts	1.71 (0.81, 3.63)	1.25 (0.60, 2.59)
					Meniscal tear	1.26 (0.58, 2.74)	0.64 (0.30, 1.39)
					Meniscal subluxation	1.03 (0.48, 2.21)	1.63 (0.76, 3.46)
					Effusion	9.99 (1.13, 149)	4.67 (1.11, 26.14)
					Popliteal cyst	1.68 (0.80, 3.53)	1.32 (0.64, 2.73)
					BMLs	1.36 (0.65, 2.84)	1.57 (0.76, 3.25)

MRI: magnetic resonance imaging; OR (95% CI): odds ratio with and 95% confidence interval; BML: bone marrow lesions; KOOS: Knee injury and Osteoarthritis Outcome Score; r: correlation coefficient

Reference	No of subjects	Study design	Imaging modality	Outcome				
Torres L 2006 [63]	143	Cross-sectional	MRI	MRI/VAS pain	increase in median knee pain severity (VAS pain) for 1-unit increase in lesion score			
					Bone attrition	1.91		
					Bone cysts	0.82		
					Osteophytes	0.50		
					Cartilage	0.53		
					Meniscal tears	1.99		
					Meniscal extrusion (1 vs 0)	-2.96		
					Meniscal extrusion (2 vs 0)	2.22		
					Synovitis/effusion (1 vs 0)	1.93		
Synovitis/effusion (2-3 vs 0)	9.82							
Lange AK 2007 [64]	41	Cross-sectional	MRI	Meniscal tears/WOMAC and function	Mean (sd)	Tear	No tear	p
					WOMAC pain	6.5 (3.2)	5.3 (2.6)	0.26
					WOMAC stiffness	3.8 (1.5)	3.6 (1.6)	0.71
					WOMAC function	23.9 (10.6)	21.1 (11.7)	0.47
					Habitual physical activity level	136 (56)	134.4 (60.5)	0.93
					6 minute walk	504.2 (79)	560.9 (65.6)	0.04
					Gait velocity	1.08 (0.16)	1.10 (0.21)	0.77
Balance	95.6 (18.8)	81.5 (19.1)	0.04					
Bhattacharrya R 2003 [65]	154	Cross-sectional	MRI	meniscal tears/WOMAC and VAS pain		Tear	No tear	p
					WOMAC (mean, sd)	29 (4.1)	29 (3.0)	0.91
					VAS pain (mean, sd)	31 (8.3)	34 (3.1)	0.67
Wenger A 2012 [66]	53	Cross-sectional	MRI	Area covered by the meniscus/pain	Mean, sd	pain	No pain	p
					Medial meniscus	41% (10)	44% (7)	0.01
					Lateral meniscus	-	-	>0.05
Seah S 2012 [67]	37	Cross-sectional	MRI	BMLs/pain at night	BML grade 3 vs 1 (OR 95% CI)	0.38 (0.07,2.01)		
					BML grade 3 vs 2	0.44 (0.08,2.58)		
Felson DT 2001 [68]	287	Cross-sectional	MRI	BMLs/pain		OR (95%CI)		
					BMLs	3.31 (1.54,7.41)		
					Large BMLs	(1.04, 111.11)		

MRI: magnetic resonance imaging; VAS: visual analogue scale; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index; sd: standard deviation; BML: bone marrow lesions

Reference	No of subjects	Study design	Imaging modality	Outcome	Reference		
Sower M 2003 [69]	113	Cross-sectional	MRI	BMLs and cartilage defects/pain	OR (95% CI)		
					BMLs	0.85 (0.365,1.98)	
					Cartilage defects	0.203 (0.023,1.80)	
Sengupta N 2006 [70]	217	Cross-sectional	MRI	Osteophyte BMLs/pain	OR (95% CI) 0.690 (0.245, 1.942)		
Wluka AE 2004 [71]	117	Cross-sectional	MRI	cartilage volume/WOMAC	WOMAC pain	r=-0.17 p=0.05	
					WOMAC stiffness	r=-0.16 p=0.08	
					WOMAC function	r=-0.20 p=0.03	
Hernandez Molina G 2008 [72]	1627	Cross-sectional	MRI	Bone attrition/pain	OR (95% CI)=1.2 (0.7, 2.0)		
Baker K 2010 [73]	454	Cross-sectional	MRI	synovitis/pain	OR (95% CI)=2.35 (1.57,3.50)		
Hill C 2001 [74]	458	Cross-sectional	MRI	Effusion and popliteal cysts/pain	Pain (%)	No pain (%)	
					Small effusion	37.1	62.3
					Moderate effusion	36.0	13.2
					Large effusion	18.6	1.4
					Popliteal cysts	33.0	20.8
Hill C 2005 [75]	403	Case-control	MRI	ACL tear/VAS pain	VAS pain (mean, sd): complete ACL tear 44.3 mm, intact ACL 44.1 mm (p=0.95)		
Lindsey CT 2004 [76]	53	Cross-sectional	MRI	Cartilage volume/WOMAC pain	medial tibia volume and WOMAC pain score: r= -0.31 (p=0.017)		
Cotofana S 2013 [77]	633	Cross-sectional	MRI	cartilage defects/pain	OR (95%CI)		
					Weightbearing pain	1.64 (1.20,2.19)	
					Non weightbearing pain	1.52 (1.02,2.28)	
					Infrequent pain vs no pain	1.85 (1.23,2.80)	
					Frequent pain vs no pain	2.62 (1.80,3.81)	
Moderate/severe pain	1.80 (1.32,2.47)						

MRI: magnetic resonance imaging; OR (95%CI): odds ratio, 95% confidence interval ; WOMAC: Western Ontario and MacMAster Universities Osteoarthritis Index; sd: standard deviation; BML: bone marrow lesions; OR (95%CI): odds ratio, 95% confidence interval; ACL: anterior cruciate ligament

Cross-sectional studies, MRI: structure

Reference	No of subjects	Study design	Imaging modality	Outcome												
Gale DR 1999 [78]	291	Case-control	MRI	Meniscal subluxation/JSN <table border="1"> <tr> <td>Medial meniscus</td> <td>r=0.55 p=0.0001</td> </tr> <tr> <td>Lateral meniscus</td> <td>r=0.40 p=0.0001</td> </tr> </table>	Medial meniscus	r=0.55 p=0.0001	Lateral meniscus	r=0.40 p=0.0001								
Medial meniscus	r=0.55 p=0.0001															
Lateral meniscus	r=0.40 p=0.0001															
Friedrich K 2009 [79]	37	Cross-sectional	MRI	Meniscal pathology/T2 Mean (sd) T2 in patients with meniscal tears vs patients without meniscal tears: 50.1 (6.1) vs 45.7 (4.8) p=0.0021												
Hovis K 2011 [80]	304	Cross-sectional	MRI	ACL abnormalities/WORMS Significantly higher total WORMS scores in abnormal ACL compared to normal ACL												
Jungmann P 2013 [81]	135	Case-control	MRI	Shallow trochlea/PF WORMS and T2 <table border="1"> <tr> <td></td> <td>Shallow trochlea</td> <td>Deep trochlea</td> <td>p</td> </tr> <tr> <td>PF WORMS (mean,sd)</td> <td>11.2 (0.5)</td> <td>5.7 (0.6)</td> <td><0.001</td> </tr> <tr> <td>T2 (mean,sd)</td> <td>40.9 (0.5)</td> <td>42.7 (0.6)</td> <td>0.037</td> </tr> </table>		Shallow trochlea	Deep trochlea	p	PF WORMS (mean,sd)	11.2 (0.5)	5.7 (0.6)	<0.001	T2 (mean,sd)	40.9 (0.5)	42.7 (0.6)	0.037
	Shallow trochlea	Deep trochlea	p													
PF WORMS (mean,sd)	11.2 (0.5)	5.7 (0.6)	<0.001													
T2 (mean,sd)	40.9 (0.5)	42.7 (0.6)	0.037													
Kalichman L 2007 [82]	2013	Cross-sectional	MRI	Lateral patellar tilt ratio and bisect offset/JSN and osteophytes LPTA associated with lower degree of lateral and medial patellar JSN and lateral osteophyte. BO associated with higher degree of lateral and medial patellar JSN and lateral osteophytes.												
Kornaat P 2005 [83]	205	Cross-sectional	MRI	MRI cartilage defects /MRI cartilage defects <table border="1"> <tr> <td></td> <td>OR (95%CI)</td> </tr> <tr> <td>TF cartilage defects in case of PF cartilage defects</td> <td>3.95 (2.12,7.37)</td> </tr> <tr> <td>PF cartilage defects in case of PF BME</td> <td>13.6 (4.69,39.44)</td> </tr> <tr> <td>TF cartilage defects in case of TF BME</td> <td>3.99 (1.48,10.77)</td> </tr> <tr> <td>PF cartilage defects and ipsilateral meniscal tears</td> <td>3.7 (1.0,15)</td> </tr> <tr> <td>TF cartilage defects/ipsilateral meniscal tears</td> <td>9.8 (2.5,38)</td> </tr> </table>		OR (95%CI)	TF cartilage defects in case of PF cartilage defects	3.95 (2.12,7.37)	PF cartilage defects in case of PF BME	13.6 (4.69,39.44)	TF cartilage defects in case of TF BME	3.99 (1.48,10.77)	PF cartilage defects and ipsilateral meniscal tears	3.7 (1.0,15)	TF cartilage defects/ipsilateral meniscal tears	9.8 (2.5,38)
	OR (95%CI)															
TF cartilage defects in case of PF cartilage defects	3.95 (2.12,7.37)															
PF cartilage defects in case of PF BME	13.6 (4.69,39.44)															
TF cartilage defects in case of TF BME	3.99 (1.48,10.77)															
PF cartilage defects and ipsilateral meniscal tears	3.7 (1.0,15)															
TF cartilage defects/ipsilateral meniscal tears	9.8 (2.5,38)															
Krasnutotsky K 2011 [84]	58	Cross-sectional	MRI	Quantitative synovial volume/KLG JSN JSW Significant correlation with KLG, JSN and JSW												
Lo GH 2009 [85]	160	Cross-sectional	MRI	Meniscal derangments/large BMLs <table border="1"> <tr> <td>OR (95%CI)</td> <td>Medial compartment</td> <td>37.21 (2.22,621.64)</td> </tr> <tr> <td></td> <td>Lateral compartment</td> <td>4.47 (1.61,12.37)</td> </tr> </table>	OR (95%CI)	Medial compartment	37.21 (2.22,621.64)		Lateral compartment	4.47 (1.61,12.37)						
OR (95%CI)	Medial compartment	37.21 (2.22,621.64)														
	Lateral compartment	4.47 (1.61,12.37)														
Oda H 2008 [86]	161	Cross-sectional	MRI	Bone bruise/effusion OR (95%CI)= 2.74 (1.31,5.72)												
Stein V 2010 [87]	160	Cross-sectional	MRI	Intecondylar notch/ACL tear OR (95%CI)= 1.00 (0.058,17.01)												

MRI: magnetic resonance imaging; JSN: joint space narrowing; T2: T2 relaxation time; sd: standard deviation; PF: patellofemoral; WORMS: whole organ magnetic resonance imaging score; LPTA: Lateral patellar tilt ratio; TF: tiobiofemoral; BME: bone marrow edema;KLG: Kellgren and Lawrence grade; JSN: joint space narrowing; JSW: joint space width; BMLs: bone marrow lesions; OR (95%CI): odds ratio and 95% confidence interval

Cross-sectional studies, US: all outcomes

Reference	No of subjects	Study design	Imaging modality	Outcome						
Song IH 2008 [8]	41	Cross-sectional	US (CEUS)	Effusion, synovitis, PD, slope/VAS pain	Slope		r=0.386; p=0.018			
					Superior recess PD		r=0.366; p=0.020			
Naredo E 2005 [88]	50	Cross-sectional	US	Effusion, meniscal displacement and popliteal cyst/symptoms	Effusion		OR (95% CI)= 18.4 (1.04,323.52)			
					medial meniscal displacement		OR (95% CI)= 32.83 (1.86,577.99)			
					Popliteal cyst		OR (95% CI)= 6.10 (108.71)			
D'Agostino M 2005 [89]	600	Cross-sectional	US	effusion/sudden knee pain	OR (95% CI) = 1.39 (1.008,1.93)					
De Miguel Mendieta E 2006 [90]	121	Cross-sectional	US	Effusion and popliteal cyst/pain	Suprapatellar effusion		OR (95%CI) = 6.5 (1.87,22.37)			
					Popliteal cyst		OR (95%CI) = 5.5 (1,31.05)			
Hall M 2014 [91]		Case-control	US	Effusion, synovitis and popliteal cyst/VAS pain and WOMAC pain			VAS pain		WOMAC pain	
					Effusion		r=0.3 p<0.01		r= 0.3 p<0.01	
					Synovitis		r= 0.3 p<0.01		r= 0.3 p<0.01	
					Popliteal cyst		r= 0.1 n.s.		r= 0.1 n.s.	
Iagnocco A 2010 [92]	82	Cross-sectional	US	US score/VAS pain and Lequesne Index	Associations between total US score and VAS pain (p=0.004) and Lequesne Index (p<0.0001)					
Wu P 2012 [93]	112	Cross-sectional	US	US/VAS pain at motion and rest, WOMAC pain	p		VAS pain (motion)	VAS pain (rest)	WOMAC pain	
					Suprapatellar synovitis		0.45	0.17	0.23	
					Suprapatellar effusion		0.008	0.20	0.55	
					Medial meniscal extrusion		0.19	0.40	0.27	
					Medial compartment synovitis		0.001	0.001	0.001	
					Lateral compartment synovitis		0.60	0.64	0.77	
					Marginal osteophytes		0.39	0.66	0.46	

US: ultrasonography; CEUS; contrast enhanced ultrasound; PD: power Doppler; VAS: visual analogue scale; OR (95%CI): odds ratio and 95% confidence interval; r: correlation coefficient; KOOS: Knee injury and Osteoarthritis Outcome Score; ADL: activities of daily living; QoL: quality of life; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index
 US: ultrasonography; CEUS; contrast enhanced ultrasound; PD: power Doppler; VAS: visual analogue scale; OR (95%CI): odds ratio and 95% confidence interval; r: correlation coefficient; KOOS: Knee injury and Osteoarthritis Outcome Score; ADL: activities of daily living; QoL: quality of life; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index

Cross-sectional studies, radioisotope scan: all outcomes

Reference	No of subjects	Study design	Imaging modality	Outcome				
				Mean, sd	symptomatic	asymptomatic	p	
Kim HR 2008 [94]	30	Cross-sectional	SPECT	SPECT score/symptoms				
				MT	1.37 (0.17)	0.21 (0.11)	<0.001	
				MF	0.91 (0.15)	0	<0.001	
				LT	0.43 (0.08)	0.29 (0.16)	>0.05	
				LF	0.41 (0.09)	0.07 (0.07)	0.006	
				Patella	1.26 (0.12)	0.43 (0.17)	0.001	
			Femoral trochlea	0.83 (0.12)	0.29 (0.16)	0.023		
Kraus VB 2009 [95]	159	Cross-sectional	radioisotope scan	Captation/severity of knee symptoms	Significant correlation between captation and symptoms severity p=0.0012			
McCrae F 1992 [96]	100	Cross-sectional	radioisotope scan	Abnormalities/pain	OR (95% CI)= 45.1 (6.07, 335)			
SPECT: single proton emission computed tomography; MT: medial tibia; MF: medial femur; LT: lateral tibia; LF: lateral femur; OR (95% CI): odds ratio and 95% confidence interval								

Longitudinal studies, CR: symptoms, joint replacement

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Bruyere O 2002 [97]	212	RCT	CR	3 years	JSW and mJSW/WOMAC		WOMAC pain	WOMAC stiffness	WOMAC function
						Baseline JSW	r = -0.10 p = 0.16	r = 0.12, p = 0.09	r = - 0.19, p = 0.11
						Baseline mJSW	r = - 0.08, p = 0.27	r = - 0.05, p = 0.51	r = - 0.12, p = 0.31
						Change in JSW	r = - 0.29, p = 0.017	r = - 0.22, p = 0.06	-
						Change in mJSW	r = - 0.24, p = 0.044	r = - 0.05, p = 0.67	-
Collins JE 2014 [98]	1753	Cohort	CR	6 years	KLG/pain trajectories		KL3 vs KL2	KL4 vs KL2	
						Mild	1.0 (0.6,1.6)	3.6 (0.9,15.0)	
						Low moderate	2.6 (1.7,4.1)	21.9 (6.1,79.1)	
						High moderate	2.6 (1.6,4.3)	25.5 (6.8,95.8)	
						Severe	4.3 (2.1,8.6)	80.6 (18.7,346.8)	
Conaghan PJ 2010 [99]	531	Cohort	US CR	3 years	KLG and effusion/joint replacement	KLG	HR (95%CI)= 4.08 (2.34 , 7.12)		
						US effusion	HR (95%CI)= 2.63 (1.70, 4.06)		
Podsiadlo P 2014 [100]	114	Cohort	CR	6 years	Osteophytes/TKR	grade of medial tibiofemoral osteophyte: OR (95%CI)= 2.0 (1.27, 3.13) p=0.003			

OR (95% CI): odds ratio and 95% confidence interval; JSW: joint space width; mJSW: minimal joint space width; r: correlation coefficient; KL/KLG: Kellgren and Lawrence grade; US: ultrasonography; CR: conventional radiography; TKR: total knee replacement; RCT: randomized controlled trial

Longitudinal studies, CR: structure

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Cahue S 2004[101]	217	Cohort	CR	18 months	Alignment/PF progression	Medial PF progression varus/non varus (OR, 95%CI)= 1.98 (1.15,3.43)		
						Lateral PF progression valgus/non valgus (OR 95% CI)= 1.25 (0.79–1.96)		
Cerejo R 2002[102]	230	Cohort	CR	18 months	Alignment/CR progression	KLG 0-1	KLG 2	KLG 3
					Varus/medial progression (OR 95% CI)	2.50 (0.67,9.39)	4.12 (1.92,8.82)	10.96 (3.10,38.77)
					Valgus/lateral progression (OR 95% CI)	-	2.46 (0.95,6.34)	10.44 (2.76,39.49)
Brouwer B 2007[103]	1501	Cohort	CR	6 years	Alignment/KLG	Varus (OR, 95%CI)	1.39 (0.48,4.05)	
						Valgus (OR, 95%CI)	2.90 (1.07,7.88)	
Felson DT 2009 [104]	160	Cohort	CR	1 year	Alignment/medial JSN progression	Varus vs neutral (OR, 95%CI)	0.33 (0.05, 2.24)	
						Valgus vs neutral (OR, 95%CI)	4.82 (1.93, 12.00)	
Felson DT 2005 [105]	270	Cohort	CR	30 months	Increase in osteohypte score/ radiographic progression	OR (95% CI)= 1.3 (1.0,1.7)		
Miyazaki T 2002 [106]	74	Cohort	CR	6 years	Alignment/mJSW	sensitivity 0.66, specificity 0.62		
Moisio K 2011 [107]	159	Cohort	CR		Alignment/MRI cartilage loss	Medial (OR 95% CI)	Lateral (OR 95% CI)	
					Neutral vs varus	0.49 (0.24,1.01)	-	
					Valgus vs varus	0.24 (0.10,0.58)	-	
					Neutral vs valgus	-	0.32 (0.16,0.66)	
					Varus vs valgus	-	0.41 (0.20,0.85)	

OR (95% CI): odds ratio and 95% confidence interval; CR: conventional radiography; PF: patellofemoral; KLG: Kellgren and Lawrence grade; JSN: joint space narrowing; mJSW: minimal joint space width

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome		
Sharma L 2001[108]	230	Cohort	CR	18 months	Alignment/progression	Medial progression varus/non varus	OR 95% CI= 5.00 (2.77,9.02)
						Lateral progression valgus/non valgus	OR 95% CI= 3.88 (1.82,8.24)
Cicuttini F 2004 [109]	117	Cohort	CR	2 years	Knee angle/MRI cartilage volume	Change in CV per degree of angle increase	p
						MT	0.16
						MF	0.004
						LT	0.05
						LF	0.57
Davies Tuck M 2008 [110]	100	Cohort	CR	2 years	Femoral sulcus angle/MRI patellar cartilage loss	No significant association between femoral sulcus angle and subsequent cartilage loss	
Eckstein F 2011 [111]	152	Cohort	CR MRI	2 years	CR and MRI/MRI cartilage thickness loss	predictors	AUC (p)
						mJSW (mm)	0.738 (0.0035)
						alignement	0.767 (0.0012)
						OP_score_cMF	0.637 (0.0163)
						JSN_score_MFT C	0.741 (0.0001)
						ThCtAB_MT (mm)	0.568 (0.0001)
						ThCtAB_cMF (mm)	0.805 (0.0193)
Bruyere O 2003[112]	105	Cohort	CR	3 years	Baseline JSW/JSN	r= -0.34, p=0.003	
Felson DT 2013[113]	5147	Cohort	CR	4 years	Radiographic progression in the previous year/radiographic progression	OR (95% CI)= 4. (2.4,6.7)	

CR: conventional radiography; MRI: magnetic resonance imaging; OR (95% CI): odds ratio and 95% confidence interval; CV: cartilage volume; MT: medial tibia, MF: medial femur; LT: lateral tibia; LF: lateral femur; AUC: area under the curve; mJSW: minimal joint space width; OP: osteophyte; cMF: centra medial femur; JSN: joint space narrowing; MFTC: medial femorotibial compartment; ThCtAB: mean cartilage thickness over the bone area; MT: medial tibia; r: correlation coefficient; ROI: region of interest

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Woloszynski T 2012[114]	68	Cohort	CR (fractal analysis)		Fractal signature analysis/radiographic progression	AUC (95% CI) for radiographic progression: Medial ROI 0.77 (0.68, 0.86); Lateral ROI 0.71 (0.63, 0.79)		
Kraus VB 2009[115]	138	Cohort	CR	3 years	Fractal signature analysis/radiographic progression	AUC 0.74 (0.65–0.84)		
Mazzuca S 2005 [116]	174	Cohort	CR radioisotope scan	30 months	KLG and bone uptake/radiographic progression	Se	Sp	
						MT uptake (50th percentile) Baseline KLG	0.57 0.65	0.60 0.64
Mazzuca S 2006[117]	319	Cohort	CR	30 months	CR/radiographic progression (JSN and osteophytes)	JSN (OR 95%CI)	Osteophytes (OR 95% CI)	
						JSW	0.67 (0.49, 0.91)	-
						Controlateral OA	1.53 (0.82, 2.85)	-
					Ipsilateral PF OA	3.01 (1.63, 5.57)	2.31 (1.37, 3.88)	
Bijsterbosch J 2014[118]	236	Cohort	CR		Hand OA/knee OA progression	hand OA with baseline knee OA 2.5 (0.9 to 6.9); hand OA without baseline knee OA 2.7 (1.3 to 5.9)		
Neogi T 2006 [119]	495	Cohort	CR		Chondrocalcinosis/MRI cartilage loss	RR (95% CI)= 0.4 (0.2,0.7)		
Buck RJ 2013 [120]	869	Cohort	CR	1 year	KLG/MRI cartilage loss	cKLG moderate+severe vs minimal+doubtful OR (95% CI) 2.49 (1.46,4.26)		
Wirth W 2014 [121]	922	Cohort	CR	1 year	JSN/change in cartilage thickness	LFTC (mean, sd)	MFTC (mean, sd)	
						No JSN	-8 (84)	-2 (87)
						Lateral JSN	-81 (168)	15 (117)
					Medial JSN	-20 (100)	-63 (154)	

CR: conventional radiography; MRI: magnetic resonance imaging; AUC: area under the curve; Se: sensitivity; Sp: specificity; KLG: Kellgren and Lawrence grade; MT: medial tibia; JSN: joint space narrowing; OR (95% CI): odds ratio and 95% confidence interval; LFTC: lateral femorotibial compartment; MFTC: medial femorotibial compartment; JSW: joint space width, BML: bone marrow lesions; r: correlation coefficient

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Pelletier J 2007 [122]	110	Cohort	CR MRI	2 years	CR and MRI/medial cartilage loss and JSW		Cartilage loss	JSW
						JSW	r= 0.32 p=0.001	-
						Meniscal extrusion	r= -0.32 p=0.001	r= -0.32 p=0.001
						Medial meniscal tear	r= -0.21 p=0.03	r= -0.32 p=0.001
						Lateral meniscal tear	r= 0.15 p=0.13	r= -0.09 p=0.33
						BMLs	r= -0.27 p=0.005	r= 0.11 p=0.25
Sharma L 2008 [123]	153	Cohort	CR MRI	2 years	MRI/medial cartilage loss		Cartilage volume (MT)	Cartilage thickness (MT)
						Medial meniscal damage (OR 95%CI)	1.57 (1.29, 1.91)	1.40 (1.16, 1.69)
						Medial meniscal extrusion (OR 95%CI)	1.99 (1.36, 2.91)	1.81 (1.23, 2.65)
						Varus (OR 95%CI)	1.16 (1.07, 1.26)	1.20 (1.10, 1.31)
Wang Y 2005 [124]	132	Cohort	CR	2 years	JSN and osteophytes/change in plateau area		Medial plateau (p)	Lateral plateau (p)
						JSN	0.98	0.24
						Osteophytes	0.02	0.25
						Basal area	<0.001	0.11
Cicuttini F 2005 [125]	28	Cohort	CR MRI	2 years	JSW/cartilage loss Cartilage loss and JSN/joint replacement	JSW and cartilage volume r= 0.35 (p=0.06)	loss of medial TF cartilage volume to predict joint replacement at 4 years: OR=9.0, P<0.07 change in JSW: OR=1.1, P<0.92	
Tanamas S 2009 [126]		SLR	CR		Alignment/radiographic progression	No quantitative synthesis, references were reviewed		

CR: conventional radiography; MRI: magnetic resonance imaging; OR (95% CI): odds ratio and 95% confidence interval; MT: medial tibia; JSN: joint space narrowing; JSW: joint space width; r: correlation coefficient; TF: tibiofemoral; SRL: systematic literature review

Longitudinal studies, MRI: joint replacement

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Cicuttini F 2004 [127]	123	Cohort	MRI	2 years	Cartilage volume loss/joint replacement	Rate of cartilage loss 3-8% year OR (95%CI)= 2.3 (0.4, 12.2)			
						Rate of cartilage loss >8% year OR (95%CI)= 7.1 (1.4, 36.5)			
Eckstein F 2013 [128]	110	Case-control	MRI	1 year	Change in cartilage thickness/joint replacement	Longitudinal change in cMFTC 1 year before TKA and immediately before TKA was greater in TKA case AUC=0.59 (0.52,0.57)			
Raynauld J 2011[129]	123	RCT	CR MRI	4-7 years	CR and MRI/joint replacement	OR (95%CI)			
						JSW	0.579 (0.42, 0.79)		
						Medial meniscal tear	5.687 (1.75, 18.49)		
						Medial meniscal extrusion	4.058 (1.34, 12.23)		
					BMLs score	1.196 (1.04, 1.37)			
Wluka AE 2005 [130]	117	Cohort	MRI	2 years	Cartilage defects/ annual cartilage loss and joint replacement	Cartilage defects	No cartilage defects	p	
						Lateral TF	4.6% (3.2%, 5.9%)	5.8% (3.2%, 8.4%)	0.42
						Medial TF	4.7% (3.1%, 6.4%)	6.5% (4.3%, 8.6%)	0.21
						Joint replacement	OR (95%CI) = 6.0, (1.6, 22.3)		
Wang Y 2012 [131]	107	Cohort	MRI	2-4.5 years	Vastus medialis area and cartilage volume/WOMAC pain, radiographic progression, joint replacement	inverse association between vastus medialis CSA and WOMAC knee pain	Radiographic progression: Baseline tibial cartilage volume Medial p=0.001; Lateral p<0.001	Knee replacement over 4 years (change in vastus medialis area between 0 and 2 years) 0.62 (0.41, 0.96) OR 95% CI	
Tanamas S 2010 [132]	88	Cohort	MRI	4 years	BMLs and cysts/joint replacement	Medial compartment OR 95%CI 1.72 (0.93, 3.18) Lateral compartment 0.95 (0.48, 1.88)			
Scher C 2008 [133]	65	Cohort	MRI	3 years	BMLs/joint replacement	OR 95% CI 5.22 (1.08,25.08)			
Eckstein F 2009 [134]	156	Cohort	MRI	1 year	Denuded areas and cartilage thickness/cartilage loss	denuded area in MFTC : significantly greater cartilage loss in MT (P < 0.01) and cMT (P< 0.001), MF(P < 0.05) and cMF (P < 0.001); smaller baseline cartilage thickness in MT: greater cartilage loss in MT (P < 0.05) and cMT (P < 0.05); smaller baseline cartilage thickness in cMF : greater cartilage loss in cMF (P < 0.05)			

CR: conventional radiography; MRI: magnetic resonance imaging; OR (95% CI): odds ratio 95% confidence interval, sd: standard deviation ; cMFTC: central medial tibiofemoral compartment; TKA: total knee arthroplasty; AUC: area under the curve; JSW: joint space width; BMLs: bone marrow lesions; TF: tibiofemoral; WOMAC: Western Ontario and MacMAster Universities Osteoarthritis Index; CSA: cross-sectional area; MFTC: medial tibiofemoral compartment; MT: medial tibia; cMT: central medial tibia; cMF: central medial femur; RCT: randomized controlled trial.

Longitudinal studies, MRI: symptoms and function

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Amin S 2008 [135]	265	Cohort	MRI	30 months	ACL tears/VAS pain, cartilage loss, WOMAC function		Complete tear	No complete tear
						VAS pain (sq mean 95% CI)	45.3 (38.1, 52.5)	39.8 (36.4, 43.2)
						Cartilage loss OR(95% CI)	1.1 (0.6, 1.8) (medial), 1.0 (0.4, 2.2) lateral, 0.8 (0.5, 1.4) PF	
					WOMAC function (sq mean 95% CI)	25.8 (22.3, 29.3)	22.3 (21.8, 25.2)	
Everhart J 2014 [136]	1433	Cohort	MRI	4 years	SSR/incident symptoms, radiographic progression		Incident symptoms (OR 95% CI)	Radiographic progression
						Medial SSR	0.48 (0.30, 0.75)	0.91 (0.55, 1.24)
						Lateral SSR	1.27 (0.86, 1.88)	0.89 (0.69, 1.14)
Moio K 2009 [137]	182	Cohort	MRI	2 years	Denuded areas/incident pain	No significant association between baseline denuded areas and incident pain		
Pelletier J 2008 [138]	27	Cohort	MRI	2 months	Synovitis score/VAS pain, WOMAC, MRI	WOMAC pain 0.07 p=0.71; WOMAC stiffness 0.10 p=0.63; VAS pain 0.01 p=0.93	Cartilage volume loss 0.20 p=0.34; BME score 0.04 p=0.83	WOMAC function 0.20 p=0.34
Hill CL 2007 [139]	270	Cohort	MRI	30 months	Change of synovitis and change of effusion/pain and cartilage loss	Change of synovitis: r=0.21, p=0.0003 Change of effusion: n.s.		
						Baseline synovitis/ cartilage loss in the medial compartment 2.9 (1.3, 6.5). Not significant for lateral and PF compartments. No significant relationship between change in score and cartilage loss		

MRI: magnetic resonance imaging; ACL: anterior cruciate ligament; VAS: visual analogue scale; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index; OR (95% CI): odds ratio and 95% confidence interval; SSR: subchondral bone surface; r: correlation coefficient; PF: patellofemoral

Longitudinal studies, MRI: structure

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Berthiaume M 2005 [140]	32	Cohort	MRI	2 years	Meniscal extrusion/cartilage volume loss Anterior horn extrusion p=0.01; Middle horn extrusion p=0.02; Meniscal tear p=0.83; Meniscal degeneration p=0.94.			
Chang AM 2011 [141]	159	Cohort	MRI	2 years	meniscal tears/cartilage loss	Medial meniscal tear	body tear: central subregion 4.18 (1.80, 9.72); External subregion 5.39 (2.44, 11.88). Anterior subregion 3.22 (1.49, 6.96); Posterior horn, posterior subregion 2.63 (1.23, 5.62)	
						Lateral meniscal tear	body tear : Central subregion 3.81 (1.12, 13.00); External subregion 3.99 (1.01, 15.85); Anterior subregion 3.24 (1.02, 10.35); Posterior subregion 6.47 (1.74, 24.05); Posterior horn 3.94 (1.26, 12.38)	
Crema MD 2014 [142]	148	Cohort	MRI (dGEMRIC)	1 year	Meniscal pathology/dGEMRIC	Regions with meniscal pathology had significantly lower dGEMRIC compared to those without meniscal pathology both cross-sectionally and longitudinally		
Wang Y 2010 [143]	126	Cohort	MRI	2 years	Medial meniscal extrusion/increase in BMLs and cysts	Increase in medial TF BMLs OR 95% CI= 3.3 (1.2, 9.1) Increase in medial TF cysts OR 95% CI= 6.8 (2.0, 23.8)		
Raynauld J 2006 [144]	110	Cohort	MRI	2 years	Meniscal extrusion, cartilage volume, BMLs/cartilage loss	Severe medial meniscal extrusion -0.28 (p= 0.08); Medial compartment cartilage volume -0.36 (p= 0.012); Bone oedema -0.30 (p= 0.06)		
Hunter DJ 2006 [145]	257	Cohort	MRI	30 months	Meniscal damage and position/cartilage loss	OR (95%CI)	Medial compartment	Lateral compartment
						subluxation: 3-4 mm	0.9 (0.4,1.9)	1.7 (0.6,5.0)
						5-6 mm subluxation	3.2 (1.5,6.9)	3.0 (1.3,7.0)
						≥7 mm subluxation	2.4 (1.1,5.0)	4.6 (2.0,10.8)
Crema MD 2014 [146]	163	Cohort	MRI	3 years	subchondral sclerosis/cartilage loss	OR 95% CI		
						MF	1.7 (0.5, 5.9)	
						MT	1.87 (0.,7.1)	
						LF	0.5 (0.1,3.8)	
						LT	0.7 (0.1,8.2)	
			patella	2.1 (0.5,8.5)				

MRI: magnetic resonance imaging; dGEMRIC: delayed gadolinium contrast enhanced magnetic resonance imaging; TF: tibiofemoral; BMLs: bone marrow lesions; OR (95% CI): odds ratio and 95% confidence interval

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Eckstein F 2009 [147]	80	Cohort	MRI	1 year	JSN/cartilage thickness	No significant differences between knees with more JSN compared to those with less JSN			
Driban JB 2011 [148]	44	Cohort	MRI	2 years	BMLs volume/cartilage lesions		Cartilage volume	Cartilage thickness	Full-thickness lesions
						Femur BMLs	r= 0.03 (-0.29, 0.34)	r= -0.11 (-0.41, 0.21)	r= 0.48 (0.20, 0.69)
						Tibial BMLs	r= 0.16 (-0.18, 0.46)	r= -0.13 (-0.43, 0.21)	r= 0.43 (0.12, 0.66)
Carrino JA 2006 [149]	32	Cohort	MRI	1.5 years	BME/subchondral cysts	11/12 newly developed subchondral cysts developed in areas of previous BME			
Hunter DJ 2006 [150]	217	Cohort	MRI	30 months	BMLs/cartilage loss	Medial TF p<0.0001, lateral TF cartilage p<0.0001			
Hernandez Molina G 2008 [151]	258	Cohort	MRI	30 months	BMLs/medial cartilage loss and ACL tears	Cartilage loss (OR 95% CI) 1.6 (0.8,3.1); ALC tears 5.9 (2.2,16.2)			
Felson DT 2003 [152]	223	Cohort	MRI	30 months	BMLs/radiographic progression	Lateral BMLs	OR (95% CI) 6.5 (3.0,14.0)		
						Medial BMLs	OR (95% CI) 6.1 (2.2,16.5)		
Kothari A 2010 [153]	177	Cohort	MRI	2 years	BMLS, subchondral cysts and bone attrition/cartilage loss in the same subregion		OR (95% CI)		
						BMLs	3.74 (1.59,8.82)		
						Subchondral cysts	0.47 (0.11,2.03)		
						Bone attrition	1.85 (0.71,4.82)		
Felson DT 2010 [154]	113	Cohort	CR MRI	2 years	Alignment and BMLs/WORMS and BLOKS cartilage score		BLOKS medial cartilage (OR 95% CI)	WORMS medial cartilage (OR 95% CI)	
						alignment	5.9 (1.5,24.0)	2.1 (0.7,6.3)	
						BMLs	1.9 (0.5,7.2)	5.5 (1.2, 25.4)	
MRI: magnetic resonance imaging;MF: medial femur, MT: medial tibia; LF: lateral femur; LT: lateral tibia; JSN: joint space narrowing; BMLs: bone marrow lesions; BME: bone marrow lesions; r: correlation coefficient; TF: tibiofemoral; ACL: anterior cruciate ligaments; OR (95% CI): odds ratio and 95% confidence interval									

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome		
Wluka AE 2002 [155]	123	Cohort	MRI	2 years	Initial cartilage volume/cartilage volume loss	Initial cartilage volume predicted subsequent cartilage volume loss p=0.01	
Madan-Sharma L 2008 [156]	186	Cohort	MRI	2 years	BMLs, cysts, cartilage loss, osteophytes, meniscal tears, meniscal subluxation, joint effusion/JSN progression	OR (95% CI)	
						BMLs	1.8 (0.51,5.17)
						Subchondral cysts	2.7 (0.8,6.4)
						Cartilage loss	1.7 (0.4,6.7)
						Osteophytes	2.1 (0.6,6.4)
						Meniscal tears	3.57 (1.1–10.0)
Meniscal subluxation	2.73 (1.2,5.4)						
effusion	1.35 (0.7,2.4)						
Yusuf E 2011 [157]		SLR	MRI				

MRI: magnetic resonance imaging ; CR: conventional radiography; BMLs: bone marrow lesions; BLOKS: Boston Leeds osteoarthritis knee score; WOMRS: whole organ magnetic resonance imaging score; OR (95% CI): odds ratio and 95% confidence interval; cMF: central medial femur; JSN: joint space narrowing

Longitudinal studies, US and radioisotope scan

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome
Dieppe P 1993 [158]	94	Cohort	Radioisotope scan	5 years	Radiographic progression OR (95% CI)= 1.86 (0.82,4.23)

US: ultrasonography; OR (95% CI): odds ratio and 95% confidence interval

Q6: What is the prognostic (prediction of therapeutic response) value of imaging in OA?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Barrett JP 1990 [159]	248	Cohort	CR	6 months	Radiographic severity/response to intraarticular HA		Pain on walking	Pain at rest	Pain at night
						Radiographic grade 2	r=0.80	r= 0.67	r= 0.42
						Radiographic grade 3	r= 0.08	r= 0.14	r= -0.20
Gaffney K 1995 [160]	84	RCT	CR	6 weeks	OA severity 0-3/response to intraarticular trimacynolone vs placebo	No association between improvement in VAS pain and radiographic score			
Gudbergesen H 2012 [161]	192	RCT	CR MRI	16 weeks	mJSW, alignment and MRI scores/pain reduction in response to very low energy diet vs low energy diet	mJSW r =-0.02 (p =0.81); KL r =-0.07 (p = 0.29); r = 0.08 (p= 0.27); cartilage score r =-0.09 (p= 0.24); BML score r =-0.04 (p= 0.62); r =-0.08 (p= 0.29); effusion score r = 0.02 (p= 0.72); synovitis score r = 0.01 (p=0.94); meniscal extrusion score medial r =0.09; (p = 0.23) lateral r = 0.04 (p= 0.60); meniscal pathology score medial r = 0.04 (p = 0.60) Lateral r =-0.06 (p= 0.44)			
Gudbergesen H 2011 [162]	30	RCT	CR MRI	32 weeks	KLG and MRI score/change in WOMAC pain and function during weight reduction		Change in WOMAC pain	Change in WOMAC function	
						KLG	r = 0.13; (p= 0.50)	r = 0.03; (p= 0.88)	
						MRI score	r = 0.19; (p = 0.31)	r = -0.003; (p= 0.98)	
Hellio le Graverand M 2013 [163]	1452	RCT	CR	96 weeks	KLG/structural progression in patients treated with cidunistat or placebo	the loss of JSW after 48weeks in KLG2 patients was smaller with cindunistat 50 mg/day than with placebo (least squares (LS)-mean vs placebo (95% CI): 0.072 (0.006 to 0.138); p=0.032), but did not differ for cindunistat 200 mg/day versus placebo. No significant differences at 96 weeks.			

MRI: magnetic resonance imaging; CR: conventional radiography; US: ultrasonography; mExt: meniscal extrusion; BMLs: bone marrow lesions; Str Ran: strontium ranleate; LB: placebo; r: correlation coefficient; HA: hyaluronic acid; VAS: visual analogue scale; mJSW: minimal joint space width; KL/ BML: bone marrow lesions; WOMAC: Western Ontario and MacMAster Universities Osteoarthritis Index; r: correlation coefficient; RCT: randomized controlled trial

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome							
Keurentjes J 2013 [164]	723	Cohort	CR	5 years	KLG/improvement of Qol after knee replacement		KLG 0-2 vs KLG 3-4 Md (95% CI)	p				
						Physical functioning	19.1 (8.48–29.7)	<0.001				
						Role physical	17.4 (26.32–41.1)	0.15				
						Bodily pain	9.02 (23.43–21.5)	0.15				
						General Health	9.23 (1.31–17.2)	0.02				
						Vitality	8.44 (20.28–17.2)	0.06				
						Social functioning	7.44 (24.18–19.1)	0.21				
						Role emotional	8.87 (211.8–29.6)	0.40				
						PCS	5.64 (1.26–10.0)	0.01				
	MCS	20.18 (24.45–4.10)	0.94									
	NRS satisfaction	1.2 (0.1–2.4)	0.04									
Case JP 2003 [165]	82	RCT	CR	12 weeks	KLG and medial JSN/WOMAC response to diclofenac vs paracetamol		WOMAC pain (p)		WOMAC stiffness		WOMAC function	
							Dicl/par	Dic/plb	Dicl/par	Dic/plb	Dicl/par	Dic/plb
						KLG 1-2	0.002	0.003	<0.001	0.002	<0.001	0.002
						KLG 3-4	0.85	0.37	>0.999	>0.999	0.28	0.48
						JSN 0-1	0.005	0.002	<0.001	<0.001	0.002	<0.001
	JSN 2	0.63	0.29	0.99	0.99	0.41	0.70					
Sawitzke A 2008 [166]	375	RCT	CR	24 months	KLG/radiographic progression during treatment with glucosamine, chondroitin sulphate and celecoxib	OR for radiographic progression compared with the placebo group was <1 in patients with K/L grade 2 knees in all treatment groups, whereas it was >1 in patients with K/L grade 3 knees in all treatment groups						
Mazzuca S 2010 [167]	379	RCT	CR	30 months	Alignment/radiographic progression in doxycycline vs placebo	varus knees exhibited a greater loss of JSW than non-varus knees						
CR: conventional radiography; KLG: Kellgren and Lawrence scale; JSW: joint space width; MD: mean difference; PCS: physical component score; MCS: mental component score; NRS: numerical rating scale; JSN: joint space narrowing; WOMAC: Western Ontario and MacMaster Universities Osteoarthritis Index; Dic: diclofenac, plb: placebo; OR: odds ratio; RCT: randomized controlled trial.												

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome		
Toh E 2002 [168]	60	Cohort	CR	12 weeks	Alignment, sclerosis, cysts, osteophytes, JSN/WOMAC response to intraarticular HA	OR (95% CI)	
						Alignment	0.91 (0.29,2.80)
						Subchondral sclerosis	0.97 (0.32,2.93)
						Subchondral cysts	2.94 (0.89,9.67)
						osteophytes	0.41 (0.13,1.26)
						Lateral JSN	0.17 (0.05,0.619)
						Medial JSN	0.037 (0.008,0.15)
Pendleton 2008 [4]	86	Cohort	US	6 weeks	US/WOMAC response to intraarticular methylprednisolone	Higher baseline US scores: significant improvements in outcome at both week 1 (p<0.01 for WOMAC pain and function, p<0.05 for stiffness) and week 6 (p<0.01 all subscales)	
Chao J 2010 [169]	67	RCT	US	12 weeks	US inflammation/WOMAC response to triamcinolone	statistically significant improvement in pain subscales among non-inflammatory patients than among inflammatory patients at 12 weeks (p=0.03)	
Hirsch G 2013 [170]		SLR				No quantitative synthesis, references were reviewed	
Anandacoomara samy A 2008 [171]	32	Cohort	MRI	6 months	Cartilage volume/response to intraarticular HA	no correlation between baseline MRI measures and clinical response	
Knoop JD 2014 [172]	91	Cohort	MRI	12 weeks	MRI/change in WOMAC function in response to exercise program	Cartilage integrity TF p=0.92, PF -0.25 p=0.01; BMLs TF p=0.28; PF -p=0.45; osteophytes TF p=0.88 PF p=0.83; Inflammation -p=0.45; Meniscal abnormalities p=0.51	
CR: conventional radiography; JSN: joint space narrowing; WOMAC: Western Ontario and MacMAster Universities Osteoarthritis Index; HA: hyaluronic acid; US: ultrasonography; NSAIDs: non-steroidal anti-inflammatory drugs; Glu: glucosamine; CS: chondroitin sulphate; MRI: magnetic resonance imaging; TF: tibiofemoral; BMLs: bone marrow lesions; PF: patellofemoral.							

FOLLOW-UP OF OA

Q7: When, how often and under what clinical circumstance should imaging be used in the follow-up of osteoarthritic joints?

CR: sensitivity to change

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Pessis E 2003 [173]	20	Cohort	CR MRI	1 year	Sensitivity to change JSW and MRI score SRM: JSW -0.11 MRI score 1.03			
Wirth W 2014 [121]	922	Cohort	MRI	1 year	Sensitivity to change ThCtAb depending on JSN	SRM	Medial compartment	Lateral compartment
						No JSN	-0.02	-0.09
						JSN	Medial	-0.41
						Lateral	-0.13	-0.48
Conrozier T 2005 [174]	160	Cohort	CR	1 year	Sensitivity to change mJSW SRM: 0.40			
Hellio Le Graverand MP 2008 [175]	62	Cohort	CR	1 year	Sensitivity to change of mJSW	SRM	Lyon Schuss	Fixed flexion
						KLG1	0.51	0.0
						KLG2	0.0	0.0
						KLG3	0.0	0.01
						KLG4	0.65	0.01
Mazzuca S 2008[176]	74	Cohort	CR	1 year	Sensitivity to change of mJSW • Lyon Schuss: SRM 0.46 • Fixed flexion: SRM 0.05			
Piperno M 1998[177]	44	Cohort	CR	1 year	Sensitivity to change of mJSW	Lyon Schuss	Full extension	
						KLG2	3.08	2.5
						KLG3	2.5	2.3
Spector TD 2005 [178]	231	RCT	CR	1 year	Sensitivity to change of medial mJSW placebo -0.28 Risedronate 5 -0.18 Risedronate 15 0.24			
Mazzuca S 2003 [179]	86	Cohort	CR	14 months	Sensitivity to change of medial mJSW SRM -0.29			
Botha-Scheepers S 2007 [180]	378	Cohort	CR	2 years	Sensitivity to change of mJSW SRM 0.56			

CR: conventional radiography; MRI: magnetic resonance imaging; JSW: joint space width; SRM: standardized response mean; JSN: joint space narrowing; ThCtAb: mean cartilage thickness over the bone area; mJSW: minimal joint space width; KLG: Kellgren and Lawrence grade

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Cicutini F 2004 [181]	102	Cohort	CR MRI	2 years	Sensitivity to change of cartilage volume and JSN at the patellofemoral		Lateral view	Skyline view	
						Midpatellar JSW	0.05	0.13	
						mJSW	0.02	0.3	
						Cartilage volume	0.96		
Hellio Le Graverand MP 2010 [182]	55	Cohort	CR MRI	2 years	Sensitivity to change of cartilage thickness and mJSW		Cartilage thickness	Lyon Schuss	Fixed flexion
						KL2	-0.15	-0.32	-0.62
						KL3	-0.48	-0.62	-0.20
Hellio Le Graverand MP 2013 [163]	60	RCT	CR MRI	2 years	Sensitivity to change of cartilage volume, JSW and JSA	JSA	JSW	CV medial tibia	CV medial femur
						-0.30	-0.27	-0.45	-0.15
Mazzuca S 2001 [183]	255	Cohort	CR	2-3 years	Sensitivity to change of JSN	SRM 0.29			
Pavelka K 2004 [184]	89	Cohort	CR	2 years	Sensitivity to change of mJSW	SRM -0.50			
Vignon E 2003 [185]	32	Cohort	CR	2 years	Sensitivity to change of mJSW	SRM: Standing AP 0.23; Lyon Schuss 0.48			
Wirth W 2013 [186]	445	Cohort	CR MRI	2 years	Sensitivity to change of mJSW and cartilage thickness	SRM: MFTC -0.43; mJSW -0.31			
Boegard T 2003 [187]	55	Cohort	CR	25 months	Sensitivity to change of mJSW	medial MJS: md= 0.056 (-0.070, 0.182), p=0.37 lateral MJS: md=-0.080 (-0.226, 0.067), p=0.28 PF joint: md= 0.019 (-0.16, 0.20), p=0.84			
Mazzuca S 2004 [188]	86	Cohort	CR	16 and 30 months	Sensitivity to change of JSN	SRM 16 months: 0.34 SRM 30 months 0.48			
Miyazaki T 2002 [189]	174	Cohort	CR	6 years	Sensitivity to change of mJSW	SRM 1.16			
Bruyere O 2003 [190]	212	RCT	CR	3 years	Sensitivity to change of mJSW	SRM (computer assisted) 0.44 SRM (manual) 0.40			
Gossec L 2008 [191]	100	Cohort	CR	30 months	Sensitivity to change of KLG, OARSI JSN and JSW	SRM	Full extension	Semiflexed	
						KLG	0.10	0.22	
						OARSI JSN	0.15	0.34	
						JSW	0.20	0.49	
CR: conventional radiography; MRI: magnetic resonance imaging; JSW: joint space width; SRM: standardized response mean; JSN: joint space narrowing; JSA: joint space area; CV: cartilage volume; MFTC: medial femorotibial compartment; MJS: medial joint space; PF: patellofemoral									

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Lanyon P 1996 [192]	108	Cohort	CR	31 months	Sensitivity to change of PF mJSW		Skyline view	Lateral view
						Mean change (sd) in JSW	-0.5 (-0.1,-0.8) lateral facet -0.5 (-0.2,-0.8) medial facet	-0.2(0.1,-0.5)
La Valley P 2005 [193]	482	Cohort	CR	30 months	Sensitivity to change of JSN	No change 0%; Change on both views 43%; Change on PA view 23% ; Change on lateral view 34%		
Nevitt M 2007 [194]	153	Cohort	CR	37 months	Sensitivity to change of mJSW	SRM 0.41		
Reginster JY 2001[195]	212	RCT	CR	3 years	Sensitivity to change of mJSW	Mean (sd) change in mJSW: placebo -0.31 (-0.57, -0.04) Glucosamine 0.07 (-0.17, 0.32)		
Sugiyama S 2003 [196]	172	Cohort	CR	4 years	Sensitivity to change of mJSW	SRM 1.23		
Reichmann W 2011 [197]		SLR	CR		Sensitivity to change of JSW	SRM 0.33 (0.26, 0.41)		
Duryea J 2010 [198]	150	Cohort	CR MRI	1 year	Sensitivity to change of JSW, cartilage volume and cartilage thickness	SRM: mJSW -0.15; JSW -0.31; MT volume -0.10; LT volume -0.22; MF volume -0.39; LF volume -0.06; MT thickness -0.04; LT thickness -0.22; MF thickness -0.34; LF thickness -0.13		

CR: conventional radiography; MRI: magnetic resonance imaging; JSW: joint space width; SRM: standardized response mean; JSN: joint space narrowing; mJSW: minimal joint space width; sd: standard deviation; MT: medial tibia; LT: lateral tibia; MF: media femur; LF: lateral femur; SLR: systematic literature review

CR: structural progression

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome		
Bartlett SJ 2011 [199]	312	Cohort	CR	1-2 years	Evaluation of trajectories of JSN progression	<ul style="list-style-type: none"> • 71% patients presented stability over 2 years • 7% moderate progression • 20% low progression • 2% rapid progression 	
Bruyere O 2007 [200]	62	Cohort	CR MRI	1 year	Correlation between JSN and WOMMS progression	<ul style="list-style-type: none"> • correlation between the progression of the WOMMS and radiographic medial JSN: (r=0.35, p=0.006) • ROC measuring the value of WOMMS change over 1 year for the prediction of JSN >0.3 mm AUC (95%CI) 0.73 (0.56,0.89, p=0.013) 	
Cicuttini F 2005 [125]	28	Cohort	CR MRI	2 years	Correlation between cartilage volume and JSW changes	r= -0.11 (p=0.58)	
Crema MD 2014 [201]	267	Cohort	CR MRI	30 months	Correlation between MRI features and JSN progression	OR (95%CI)	
						Worsening of cartilage damage	4.9 (2.6, 9.3)
						Worsening of meniscal damage	4.8 (1.8, 13.0)
						Worsening of meniscal extrusion	5.1 (1.5, 17.0)
Fukui N 2010 [202]	68	Cohort	CR	36 months	Correlation between radiographic progression and pain and function scores	Baseline JKOM (pain and function) score similar for progressors and non progressors. End of follow-up: the score for non progressors declined, while that for the subjects with progressed joints remained high	
Teichtahl A 2009 [203]	78	Cohort	CR	2 years	Correlation between TF cartilage loss and alignment	Annual change in cartilage volume per degree of angle increase: medial tibial cartilage volume (%) -0.44 (-0.95, -0.04) p=0.03 Lateral tibial cartilage volume (%) 0.01 (-0.4, 0.4) p=0.95	
Teichtahl A 2008 [204]	99	Cohort	CR	2 years	Correlation between PF cartilage loss and alignment	Annual change of knee angle for annual change in total patella cartilage volume -23.4 (-38.7, -8.1) p= 0.0003	

CR: conventional radiography; MRI: magnetic resonance imaging; JSN: joint space narrowing; WOMMS: whole organ magnetic resonance imaging score; OR (95% CI) odds ratio and 95% confidence interval; JSW: joint space width; JKOM: Japanese knee osteoarthritis measure; TF: tibiofemoral; PF: patellofemoral.

MRI: sensitivity to change

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome					
Eckstein F 2009 [134]	156	Cohort	MRI	1 year	Sensitivity to change of cartilage volume and thickness	SRM	MFTC	LTFC		
						CV	-0.21	-0.12		
						ThCtAb	-0.31	-0.11		
Eckstein F 2009 [147]	156	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	cMF -0.50; ccMF -0.45 ; CMTFC -0.47				
Eckstein F 2008 [205]	158	Cohort	MRI	3 months	Sensitivity to change of cartilage volume and thickness	<ul style="list-style-type: none"> • Mean differences in ThCtAB (3 month vsbaseline) were +0.13% in MT, +0.09% in cMF, +0.58% in LT and +0.98% in cLF • ThCtAB between the first and second measurement was not significant in MT and cMF • The tAB increased significantly in all cartilage plates 				
Eckstein F 2009 [206]	80	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness according to the presence of baseline JSN	SRM	MFTC	MT	MF	cMF
						JSN 0-1	-0.23	-0.28	-0.13	-0.19
						JSN 2-3	-0.46	-0.45	-0.25	-0.37
Blumenkrantz G 2004 [207]	38	Cohort	MRI	2 years	Sensitivity to change of cartilage thickness and volume	Mean change (sd)		KLG 1-2	KLG 3-4	
						lateral condyle cartilage thickness		-10.5 (11.0)	-9.4 (11.2)	
						lateral condyle cartilage volume		-7.0 (5.8)	-5.3 (7.3)	
						Medial condyle cartilage thickness		-7.7 (10.3)	-2.5 (10.4)	
Medial condyle cartilage volume		-10.2 (14.2)	-5.4 (14.5)							
Eckstein F 2011 [208]	719	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	SRM	MFTC	LTFC		
						KLG2	-0.11	-0.13		
						KLG3	-0.31	-0.23		
						KLG4	-0.38	-0.39		
Hunter DJ 2009 [209]	150	Cohort	MRI	1 year	Sensitivity to change of cartilage volume and normalized cartilage volume	SRM	Cartilage volume	Normalized Cartilage volume		
						Femur	-0.10	-0.09		
						Lat tibia	-0.24	-0.26		
						Med tibia	-0.04	-0.01		
						Patella	-0.18	-0.16		
Trochlea	-0.006	-0.01								

MRI: magnetic resonance imaging; SRM: standardized response mean; MFTC: medial tibiofemoral compartment; LTFC: lateral tibiofemoral compartment; CV: cartilage volume; ThCtAb: mean cartilage thickness over the bone surface; MT: media tibia; cMF: central medial femur; JSN: joint space narrowing; MF: medial femur; KLG: Kellgren and Lawrence grade

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome														
Hunter DJ 2010 [210]	150	Cohort	MRI	1 year	Sensitivity to change of cartilage volume SRM: -0.39														
Maschek S 2014 [211]	724	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	<table border="1"> <tr> <td></td> <td>KL2</td> <td>KL3</td> <td>KL4</td> </tr> <tr> <td>cartilage thickness SRM</td> <td>-0.14</td> <td>-0.35</td> <td>-0.50</td> </tr> </table>		KL2	KL3	KL4	cartilage thickness SRM	-0.14	-0.35	-0.50					
	KL2	KL3	KL4																
cartilage thickness SRM	-0.14	-0.35	-0.50																
Cromer T 2014 [212]	23	RCT	CR MRI	1 year	Sensitivity to change	<table border="1"> <tr> <td colspan="2">MJSW (SRM)</td> <td>-0.46</td> </tr> <tr> <td rowspan="2">Cartilage volume</td> <td>Medial tibia</td> <td>-0.56</td> </tr> <tr> <td>Medial femur</td> <td>-0.74</td> </tr> <tr> <td rowspan="2">Cartilage thickness</td> <td>Medial tibia</td> <td>-0.61</td> </tr> <tr> <td>Medial femur</td> <td>-0.59</td> </tr> </table>	MJSW (SRM)		-0.46	Cartilage volume	Medial tibia	-0.56	Medial femur	-0.74	Cartilage thickness	Medial tibia	-0.61	Medial femur	-0.59
MJSW (SRM)		-0.46																	
Cartilage volume	Medial tibia	-0.56																	
	Medial femur	-0.74																	
Cartilage thickness	Medial tibia	-0.61																	
	Medial femur	-0.59																	
Buck RJ 2010 [213]	70	Case-control	MRI	2 years	Proportion of patients presenting with cartilage thinning or thickening	20/70 patients cartilage thinning in the medial compartment 14/70 patients cartilage thickening in the medial compartment 13/70 patients cartilage thinning in the lateral compartment 6/70 patients cartilage thickening in the lateral compartment													
Eckstein F 2007 [214]	9	Cohort	MRI	2 years	Sensitivity to change of cartilage thickness and volume	Cartilage volume: MFTC SRM=0.76 LTFC SRM= 0.92 Cartilage thickness: MFTC SRM= 0.92													
Hudelmaier M 2010 [215]	23	Cohort	MRI	2 years	Sensitivity to change of cartilage thickness and volume	Cartilage volume: -0.44 Cartilage thickness: -0.58													
Iranpour-Bourjeni T 2011 [216]	24	Cohort	MRI	2 years	Sensitivity to change of cartilage volume	SRM 0.51													
Raynauld J 2004 [217]	32	Cohort	MRI	2 years	Sensitivity to change of cartilage thickness	SRM 0.84													
Raynauld J 2006 [144]	110	RCT	MRI	2 years	Sensitivity to change of cartilage volume	SRM -1.3													
Raynauld J 2008 [218]	107	RCT	MRI	2 years	Sensitivity to change of BMLs and cysts size	SRM cysts 0.13 SRM BMLs 0.13													
Raynauld J 2008 [219]	301	RCT	MRI	2 years	Sensitivity to change of cartilage thickness and volume	<table border="1"> <tr> <td></td> <td>Licofelone</td> <td>Naproxen</td> </tr> <tr> <td>CV</td> <td>2.11</td> <td>-2.8</td> </tr> <tr> <td>ThCtAb</td> <td>-2.3</td> <td>-2.9</td> </tr> </table>		Licofelone	Naproxen	CV	2.11	-2.8	ThCtAb	-2.3	-2.9				
	Licofelone	Naproxen																	
CV	2.11	-2.8																	
ThCtAb	-2.3	-2.9																	

MRI: magnetic resonance imaging; SRM: standardized response mean; KLG: Kellgren and Lawrence grade MFTC: medial tibiofemoral compartment; LTFC: lateral tibiofemoral compartment; CV: cartilage volume; ThCtAb: mean cartilage thickness over the bone area; BMLs: bone marrow lesions

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome						
Raynauld J 2009 [220]	355	RCT	MRI	2 years	Sensitivity to change of cartilage volume	CV	Licofelone		Naproxen		
							8.1			9.5	
Eckstein F 2012 [221]	441	Cohort	MRI	4 years	Yearly sensitivity to change of cartilage thickness from baseline to year 4		BL/Y1	Y1/Y2	BL/Y2	Y1/Y3	BL/Y4
						SRM	-0.42	-0.28	-0.56	-0.55	-0.78
Amin S 2005 [222]	323	Cohort	CR	30 months	Performance of CR in detecting MRI progression	Medial compartment: OR(95%CI)= 3.7 (2.2,6.3) Lateral compartment: OR (95% CI) 5.7 (2.3,14.0)					
Gandy S 2002 [223]	11	Cohort	MRI	30 months	Sensitivity to change of cartilage volume	SRM 0.26					
Hunter DJ 2006 [150]	217	Cohort	MRI	30 months	Correlation of cartilage loss and BMLs	Ipsilateral change in score: medial TF cartilage: 0.65 (0.003); lateral TF cartilage: 0.68 (0.01)					
Pelletier J 2008 [224]	27	Cohort	MRI	2 months	Sensitivity to change of synovitis and BMLs	Synovitis score: SRM 0.67 BMLs score: SRM 0.76					
Brandt KD 2006 [225]	30	Cohort	MRI	14 days	Sensitivity to change of effusion and synovitis volume			NSAIDs		ACET	
						Synovitis (SRM)		0.55		0.79	
						Effusion (SRM)		0.97		0.65	
Driban JB 2013 [226]	629	Cohort	MRI	6 months	Subchondral sclerosis progression depending on BMLs progression	subchondral sclerosis in BML progression OR 95% CI = 3.1 (1.10, 9.61) subchondral sclerosis in BML regression OR 95% CI= 1.74 (0.56,5.37)					
Eckstein F 2014 [227]	189	Case-control	MRI	16-20 months	Cartilage loss in patients undergoing TKA vs controls	OR = 1.36 (1.08,1.70)					

MRI: magnetic resonance imaging; CR: conventional radiography; SRM: standardized response mean; OR (95% CI) odds ratio and 95% confidence interval; CV: cartilage volume; ThCtAb: mean cartilage thickness over the bone area; BL: baseline; Y: year; TF: tibiofemoral

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Felson DT 2012 [228]	62	Cohort	MRI	6-12 weeks	Rate of progression in BMLs	<ul style="list-style-type: none"> At 6 weeks: 40.8% (20/49) knees showed BMLs changes beyond the MDD At 12 weeks: 38.8% (19/49) knees showed BMLs changes beyond the MDD 			
Hunter DJ 2006 [229]	217	Cohort	MRI	24 weeks	Sensitivity to change of cartilage morphology, BMLs, osteophytes, synovitis/effusion	SRM: Cartilage morphology -0.46; BME -0.18; osteophytes -0.33; Synovitis/effusion-0.36; Cartilage volume: medial tibia -0.05; lateral tibia -0.27; patella -0.20			
Hunter DJ 2010 [230]	29	Cohort	MRI	6 months	Change in cMF ThCtAb	no significant changes in cartilage thickness for any of the regions of interest on the TF joint over 3 or 6 months			
Hunter DJ 2011 [231]		SLR	MRI		Sensitivity to change of cartilage, synovium measures and BMLs		Medial TF	Lateral TF	PF
						Quantitative measures:	-0.86	-1.01	-0.63
						Semiquantitative measures	0.55	0.37	0.29
						Semiquantitative measures for the synovium	0.47		
		BMLs	0.43						
Kubota M 2010 [232]	180	Cohort	MRI	6 months	Change in BME score in radiographic progressors vs non progressors	Mean (sd)	Progressors	Non progressors	p
						KLG2	5.1 (5.2)	5.0 (2.6)	0.06
						KLG3	2.7 (1.8)	11.0 (6.3)	0.04
						KLG4	0.6 (1.2)	1.2 (2.6)	0.57

MRI: magnetic resonance imaging; SRM: standardized response mean; OR (95% CI) odds ratio and 95% confidence interval; BMLs: bone marrow lesions; NSAIDs: non-steroidal anti-inflammatory drugs; ACET: RCT on paracetamol; MDD: minimal detectable difference; BME: bone marrow edema; TKA: total knee arthroplasty

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
							Se	Sp	
Felson DT 2010 [154]	113	Cohort	MRI	24 months	WORMS and BLOKS cartilage loss vs radiographic JSN for monitoring cartilage loss				
						JSN medial compartment WORMS	0.86 (0.76,0.93)	0.42 (0.23,0.63)	
						JSN medial compartment BLOKS	0.87 (0.78,0.93)	0.42 (0.23,0.63)	
						JSN lateral compartment WORMS	0.96 (0.90,0.98)	0.50 (0.12,0.87)	
						JSN lateral compartment BLOKS	0.90 (0.83,0.95)	0.50 (0.12,0.87)	
Crema MD 2014 [233]	148	Cohort	MRI (dGEMRIC)	2 years	Correlation between dGEMRIC and changes in cartilage thickness	KL grade 0 and 1 and 3: no significant associations KL grade 2: significant association at the central medial femur between the decrease in dGEMRIC indices and the increase in cartilage thickness			
Kornaat P 2007 [234]	182	Cohort	MRI	2 years	Change in BMLs/change in symptoms, function and increase in cystic lesions		WOMAC pain	WOMAC function	
						unchanged BML	2 (-8, 12)	-2 (12, 8)	
						increasing BML	2 (-8, 11)	-4 (13, 6)	
						decreasing BML	1 (-11, 12)	-4 (15, 8)	
change in size of an adjacent cyst or BML	OR 95% CI=6.2 (3.2,12.3)								
Phan C 2006 [235]	34	Cohort	MRI	1 year-2 years	Cartilage and BMLs/WOMAC	r	Baseline (cartilage/BMLs)	12 months (cartilage/BMLs)	24 months (cartilage/BMLs)
						WOMAC pain	n.s./ n.s.	0.42/0.42	n.s./0.45
						WOMAC stiffness	n.s./ n.s.	0.54/0.51	n.s./ n.s.
						WOMAC function	n.s./ n.s.	0.51/0.45	0.63 /n.s.
MRI: magnetic resonance imaging; BMLs: bone marrow lesions; dGEMRCI: delayed gadolinium enhanced magnetic resonance imaging; WOMAC: Western Ontario McMaster Universities osteoarthritis index; r: correlation coefficient; KLG: Kellgran and Lawrence grade; PF: patellofemoral; n.s. not significant; HA: hyaluronic acid; cMF: central medial femur; ThCtAb: mean cartilage thickness over the bone area; BMLs: bone marrow lesions; BME: bone marrow edema; sd: standard deviation; JSN: joint space narrowing; Se: sensitivity; Sp: specificity; WORMS: whole organ magnetic resonance imaging score; BLOKS: Boston Leeds osteoarthritis knee score									

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Cubukcu D 2005 [236]	40	RCT	MRI	8 weeks	PF cartilage changes	no significant change in PF cartilage in the saline group ($p>0.05$), but statistically significant improvement in the HA group ($p<0.05$)
Stahl R 2007 [237]	8	Cohort	MRI	1 year	Sensitivity to change of cartilage volume and cartilage thickness	SRM cartilage volume: - 1.17 SRM cartilage thickness: -0.33
Stahl R 2011 [238]	30	Cohort	MRI	1 year	Prevalence of progression in cartilage lesions and BMLs	<ul style="list-style-type: none"> Using the WORMS 3.8% of the cartilage lesions in OA patients increased, while 96.2% of the lesions remained unchanged. BMEP decreased in 3.4% and increased in 4.9% of the lesions in the OA patients.
Wirth W 2009 [239]	156	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	MFTC.ThCtAB -0.24 (-0.30,-0.17)
Wirth W 2011 [240]	346	Cohort	MRI	2 years	Change in cartilage thickness	Change of ThCtAB TFJ: baseline-year 1 -0.32; year 1- year 2 -0.28; baseline year 2: -0.44
Wirth W 2011 [241]	290	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	MFTC (mcm) no JSN: SRM -0.12; JSN: -0.31; LFTC (mcm) JSN: -0.09 ; no JSN:-0.23
Wirth W 2010 [242]	80	Cohort	MRI	1 year	Sensitivity to change of cartilage thickness	SRM -0.29

MRI: magnetic resonance imaging; SRM: standardized response mean; WORMS: whole organ magnetic resonance imaging score; MFTC: medalfemorotibial compartment; ThCtAb: mean cartilage thickness over the bone area; TFJ: tibiofemoral joint; MFTC: medial femorotibial compartment; JSN: joint space narrowing; LFTC: lateral femorotibial compartment

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Zhang Y 2011 [243]	651	Cohort	MRI	3 months	Change in pain status according to change in BML and effusion/synovitis score	PAIN (OR 95%CI)	Absent/present	Present/absent
						BMLS 0	1	0.3 (0.1–0.9)
						BMLS 1	1.1 (0.6–2.3)	0.3 (0.1–1.0)
						BMLS 2	0.7 (0.3–1.6)	0.3 (0.1–1.1)
						BMLS 3	0.3 (0.3–1.5)	0.5 (0.2–1.6)
						BMLS 4	0.9 (0.3–2.3)	1.0 (0.3–3.4)
						BMLS 5-6	1.3 (0.5–3.4)	0.7 (0.3–1.8)
						BMLS 7-18	1.3 (0.5–3.6)	1
						Synovitis 0	1	0.4 (0.1–1.4)
						Synovitis 1	1.2 (0.7–2.3)	1.0 (0.3–2.8)
						Synovitis 2	2.1 (0.9–5.1)	0.6 (0.2–1.8)
						Synovitis 3-6	4.4 (1.4–14.0)	1
						Effusion 0	1	3.0 (1.1–8.2)
Effusion 1	1.4 (0.8–2.6)	0.8 (0.4–1.5)						
Effusion 2-3	2.3 (1.0–5.3)	1						
Cremer P 1994 [244]	12	RCT	MRI radioisotope scan	6 weeks	Sensitivity to change	No differences between baseline and follow up in Tc 99 scan MRI: non consistent differences between pre and post images		

MRI: magnetic resonance imaging; BMLs; bone marrow lesions; Tc: technetium; OR 95%CI: odds ratio and 95% confidence interval

US: all outcomes

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome					
Hall M 2014 [245]	243	Case-control	US	3 months	Sensitivity to change of effusion, synovial hypertrophy, popliteal cysts	Mean (sd)	baseline	3 months	p	
						Effusion	8.6 (4.6)	8.6 (4.9)	0.96	
						Synovial hypertrophy	7.1 (4.0)	5.7 (4.3)	0.003	
						Popliteal cysts	0 (0-14.3)	1.15 (0-13.4)	0.45	
						no correlation between change in pain and change in US measures for either group				
Acebes JC 2006 [246]	30	Cohort	US	4 weeks	Change in popliteal cyst sagittal area/ROM	r=0.380, p<0.05				
Bandinelli F 2012 [247]	40	Cohort	US	8 weeks	Change in popliteal cyst measures after steroid injection	Mean(sd)	baseline	2 weeks	4 weeks	8 weeks
						Longitudinal diameter	4.84 (1.42)	1.8 (1.22)*	1.69 (1.1)*	1.8 (1.22)*
						Transverse diameter	2.18 (0.8)	0.56 (0.51)*	0.75 (0.59)*	0.94 (0.68)*
						Thickness	1.53 (0.57)	0.41 (0.53)*	0.66 (0.66)*	0.76 (0.69)*
					*p<0.05					
Hall M 2014 [248]	25	RCT	US		Sensitivity to change of effusion and synovial hypertrophy	Synovial hypertrophy: SRM -0.40 Effusion: SRM -0.04				
Jan M 2006 [249]	44	Cohort	US	1 year	Change in synovial thickness	the synovial thickness in both treatment groups decreased to 81% to 84% of the initial thickness; after 30 treatment sessions, it was 67% to 72% of the initial thickness				
US: ultrasonography; sd: standard deviation; ROM: range of motion; r: correlation coefficient; SRM: standardized response mean										

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Kawaguchi K 2012 [250]	58	Cohort	US	1 year	Change in meniscal radial displacement	The MRD values in the standing position were significantly increased during the followup period, except for the KL grade 4 knees		
Song IH 2009 [251]	41	RCT	US (CEUS) MRI	16 weeks	Sensitivity to change of effusion, synovial hypertrophy, CEUS slope, MRI effusion	SRM	Superior recess	Lateral recess
						effusion	0.09	0.09
						synovitis	0.06	0.03
						CEUS slope	0.18	
MRI effusion	0.05	0.003						
Keen HI 2011 [252]		SLR	US					

US: ultrasonography; CEUS: contrast enhanced ultrasonography; MRD: meniscal radial displacement; KL: Kellgren and Lawrence; SRM: standardize response mean; MRI: magnetic resonance imaging; SLR: systematic literature review

TREATMENT OF OA - KNEE

Q9: What is the value of imaging-guided interventions in OA with respect to accuracy, therapeutic response, clinical and imaging outcome?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Bum Park Y 2012 [253]	99	RCT	US		Accuracy of HA injection vs blind injection OR (95%CI)= 4.68 (0.94,23.30)			
Im SH 2006 [254]	99	RCT	US		Accuracy of HA injection vs blind injection Accuracy: 95.5% (US) vs 77.2% (blind) p 0.01			
Jang SH 2013 [255]	126	RCT	US		US guided in plain injection vs US guided out of plane vs blind Triamcinolone exhacetonide Accuracy: US guided in plain 95.1%; US guided out of plain 97.7%; Blind 78% P<0.05 blind vs remaining procedures			
Sibbitt W 2011 [256]	92	RCT	US	2 weeks – 6 months	US guided vs blind trimacinolone			
						US	blind	p
					VAS pain	-42 (-78,-5) p=0.025	-0 (-18,+18) p=1.0	-
					Reinjection	24/46	34/46	0.03
VAS pain related to the injection	2.3 (2.4)	4.4 (2.9)	0.0003					
Cost per year (\$)	460 ± 207	173±81	0.0001					
Maricar N 2013 [257]		SLR	US		No quantitative synthesis, references reviewed			

US: ultrasonography; VAS: visual analogue scale

MAKING A DIAGNOSIS OF OSTEOARTHRITIS – FOOT

Q1: What is the added value of imaging to the diagnosis of OA?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome	
Matsos M 2009 [258]	62	Cross-sectional	US	Clinical diagnosis	Certainty pre-US vs certainty post-US	primary OA: 29 (46.8) vs 45 (73.0) p=0.002; inflammatory OA 29 (46.8) vs 54 (87.1) p<0.001

CR: conventional radiography, US: ultrasonography

Q2: What is the accuracy of different imaging modalities for detecting OA features (soft tissues and bone involvement)?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome		
Moon J 2010 [259]	83	Cross-sectional	CR	surgery	concordance between radiographic scores and cartilage damage	KLG 0.53 (0.38,0.66) Takakura 0.42 (0.24,0.57) van Dijk 0.42 (0.25,0.57)	
Tol J 2004 [260]	60	Surgery	CR	surgery	Accuracy of CR to detect osteophytes	Se	Sp
					Tibial OP	0.85	0.45
					Talar OP	0.73	0.68

CR: conventional radiography; Se: sensitivity; Sp: specificity; OP: osteophytes; KLG: Kellgran and Lawrence grade

PROGNOSIS OF OA- FOOT

Q5: What is the prognostic (prediction of outcome) value of imaging in OA?

Reference	No of subjects	Study design	Imaging modality	Outcome			
Bouaicha S 2010 [261]	198	Case-control	CR		Hallux rigidus	Hallux valgus	p
				MPE (mean 95% CI)	+5.2 mm(4.7, 5.7)	+2.8 mm (2.2, 3.4)	< 0.0001
				DFA	9° (8.10)	14° (13, 16)	p < 0.0001
Iagnocco A 2011 [262]	100	Cross-sectional	US	US abnormalities/joint tenderness and swelling			OR (95% CI)
						US abnormalities midfoot	16.75 (6.21,45.15)
						US abnormalities forefoot	4.28 (1.73,10.60)
Keen HI 2011 [263]	53	Case-control	US	GS, PD, Osteophytes and JSN at MTPs/ VAS pain and foot function index	p	VAS pain	Foot function index
					GS	0.55	0.40
					PD	0.43	0.03
					osteophytes	0.04	0.03
					JSN	0.47 (neg)	0.01
McDaniel G 2011[58]	138	Cross-sectional	CR	JSN, osteophytes and KLG in the tibiotalar and subtalar joint/		Gait velocity	Standing balance
					JSN tibiotalar	r=-0.06 p=0.45	r=-0.15 p>0.05
					JSN subtalar	r= -0.13 p=0.12	r= -0.29 p=0.0006
					Osteophytes tibiotalar	r= 0.03 p=0.69	r= -0.03 p=0.72
					Osteophytes subtalar	r= -0.03 p=0.74	r= -0.10 p=0.23
					KLG tibiotalar	r= -0.10 p=0.27	r= -0.19 p=0.02
					KLG subtalar	r= -0.15 p=0.08	-
Menz HB 2009 [264]	197	Cross-sectional	CR	CR/symptoms			OR (95% CI)
					OA at MTP1		1.2 (0.7,2.2)
					OA at 1st CMJ		1.1 (0.5,2.2)
					OA at 1st CJ		2.4 (1.3,4.3)
					OA at TNJ		0.8 (0.4,1.6)
					OA at 2nd CMj		2.2 (1.2,4.3)

CR: conventional radiography; US: ultrasonography; MPE: metatarsus primum elevatum; DFA: metatarsophalangeal dorsiflexion angle; OR (95% CI) ratio with 95% confidence interval; VAS: visual analogue scale; JSN: joint space narrowing; GS: grey scale; PD: power Doppler; KLG: Kellgren and Lawrence grade; MTP: metatarsophalangeal; CMJ: cuneiform-metatarsal joint; CJ: 1st navicular-cuneiform joint; TNJ: talonavicular joint; r: correlation coefficient

Q6: What is the prognostic (prediction of therapeutic response) value of imaging in OA?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Drakonaki E 2011 [265]	51	Cohort	US CR	12 months		US degenerative signs	US and CR degenerative signs	p
					Positive therapeutic response (i.a. methylprednisolone)	76.9%	75%	1
Han SH 2014 [266]	40	Cohort	CR	12 months	Response to intraarticular HA (VAS pain)	OR (95%CI)	Early stage 1-2 vs 3-4	
						3 months	11.00 (2.24,53.86)	
						6 months	10.55 (1.90,58.55)	
						12 months	5.00 (0.92,27.08)	
Sun S 2011[267]	46	Cohort	CR		KLG 2 and 3/AOS, AOFAS scores and	no significant difference in the AOS, AOFAS, or clinical balance test scores between KLG and 3 at any time point		

CR: conventional radiography; US: ultrasonography; OR (95% CI) ratio with 95% confidence interval; HA: hyaluronic acid; VAS: visual analogue scale; KLG: Kellgren and Lawrence scale, AOFAS: America Orthopedic ankle and foot society score; AOS: foot and ankle outcome score

MAKING A DIAGNOSIS OF OSTEOARTHRITIS – HAND

Q1: What is the added value of imaging to the diagnosis of OA?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome
Matsos M 2009 [258]	62	Cross-sectional	US	Clinical diagnosis	n (%) Certainty pre-US vs certainty post-US primary OA: 29 (46.8) vs 45 (73.0) p=0.002; inflammatory OA 29 (46.8) vs 54 (87.1) p<0.001

CR: conventional radiography; US: ultrasonography; Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; AUC: area under the curve

Q2: What is the accuracy of different imaging modalities for detecting OA features (soft tissues and bone involvement)?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome					
Cicuttini F 1998 [268]	6590 DIP	Cross-sectional	CR DIP	Physical examination	osteophytes	Se	Sp	LR+	LR-	
						0.49 (0.46,0.53)	0.88 (0.88,0.89)	4.51 (4.08,4.98)	0.565 (0.53,0.60)	
Grainger A 2007 [269]	120 joints	Cross-sectional	MRI DIP PIP	CR	erosions	Se	Sp	LR+	LR-	
						1 (0.66,1)	0.74 (0.65,0.82)	3.96 (2.88,5.46)	0	
Haims AH 2004 [270]	85	Cross-sectional	MRI wrist	Surgery	cartilage	Se	Sp			
						Distal radius	0.27	0.91		
						Scaphoid	0.31	0.90		
						Lunate	0.41	0.75		
						Triquetrum	0.18	0.93		
Haugen IK 2012 [271]	106	Cross-sectional	MRI DIP PIP	CR	Osteophytes Central erosions Marginal erosions JSN Cysts	Se	Sp			
						Osteophytes	1	0.22		
						Central erosions	0.78	0.72		
						Marginal erosions	0.83	0.94		
						JSN	0.95	0.63		
						Cysts	0.16	0.96		
Mutimer J 2008 [272]	20	Cross-sectional	MRI	Surgery	cartilage	Se	Sp	LR+	LR-	
						0.71 (0.58,0.82)	0.77 (0.61,0.89)	3.19 (1.75,5.78)	0.37 (0.24,0.57)	
Keen HI 2008 [273]	37	Cross-sectional	US	CR	JSN osteophytes	Se	Sp	LR+	LR-	
						JSN	0.82 (0.77,0.86)	0.72 (0.69,0.75)	2.92 (2.62,3.35)	0.25 (0.19,0.32)
						osteophytes	0.83 (0.78,0.87)	0.75 (0.72,0.78)	3.42 (2.99,3.90)	0.22 (0.17,0.28)
Koutroumpas AC 2010 [274]	18	Cross-sectional	US DIP PIP	Physical examination	synovitis	PD as reference standard for inflammation: Se of clinical assessment = 0.15, Sp = 0.96				

CR: conventional radiography; US: ultrasonography; Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; JSN: joint space narrowing

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome					
Mathiessen A 2013 [275]	127	Cross-sectional	US DIP PIP MCP CMC	Physical examination CR MRI	osteophytes		Se	Sp	LR+	LR-
						US vs MRI	0.95 (0.93,0.96)	0.86 (0.81,0.90)	7.08 (5.04,9.93)	0.05 (0.04,0.08)
						US vs CR	0.04 (0.029,0.05)	0.35 (0.33,0.36)	0.06 (0.05,0.08)	2.74 (2.59,2.90)
						US vs PE	0.89 (0.87,0.91)	0.68 (0.67,0.70)	2.89 (2.71,3.08)	0.15 (0.13,0.18)
Vlychou M 2013 [276]	20	Cross-sectional	US DIP PIP MCP CMC	MRI	Osteophytes Erosions Cysts Synovitis Tenosynovitis Effusion		Se	Sp		
						osteophytes	0.9 (0.84,0.94)		0.95 (0.85,0.99)	
						erosions	0.88 (0.81,0.93)		0.96 (0.89,0.98)	
						cysts	0.87 (0.75,0.94)		0.97 (0.92,0.99)	
						synovitis	0.84 (0.75,0.90)		0.96 (0.91,0.99)	
						tenosynovitis	0.8 (0.57,0.94)		0.99 (0.96,1)	
effusion	0.92 (0.73,0.99)		0.98 (0.95,0.99)							
Wittoek R 2010 [277]	38	Cross-sectional	US DIP PIP IP1	CR	Erosions	Se	Sp	LR+	LR-	
						0.94(0.86,0.98)	0.90 (0.87,0.93)	10.20 (7.68,13.55)	0.06 (0.02,0.16)	
Iagnocco A 2005 [278]	110	Cross-sectional	US DIP PIP	CR	Erosions	Se	Sp	LR+	LR-	
						0.72 (0.49,0.89)	1 (0.95,1)	-	0.27 (0.14,0.54)	
Vlychou M 2009 [279]	22	Cross-sectional	US DIP PIP MCP CMC	CR	Erosions osteophytes	<ul style="list-style-type: none"> Erosions were detected in 231/660 (35%) small joints by US imaging, and in 115/660 (17.4%) small joints with conventional radiographs [P<0.05]. Osteophytes were detected in 360/660 (54.5%) small joints by US imaging and in 310/660 (47%) small joints with conventional radiographs [P<0.05] 				
Wittoek R 2011 [280]	14	Cross-sectional	US DIP PIP IP1	MRI	Erosions	Se	Sp	LR+	LR-	
						0.65 (0.50,0.79)	0.90 (0.80,0.96)	7.15 (3.24,15.74)	0.38 (0.25,0.56)	
Iagnocco A 2000 [281]	57	Cross-sectional	US	Synovial fluid aspiration	Effusion	Successful synovial fluid aspiration in patients with US effusion: 18/18, Se=1 Visualization of effusion in healthy controls in which saline solution was injected: 3/3, Se=1				

CR: conventional radiography; US: ultrasonography; MRI: magnetic resonance imaging; Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; MCP: metacarpophalangeal joints; CMC: carpometacarpal joints; IP1: first interphalangeal joint; PE: physical examination

PROGNOSIS OF OA - HAND

Q5: What is the prognostic (prediction of outcome) value of imaging in OA?

Cross-sectional studies

Reference	No of subjects	Study design	Imaging modality	Outcome				
Jones G 2001 [282]	522	Cross-sectional	CR CMC DIP	Altman score/AUSCAN	Beta coefficient, p	CMC score	DIP score	
					AUSCAN pain	+0.14 (0.024)	+0.17 (0.003)	
					AUSCAN function	+0.19 (<0.001)	+0.15 (0.012)	
					Grip strenght	-0.09 (0.010)	-0.12 (0.012)	
Bijsterbosch J 2010 [283]	236	Cross-sectional	CR	Erosions/pain function and QoL	Mean, sd	Erosive OA	Non erosive OA	p
					AUSCAN pain (0–20)	9.0±4.8	7.0±4.8	0.016
					MHQ pain (0–100)	47.1±18.1	37.9±22.8	0.016
					Number of painful joints (0–30)	11.3±6.8	7.9±7.8	0.009
					Pain intensity (0–90)	8.7±7.2	6.6±7.0	0.082
					AUSCAN function (0–36)	17.3±8.7	13.2±8.7	0.006
					MHQ overall function (0–100)	53.5±14.8	61.2±15.6	0.004
					MHQ ADL (0–100)	73.2±19.4	79.3±17.8	0.049
					MHQ work performance (0–100)	65.2±24.5	71.1±25.9	0.182
					Grip strength (kg)	19.7±8.4	21.7±10.7	0.241
					Pinch grip (kg)	3.2±1.8	3.2±1.5	0.872
HAMIS (0–27)	5.7±4.0	3.7±2.6	<0.001					
Fingertip to palm distance (mm)	54.0±52.1	15.1±27.3	<0.001					

CR: conventional radiography; CMC: carpometacarpal joint; DIP: distal interphalangeal joint; AUSCAN: Australian Canadian OA hand index; MHQ: Michigan hand questionnaire; ADL: activities of daily living; HAMIS: hand mobility in scleroderma; QoL: quality of life

Reference	No of subjects	Study design	Imaging modality	Outcome								
Kortekaas M 2011 [284]	55	Cross-sectional	CR US CMC MCP DIP PIP IP1	US and CR osteophytes and JSN/pain							OR 95% CI	
					US osteophytes	Grade 1 vs 0						1.6 (1.2, 2.3)
						Grade 2 vs 0						2.7 (1.8, 4.1)
						Grade 3 vs 0						4.8 (3.1, 7.5)
					CR osteophytes	Grade 1 vs 0						2.0 (1.5, 2.7)
						Grade 2 vs 0						3.2 (2.1, 4.8)
						Grade 3 vs 0						4.1 (2.4, 7.1)
					CR JSN	Grade 1 vs 0						1.8 (1.3, 2.4)
						Grade 2 vs 0						4.3 (2.6, 7.2)
Grade 3 vs 0						4.2 (2.0, 9.0)						
Kwok YK 2013[285]	80	Cross-sectional	CR DIP PIP MCP CMC ST IP1	Erosions/pain and function							Erosive vs non erosive (mean difference 95% CI)	
					AUSCAN pain						1.3 (0.3, 2.3)	
					AIMS2 pain subscale						0.8 (0.3, 1.4)	
					AUSCAN function						2.3 (0.4, 4.2)	
					AIMS2 hand finger subscale						0.8 (0.2, 1.3)	
Grip strenght						-3.0 (-7.1,3.4)						
Marshall M 2009 [286]	592	Cross-sectional	CR DIP PIP MCP CMC ST IP1	OA location/pain and function	Mean	Non OA	Finger OA	Thumb OA	Finger+ thumb OA	p		
					AUSCAN pain	8.3 (6.7-9.8)	8.2 (6.3-10.1)	8.6 (6.9-10.3)	10.5 (9.6-11.4)	0.0018		
					AUSCAN function	5.4 (4.6-6.2)	5.7 (4.7-6.8)	5.8 (4.9-6.7)	6.5 (6.1-7.0)	0.07		
Wittoek R 2012 [287]	349	Cross-sectional	CR DIP PIP CMC	number of remodeled joints and CMC involvement/pain and function	Beta coefficient, p		No of remodeled joints		CMC involvement			
					VAS pain		0.99 p<0.001		-3.31 p>0.05			
					AUSCAN pain		0.25 p<0.001		-2.72 p>0.05			
					AUSCAN function		2.01 p<0.01		-2.72 p>0.05			
					FIHOA		3.51 p<0.01		1.59 p>0.05			

CR: conventional radiography; US: ultrasonography; CMC: carpometacarpal joint; DIP: distal interphalangeal joint; MCP: metacarpophalangeal joints; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; IP1: 1st interphalangeal joint; JSN: joint space narrowing; OR (95%CI): odds ratio with 95% confidence intervals; AUSCAN: Australian Canadian OA hand index; AIMS: arthritis impact measurement scale; VAS: visual analogue scale; FIHOA: functional index for hand OA.

Reference	No of subjects	Study design	Imaging modality	Outcome			
Keen HI 2008 [288]	67	Case-control	US CMC MCP DIP PIP IP1	Osteophytes, PD, effusion and synovial thickening/pain and tenderness	Painful and tender joints were significantly more likely to have osteophytes, JSN, synovitis or PD		
Kortekaas M 2010 [289]	55	Cross-sectional	US CMC MCP DIP PIP IP1	GS and PD/ tenderness		OR (95%CI)	
					GS	1vs0	2.1 (1.5, 2.8)
						2vs0	4.7 (2.8, 7.8)
						3vs0	4.0 (1.9, 8.2)
					Effusion	1vs0	2.0 (1.5, 2.6)
						2vs0	3.8 (2.3, 6.1)
						3vs0	3.7 (1.8, 7.6)
					Synovial thickeninkg	1vs0	1.3 (0.7, 2.4)
						2-3 vs 0	2.6 (1.1, 6.3)
					PD	1vs0	1.4 (1.0, 2.1)
2-3 vs 0	2.0 (0.8, 4.9)						
Jonsson H 1999 [290]	297	Cross-sectional	Radioisotope scan	Uptake/symptoms	“Hand symptoms were related to distal interphalangeal (DIP) and CMC1 uptake, thumb symptoms with first metacarpophalangeal joint (MCP1) CMC1 uptake, and knee symptoms with the subchondral knee uptake pattern.”		
Dahaghin s 2006 [291]		SLR					
Kwok WY 2013 [292]		SLR					
US: ultrasonography; CMC: carpometacarpal joint; DIP: distal interphalangeal joint; MCP: metacarpophalangeal joints; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; IP1: 1 st interphalangeal joint; PD: power Doppler; GS: grey scale; OR (95% CI): odds ratio with 95% confidence interval; SLR: systematic literature review							

Reference	No of subjects	Study design	Imaging modality	Outcome				
Haugen I 2012 [293]	85	Cross-sectional	CR MRI DIP PIP	CR and MRI/pain and function		Tenderness (OR 95% CI)	FIHOA B (beta coefficients)	Grip strength (beta coefficients)
					Synovitis	2.4, (1.5,3.6)	0.45 (0.01,0.89)	-
					BMLs	1.6 (1.0, 2.5)	-	-
					Osteophytes	1.4 (0.9, 2.1)	0.23 (0.07, 0.39)	-0.39 (-0.58, -0.21)
					Bone attrition	2.5 (1.5, 4.1)	0.58 (0.18,0.97)	-0.87 (-1.34, -0.39)
					Flexor tenosynovitis	1.5 (1.0,2.3)	-	
					Erosions	-	-	-0.43 (-0.69,-0.16)
JSN	-	0.28 (0.01, 0.54)	-0.53 (-0.84,-0.23)					
Wenham C 2012 [294]	65	RCT	MRI	Synovitis and effusion/pain and tenderness	Tenderness: OR (95%CI) 4.7 (2.8, 7.9) Pain: OR (95%CI) 3.7 (95% CI 2.2, 6.2)			
CR: conventional radiography; MRI: magnetic resonance imaging; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; OR (95% CI): odds ratio with 95% confidence interval; FIHOA: functional index for hand OA; BMLs: bone marrow lesions								

Longitudinal studies

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Haugen IK 2013 [295]	190	Cohort	CR DIP PIP MCP CMC	7 years	KLG/tenderness	Joint tenderness increases with the severity of KLG. In multivariate model significant association with osteophytes, malalignment and erosions			
Ding H 2011 [296]	436	Cohort	CR DIP PIP MCP IP1	5 years	KLG≥2/incident pain	Fingers 1-3 right	RR= 1.76 (1.16,2.69)		
						Fingers 1-3 left	RR= 3.0 (1.97,4.58)		
						Fingers 4-5 right	RR= 2.30 (1.38,3.82)		
						Fingers 4-5 left	RR= 1.93 (1.06,3.53)		
Bijsterbosch J 2011 [297]	236	Cohort	CR	6 years	Osteophytes and JSN/erosive progression	Osteophyte grade 2-3 OR (95% CI) 0.7 (0.3,2.0) JSN grade 2-3 8.9 OR (95% CI) (4.8,16.4)			
Dahaghin S 2005 [298]	1235	Cohort	CR	6.6 years	Hand OA/incident knee or hip OA	OR= 2.2 (1.5,3.3) for incident hip/knee OA in case of hand OA			
Filkova M 2009 [299]	88	Cohort	CR	2 years	Erosions/Kallman score progression		Erosive OA	Non erosive OA	p
						Kallman score progression (mean, sd)	2.41(3.06)	0.57 (1.07)	0.002
Balblanc J 1995 [300]	15	Cohort	Radioisotope scan	4 years	Abnormal scan/incident OA in previously normal joints	OR (95% CI)= 5.95 (1.51,23.44)			
Hutton CW 1986 [301]	14	Cohort	Radioisotope scan	5 years	Abnormal scan/structural progression	OR 95% CI= 7.22 (4.28, 12.17)			
McCarthy M 1994 [302]	67	Cohort	Radioisotope scan	5-6 years	Abnormal scan/structural progression	OR (95% CI)= 14.53 (9.25,22.82)			

CR: conventional radiography; CMC: carpometacarpal joint; MCP: metacarpophalangeal joints; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; IP1: 1st interphalangeal joint; KLG: Kellgren and Lawrence grade; OR (95% CI): odds ratio with 95% confidence interval; RR: risk ratio; sd: standard deviation; JSN: joint space narrowing

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Kortekaas M 2014 [303]	25	Cohort	US	3 months	Synovial thickening, effusion and PD/incident pain	synovial thickening OR (95%CI) 2.0 (1.1, 3.9) effusion OR (95%CI) 4.8 (2.2, 10.5) PD OR (95%CI) 3.8 (2.7, 5.4)
Zhang W 2008 [304]		Case-control	CR		Hand finger pattern/incident hip or knee OA	Development of OA in case of type 3 finger pattern (OR 95% CI) Knee OA: OA 1.94 (1.54–2.44) Hip OA: 1.37 (1.13–1.67) IP nodes: 0.99 (0.83–1.18) IP nodes+ hip or knee OA 1.69 (1.33–2.16)

US: ultrasonography; CR: conventional radiography; DIP: distal interphalangeal joints; PIP: proximal interphalangeal joints; PD: power Doppler; JSN: joint space narrowing; OR (95% CI): odds ratio with 95% confidence interval

Q6: What is the prognostic (prediction of therapeutic response) value of imaging in OA?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Mallinson P 2013 [305]	31	Cohort	US CR	6 weeks	CR and US/ response to intraarticular triamcinolone	No significant association between treatment response and grades for osteophytes (p =0.148), joint-space narrowing (p = 0.850), and capsule thickness (grade or size; p =0.918, p =0.235)
Wenham C 2012 [294]	65	RCT	MRI	12 weeks	MRI/response to prednisolone 5 mg	The baseline number of joints with definite synovitis or effusion did not correlate with OARSI response at 4 weeks.

US: ultrasonography; CR: conventional radiography; MRI: magnetic resonance imaging; OARSI: osteoarthritis research society international

Q7: When, how often and under what clinical circumstance should imaging be used in the follow-up of osteoarthritic joints?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Bijsterbosch J 2011 [297]	172	Cohort	CR	2-6 years	Proportion of joints progressing according to KLG, VV grade or OARSI atlas	(%)	KLG	VV	OARSI
						2 years	18.9-56.2	13.3-52.2	6.7-37.8
						6 years	50.6-79.8	26.7-65.6	33.3-74.4
Botha Scheepers S 2009 [306]	90	Cohort	CR	2 years	Sensitivity to change of JSN and osteophytes	JSN: SRM= 0.35 Osteophytes: SRM= 0.34			
Botha-Scheepers S 2005 [307]	20	Cohort	CR	2 years	Sensitivity to change of JSN and osteophytes	JSN: SRM= 0.38 Osteophytes: SRM= 0.41			
Haugen IK 2013 [295]	190	Cohort	CR	7 years	Radiographic progression/incident tenderness	joints with progression had higher odds of tenderness, joints with incident KL 3 or 4 had higher odds of tenderness			
Maheu EC 2007 [308]	105	RCT	CR	12 months	Sensitivity to change of VV score, OA global score, Kallman score and KLG	KLG: SRM= 0.17 Kallman: SRM= 0.26 Verbrueggen: SRM= 0.18 Global score: SRM= 0.17			
Macfarlane DG 1991 [309]	32	Cohort	CR Radioisotope scan	1 year	Sensitivity to change of osteophytes detected by CR and radioisotope scan	Radioisotope scan SRM=0.21 CR SRM=0.20			

CR: conventional radiography; KLG: Kellgren and Lawrence grade; VV: Verbruggen-Veys score; OARSI: osteoarthritis research society international; JSN: joint space narrowing; SRM: standardized response mean; DIP: distal interphalangeal joints; MCP: metacarpophalangeal joints

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
						Mean	baseline	12 months	p
Jans L 2013 [310]	9	Cohort	MRI (DCE)	1 year	Change of DCE parameters during 1 year	Synovium maximum upslope	19.8	3.1	0.002
						Synovium enhancement curve	2.31	0.76	0.01
						Absolute enhancement	5.9	3.1	0.002
Keen HI 2011 [311]	36	Cohort	US	12 weeks	Correlation between GS and PD synovitis and decrease in pain after i.m. methylprednisolone	No significant association between changes in GS and PD at 4 and 12 weeks and response			
Klauser AS 2012 [312]	33	Cohort	US	4 weeks	Correlation between joint thickening and PD and VAS pain after intraarticular HA	decrease VAS pain statistically significantly correlated with the decrease of joint thickening (p < 0.001; r = 0.7) and PDUS score (p < 0.001; r = 0.8)			
Kortekaas MC 2014 [303]	25	Cohort	US	3 months	Change in scores for synovial thickening, effusion and PD		baseline	3 months	
						synovial thickening	5(0-23)	4 (0-19)	
						effusion	6(0-25)	8 (0-27)	
						PD	3(0-11)	2 (0-14)	
Saltzherr 2014 [313]		SLR	US						

MRI: magnetic resonance imaging; US: ultrasonography; OR (95% CI): odds ratio with 95% confidence interval; BMLs: bone marrow lesions; DCE: dynamic contrast enhanced; GS: grey scale; PD: power Doppler; i.m.: intramuscular; VAS: visual analogue scale; HA: hyaluronic acid; PDUS: power Doppler ultrasound; SLR: systematic literature review

Q9: What is the value of imaging-guided interventions in OA with respect to accuracy, therapeutic response, clinical and imaging outcome?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Karalezli N 2007 [314]	16	Cohort	CR	1 month	Fluoroscopy-guided vs blind injections of HA in TM	VAS pain related to the procedure: Fluoroscopic guide: 4.1 (range 3–6) Anatomic guide 5.6 (range 3–7) p<0.005 Safety (guided vs blind) 1.00 (0.017, 56.46)
US: ultrasonography; CR: conventional radiography; HA: hyaluronic acid; TM: trapeziometacarpal; VAS: visual analogue scale; sd: standard deviation						

MAKING A DIAGNOSIS OF OSTEOARTHRITIS – HIP

Q2: What is the accuracy of different imaging modalities for detecting OA features (soft tissues and bone involvement)?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome						
Leydet-Quilici H 2010 [315]	23	Cross-sectional	MRI	surgery	BMLs/histology	Se	Sp				
						0.80	0.95				
Taljanovic M 2008 [316]	19	Cross-sectional	MRI	Surgery	BMLs/histology	number of microfractures and focal signal/intensity ratio ($r=-0.48$, $P<0.05$). No statistically significant correlation between the calculated MRI values and other histopathologic findings, including the amount of BME					
Xu L 2013 [317]	44	Cross-sectional	CR	MRI	CR vs MRI		Se	Sp	LR+	LR-	AUC
						JSN	0.64	0.88	0.90	0.58	0.75
						osteophytes	0.84	0.71	0.94	0.45	0.77
						subchondral cysts	0.44	0.89	0.58	0.94	0.66
				bone attrition	0.78	0.86	0.58	0.94	0.81		

CR: conventional radiography; MRI: magnetic resonance imaging; BMLs: bone marrow lesions; Se: sensitivity; Sp: specificity; LR+: positive likelihood ratio; LR-: negative likelihood ratio; AUC: area under the curve; r: correlation coefficient; JSN: joint space narrowing;

MANAGEMENT OF OA - HIP

Q4: In people with OA do imaging have an impact on management?

Reference	No of subjects	Study design	Imaging modality	Comparator	Outcome	
Dolin SJ 2003 [318]	72	Cross-sectional	CR	surgery	Priority of THR assignment based on radiographic severity graded 0-15	Relative Risk (95%CI) for priority assignment: CR grading>9/15, RR (95% CI) 1.98 (1.23, 3.19)
CR: conventional radiography; THR: total hip replacement; RR (95%CI): risk ratio with 95% confidence interval						

PROGNOSIS OF OA - HIP

Q5: What is the prognostic (prediction of outcome) value of imaging in OA?

Cross-sectional studies

Reference	No of subjects	Study design	Imaging modality	Outcome			
Taljanovic M 2008 [316]	19	Cross-sectional	MRI	BMLs/symptoms	the total signal/intensity ratio of BML was significantly correlated with symptoms: pain : r -0.46 p<0.05		
Iagnocco A 2012 [319]	75	Cross-sectional	US	effusion synovitis osteophytes/pain		Previous hip pain	Current hip pain
					Joint effusion	93.3% vs 73.7% p<0.001	93.3% 57.9% p<0.0001
					Synovial hypertrophy	90% vs 82.7% p>0.05	90% vs 72.4% p>0.05
					osteophytes	82.5% vs 100% p>0.05	82.5% vs 100% p>0.05
Recnik G 2009 [320]	54	Cross-sectional	CR	Measure of centre-edge angle/age at THR	β coefficient for linear regression was 0.271 (r ² = 0.073; P = 0.024) for the correlation between centre-edge angle and age at THR, with lower angles associated to younger age		
Lievense A 2002 [321]		SLR					
Wright AA 2009 [322]		SLR					
Holla JF 2011 [57]	497	Cross-sectional	CR	CR/function	Beta coefficient (95% CI)	Internal hip rotation	External hip rotation
					severe JSN superior	-12.20 (-16.00,-8.40) p<0.0001	-5.39 (-9.00, -1.78) p=0.004
					severe JSN medial	-3.47 (-7.08, 0.14) p=0.14	-3.49 (-6.60, -0.37) p=0.03
					moderate/severe osteophytes superior	-6.48 (-9.13,-3.48) p<0.0001	-1.22 (-3.67, 1.23) p=0.33
					moderate/severe osteophytes inferior	-16.54 (-21.93,-11.15) p<0.0001	-4.95 (-9.83, -0.07) p=0.05
					subchondral sclerosis	-6.71 (-11.04, -2.38) p=0.002	-3.36 (-7.27, 0.56) p=0.09
					flattening of the femoral head	-0.69 (-3.60, 2.23) p=0.64	-4.52 (-7.07, -1.96) p=0.001
femoral buttressing	-1.46 (-3.75, 0.82) p=0.21	-4.41 (-6.44, -2.39) p<0.0001					
CR: conventional radiography; MRI: magnetic resonance imaging; US: ultrasonography; BMLs: bone marrow lesions; r: correlation coefficient; JSN: joint space narrowing							

Longitudinal studies

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Agricola R 2013 [323]	720	Cohort	CR	5 years	amount of acetabular coverage/ VAS pain, function No significant prediction of 5-years incident pain Lateral pincer deformity resulted in decreased internal rotation (29.87° vs 30.82°, P = 0.035)				
Agricola R 2013 [324]	723	Cohort	CR	5 years	active shape modeling of proximal femur and pelvis/pain, ACR criteria <ul style="list-style-type: none"> • mode 9 associated with VAS pain • no modes predicted the fulfillment of the ACR criteria • mode 7 predicted internal rotation <15° 				
Arden NK 2009 [325]	5839	Cohort	CR	5 years	CR/pain and THR	OR (95%CI)	pain	Change in lower limb disability	THR
						osteophyte grade ≥2	4.1 (2.4,6.9)	1.0 (0.8–1.4)	9.9 (6.0,16.3)
						JSN grade ≤2 lateral OR JSN grade ≤3 medial	4.8 (2.3,8.0)	2.2 (1.7–2.8)	8.2 (4.6,14.6)
						MJS ≤1.5	5.2 (3.0,9.1)	1.9 (1.5–2.5)	10.3 (6.0,17.8)
						MJS ≤2.0	3.2 (1.9,5.5)	1.7 (1.3,2.1)	6.4 (3.8,11.0)
						MJS ≤2.5	2.4 (1.5,3.9)	1.4 (1.1,1.7)	3.9 (2.3,6.7)
						KLK ≥2	5.7 (3.2,9.8)	1.1 (0.8,1.5)	13.8 (8.3,23.0)
Gossec L 2009 [326]	1404	Cohort, RCT	CR	3 years	KLG, OARSI JSN, JSW/pain, function and THR	OR (95% CI)	Symptoms (cross-sectional)	Structural progression	THR
						KLK 2 vs 1	0.87 (0.33,2.28)	1.31 (0.50,3.43)	
						KLK 3 vs 1	0.88 (0.33,2.39)	1.73 (0.64,4.68)	KLK 3 vs 2 2.62(1.67,4.13)
						KLK 4 vs 1	0.25 (0.01,4.66)	1.45 (0.10,21.07)	
						OARSI 2 vs 1	1.09 (0.77,1.54)	1.14 (0.81,1.62)	
						OARSI 3 vs 1	0.85 (0.45,1.61)	0.96 (0.51,1.82)	4.47 (1.98,10.05)
						JSW 1 vs 0	0.92 (0.55,1.53)	0.77 (0.46,1.29)	
						JSW 2 vs 0	1.14 (0.72,1.82)	1.47 (0.92,2.36)	
JSW 3 vs 0	1.00 (0.60,1.68)	1.21 (0.72,2.03)	4.02 (1.88,8.56)						

CR: conventional radiography; VAS: visual analogue scale; ACR: American college of Rheumatology; THR: total hip replacement; OR 95% CI: odds ratio with 95% confidence interval; JSN: joint space narrowing; mJS: medial joint space; KLG: Kellgren and Lawrence grade

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome			
Gossec L 2005 [327]	505	Cohort	CR	2 years	KLG and pattern of migration of the femoral head/THR OR (95% CI) of 2 year THR: KLG 3 vs 2: 3.3 (1.7,6.4); KLG 4 vs 2: 5.3 (2.6,10.8)			
Lievse A 2007 [328]	224	Cohort	CR	6 years	KLG, JSW<2.5 mm, osteophytes/THR	3 years THR (OR, p)	6 years THR (OR, p)	
						KLG≥2	13.0 p<0.01	17.8 p<0.01
						osteophytes	3.3 p=0.01	3.1 p<0.01
						JSW<2.5	6.6 p<0.01	7.1 p<0.01
Maillefert JF 2002 [329]	422	Cohort	CR	5 years	mJSW at 1 and 2 years follow up/THR Thresholds of -0.2 mm and -15%, i.e. a decrease in JSW of at least 0.2 mm (15%), were the most relevant, with sensitivity of 75 and 68% (20.2 mm) and 74 and 78% (215%) respectively.			
Lane NE 2004 [330]	745	Cohort	CR	8 years	CR/structural progression THR	OR (95% CI)	Radiographic progression	THR
						osteophytes	2.1 (1.0, 4.3)	3.8 (2.1, 7.1)
						cysts	0.6 (0.3, 1.0)	3.5 (1.8, 6.8)
						sclerosis	1.2 (0.9, 1.8)	5.1 (1.8, 14.6)
						JSN concentric	0.5 (0.2, 1.1)	9.2 (3.8, 22.5)
						JSN superolateral	0.5 (0.2, 1.1)	3.0 (1.4, 6.3)
JSN superomedial	0.6 (0.3, 0.9)	5.1 (2.2, 11.9)						
Ledingham J 1993 [331]	252	Cohort	CR	27 months	KLG and pattern of migration of the femoral head/THR, rapid, slow or no progression Rapid progression: superior migration 9.0 (1.24, 183.52); NO progression: %medial/axial migration 3.71(1.09,13.83); intermediate migration 11.41 (2.47,72.79) THR: OR 95% CI any joint progression 2.19 (1.18, 4.08); rapid joint progression 5.83 (1.90,19.11)			
Conrozier T 1998 [332]	367	Cohort	CR	1 year	JSW/JSW progression the mean JSW at 12 months was correlated with baseline JSW (r=0.66, p=0.01)			
Reijman M 2005 [333]	1904	Cohort	CR	1 year	KLG and baseline JSW/radiographic progression	OR (95%CI)		
						KLG≥2	5.8 (4.0, 8.4)	
						Baseline JSW≤2.5 mm	1.9 (1.2,2.9)	

CR: conventional radiography; KLG: Kellgren and Lawrence grade; OARSI: osteoarthritis research society international; JSN: joint space narrowing; THR: total hip replacement; OR 95% CI: odds ratio with 95% confidence interval; JSW: joint space width; mJSW: minimal joint space width

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome		
Barr RJ 2012 [334]	195	Cohort	CR	5 years	active shape modeling/symptoms and function	<ul style="list-style-type: none"> correlation between mode 2 and baseline pain duration: $r=-0.26$ ($p=0.005$) mode 4 vs baseline WOMAC function $r=-0.19$ ($p=0.034$) mode 2 OR=0.54 (0.29, 0.99) for 5 years THR 	
Agricola R 2013 [335]	865	Cohort	CR	5 years	α angle/incident OA	OR (95% CI)=2.42 (1.15,5.06) for incident OA. OR (95% CI)=9.66 (4.78,19.78) for end stage OA	
Dougados M 1996 [336]	508	Cohort	CR	1 year	pattern of migration of the femoral head, KLG and JSW/change in JSW	OR (95%CI)	
						Superolateral migration	4.25 (2.26,8.01)
						Baseline JSW\leq2 mm	2.11 (1.30,3.44)
Ahedi H 2014 [337]	198	Cohort	MRI		BMLs/cross sectional and incident pain	Cross-sectional hip pain: OR (95%CI)= 4.43 (1.46, 13.2) Incident hip pain: BMLs resolving vs absent MD (sd) -0.95 (-2.85 to +1.04) Incident vs absent +2.08 (+0.22 to +3.94). knee pain BMLs resolving vs absent MD (sd) -1.81 (-3.82 to +0.21); Incident vs absent -0.75 (-2.34 to +0.82)	
Zhang W 2008 [304]		Case-control	CR		Hand finger pattern/incident hip or knee OA	Development of OA in case of type 3 finger pattern: Knee OA: OA 1.94 (1.54-2.44) Hip OA: 1.37 (1.13-1.67) IP nodes: 0.99 (0.83-1.18) IP nodes+ hip or knee OA 1.69 (1.33-2.16)	
CR: conventional radiography; MRI: magnetic resonance imaging; KLG: Kellgren and Lawrence grade; THR: total hip replacement; OR (95% CI): odds ratio with 95% confidence interval; JSW: joint space width; BMLs: bone marrow lesions; MD: mean difference; IP: interphalangeal							

Q6: What is the prognostic (prediction of therapeutic response) value of imaging in OA?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Atchia I 2011 [338]	77	RCT	US	16 weeks	Synovitis/response to methylprednisolone	OR 95% CI= 16.7 (1.4,204)			
Hirsch G 2013[170]		SLR							
Keurentjes J 2013 [164]	723	Cohort	CR	5 years	KLG/QoL after THA or TKA	SF36 subscales Mean difference (95%CI) KLG0-2 vs 3-4			
						Physical functioning	8.93 (2.14,15.7) p=0.01		
						Role physical	6.39 (25.89,18.7) p=0.31		
						Bodily pain	0.88 (26.08,7.84) p=0.80		
						General Health	20.66 (25.66, 4.34) p=0.79		
						Vitality	23.53 (29.03,1.97) p=0.21		
						Social functioning			
						Role emotional	3.11 (28.22,14.4) p=0.59		
						Mental health	21.80 (26.13,2.50) p=0.41		
						PCS	1.94 (20.57-4.44) p=0.13		
MCS	22.03 (24.46,0.39) p=0.10								
NRS satisfaction	0.3 (20.2-0.9) p=0.19								
Lequesne M 2002 [339]	163	RCT	CR	2 years	JSW/structural progression	In patients with initial smaller JSW treated with avocado soybean, the reduction of JSW was half than in the placebo group; no differences among treatment and placebo in patients with more JSW			
Renneson- Rey B 2008 [340]	55	Cohort	US CR	6 months	Effusion and KLG/OARSI response to HA	OARSI response	KLG 1-2	KLG 3-4	p
						1 month	50%	22%	0.06
						3 months	58%		0.15
						6 months	10%	18%	
effusion	No significant difference								
CR: conventional radiography; US: ultrasonography; OR (95%CI): odds ratio with 95% confidence interval; KLG: Kellgren and Lawrence grade; QoL: quality of life; THA: total hip replacement; TKA: total knee replacement; SF39: short form 36; PCS: physical component score; MCS: mental component score; JSW: joint space width; OARSI: osteoarthritis research society international; HA: hyaluronic acid									

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome				
Rozendaal RM 2009 [341]	222	RCT	CR	2 years	KLG/WOMAC pain and function, JSN		KLG 1	KLG 2	
						WOMAC pain	-1.4 (95% CI [-6.7, 3.8])	-1.4 (95% CI [-7.4, 4.7])	
						WOMA C function	-0.09 mm (95% CI [-0.22, 0.04])	0.09 mm (95% CI [-0.07, 0.24])	
						JSN	-1.5 (95% CI [-5.8, 2.9])	-2.1 (95% CI [-7.5, 3.3])	
Hoeksma KL 2005 [342]	103	RCT	CR		KLG/Harris Hip score and range of motion in response to manual therapy vs exercise		Low radiographic grade	High radiographic grade	p
						HHS	MT -23.6 ET - 10.36	MT -16.01 ET - 3.56	p=0.52
						Hip function	MT 19.62 ET 5.07	MT 17.24 ET - 4.39	0.17
						ROM	MT 1.28 ET - 1.49	MT -0.38 ET - 3.02	p=0.01
Deshmukh A 2011 [343]	220	Cohort	CR	2 weeks	KLG/pain relief after methylprednisolone injections		Immediate pain relief	Delayed relief	
						KLG 2	2.16 p=0.03	1.14 (0.56,2.32)	
						KLG 3-4	3.94 p=0.001	5.33 (2.04,13.93)	
Robinson P 2007 [344]	120	Cohort	US CR	12 weeks	US osteophytes and capsular thickneing, KLG/WOMAC response	no baseline US or radiographic variable predictive of the outcome			

CR: conventional radiography; US: ultrasonography; KLG: Kellgren and Lawrence grade; WOMAC: Western Ontario McMaster Universities osteoarthritis index; JSN: joint space narrowing; HHS: Harris hip score; ROM: range of motion; MT: manual therapy; ET; exercise therapy; CS: corticosteroids

FOLLOW-UP OF OA - HIP

Q7: When, how often and under what clinical circumstance should imaging be used in the follow-up of osteoarthritic joints?

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Auleley G 2000 [345]	104	RCT	CR	3 years	Sensitivity to change of mJSW	SRM -0.70
Botha-Scheepers S 2008 [346]	115	Cohort	CR	2 years	Sensitivity to change of JSN score	SRM 0.43
Conrozier T 2009 [347]	50	RCT	CR	3 years	Sensitivity to change of JSN	SRM 2.05
Conrozier T 1998 [332]	61	Cohort	CR	81 months	Sensitivity to change of mean JSW, mJSW, JS area,	Mean JSW SRM 1 mJSW SRM 0.82 JS area 1.03
Conrozier T 1998 [348]	48	Cohort	CR	1 year	Sensitivity to change of JSW	SRM 1.15
Dougados M 2001 [349]	269	RCT	CR	3 years	Sensitivity to change of JSW	SRM 1
Gossec L 2009 [326]	1404	Cohort, RCT	CR	3 years	Sensitivity to change of JSW, KLG and OARSI	SRM: 0.77 JSW; 0.28 KLG; 0.35 OARSI
Maillefert JF 2002 [329]	25	Cohort	CR	3 years	Sensitivity to change of JSW (manual vs computer)	JSW: -0.77 (-1.03,-0.44) computer based mJSW -0.64 (-0.93,-0.29) computer based average JSW -0.61 (-0.9, -0.31)
Lequesne M 2002 [339]	95	RCT	CR	2 years	Sensitivity to change of JSW	mean (sd) JSW baseline 2.35 ±0.93 2 years 1.87 ± 1.10 in avocado soy/bean; baseline 2.50 ±0.94 2 years 1.90 ± 1.33
Maheu C 2005[350]	50	RCT	CR	3 years	Sensitivity to change of JSW	-0.76 (-0.77 , -0.73) (mean change)

CR: conventional radiography; mJSW: minimal joint space width; SRM: standardized response mean; JSW: joint space width; KLG: Kellgren and Lawrence grade; OARSI: osteoarthritis research society international

Reference	No of subjects	Study design	Imaging modality	Follow up	Outcome	
Maillefert JF 2002 [351]	25	Cohort	CR	5 years	Sensitivity to change of mJSW	Manual measure: -0.77 (-1.03, -0.44) Computer measure: -0.64 (-0.93, -0.29)
Papaloucas CD 2005 [352]	14	Cohort	CR	18 months	Sensitivity to change of mJSW	SRM=0.24
Pavelka K 2000 [353]	117	RCT	CR	5 years	Sensitivity to change of mJSW	mean change (SEM) 0-5 years: -0.21±0.08 treatment; -0.22±0.08 placebo
Ratzlaff C 2014 [354]	54	Case-control	CR	2 years	Sensitivity to change of mJSW	SRM 1.00
Jaremko J L 2014 [355]		SLR	CR			
Dougados M 1996 [336]	508	Cohort	CR	1 year	Sensitivity to change of mJSW	SRM 0.62
Lane NE 2004 [330]	745	Cohort	CR	8 years	Sensitivity to change of JSN	SRM 0.69
CR: conventional radiography; mJSW: minimal joint space width; SRM: standardized response mean; SEM: standard error; JSN: joint space narrowing						

References

1. Akgul O, Guldeste Z, Ozgocmen S. The reliability of the clinical examination for detecting Baker's cyst in asymptomatic fossa. *Int j rheum dis* 2014;17:204-9.
2. Esen S, Akarirmak U, Aydin FY, et al. Clinical evaluation during the acute exacerbation of knee osteoarthritis: the impact of diagnostic ultrasonography. *Rheumatol Int* 2013;33:711-7.
3. Ike RW, Somers EC, Arnold EL, et al. Ultrasound of the knee during voluntary quadriceps contraction: a technique for detecting otherwise occult effusions. *Arthritis Care Res* 2010;62:725-9.
4. Pendleton A, Millar A, O'Kane D, et al. Can sonography be used to predict the response to intra-articular corticosteroid injection in primary osteoarthritis of the knee? *Scand J Rheumatol* 2008;37:395-7.
5. Chatzopoulos D, Moralidis E, Markou P, et al. Baker's cysts in knees with chronic osteoarthritic pain: a clinical, ultrasonographic, radiographic and scintigraphic evaluation. *Rheumatol Int* 2008;29:141-6.
6. Lee CL, Huang MH, Chai CY, et al. The validity of in vivo ultrasonographic grading of osteoarthritic femoral condylar cartilage: a comparison with in vitro ultrasonographic and histologic gradings. *Osteoarthritis Cartilage* 2008;16:352-8.
7. Yoon CH, Kim HS, Ju JH, et al. Validity of the sonographic longitudinal sagittal image for assessment of the cartilage thickness in the knee osteoarthritis. *Clin rheumatol* 2008;27:1507-16.
8. Song IH, Althoff CE, Hermann KG, et al. Knee osteoarthritis. Efficacy of a new method of contrast-enhanced musculoskeletal ultrasonography in detection of synovitis in patients with knee osteoarthritis in comparison with magnetic resonance imaging. *Ann Rheum Dis* 2008;67:19-25.
9. Bhattacharya R, Kumar V, Safawi E, et al. The knee skyline radiograph: its usefulness in the diagnosis of patello-femoral osteoarthritis. *Int Orthop* 2007;31:247-52.
10. Chang CB, Seong SC, Kim TK. Evaluations of radiographic joint space - do they adequately predict cartilage conditions in the patellofemoral joint of the patients undergoing total knee arthroplasty for advanced knee osteoarthritis? *Osteoarthritis Cartilage* 2008;16:1160-6.
11. Dervin GF, Feibel RJ, Rody K, et al. 3-Foot standing AP versus 45 degrees PA radiograph for osteoarthritis of the knee. *Clin J Sport Med* 2001;11:10-6.
12. Waldstein W, Monsef JB, Buckup J, et al. The value of valgus stress radiographs in the workup for medial unicompartamental arthritis knee. *Clinical Orthop Relat Res* 2013;471:3998-4003.
13. Cicuttini FM, Wluka AE, Forbes A, et al. Comparison of tibial cartilage volume and radiologic grade of the tibiofemoral joint. *Arthritis Rheum* 2003;48:682-8.
14. Boegard T, Jonsson K. Radiography in osteoarthritis of the knee. *Skeletal Radiol* 1999;28:605-15.
15. Crema MD, Nogueira-Barbosa MH, Roemer FW, et al. Three-dimensional turbo spin-echo magnetic resonance imaging (MRI) and semiquantitative assessment of knee osteoarthritis: comparison with two-dimensional routine MRI. *Osteoarthritis Cartilage* 2013;21:428-33.
16. Gudbergesen H, Lohmander LS, Jones G, et al. Correlations between radiographic assessments and MRI features of knee osteoarthritis--a cross-sectional study. *Osteoarthritis Cartilage* 2013;21:535-43.
17. De Lange-Brokaar BJE, Ioan-Facsinay A, Yusuf E, et al. Degree of synovitis on MRI by comprehensive whole knee semi-quantitative scoring method correlates with histologic and macroscopic features of synovial tissue inflammation in knee osteoarthritis. *Osteoarthritis Cartilage* 2014;22:1606-13.
18. Fernandez-Madrid F, Karvonen RL, Teitge RA, et al. Synovial thickening detected by MR imaging in osteoarthritis of the knee confirmed by biopsy as synovitis. *Magnetic resonance imaging* 1995;13:177-83.

19. Bergman AG, Willen HK, Lindstrand AL, et al. Osteoarthritis of the knee: correlation of subchondral MR signal abnormalities with histopathologic and radiographic features. *Skeletal Radiol*. 1994;23:445-8.
20. Broderick LS, Turner DA, Renfrew DL, et al. Severity of articular cartilage abnormality in patients with osteoarthritis: evaluation with fast spin-echo MR vs arthroscopy. *AJR Am J Roentgenol* 1994;162:99-103.
21. Kalunian KC, Arnold WJ, Klashman DJ, et al. Can physical signs or magnetic resonance imaging substitute for diagnostic arthroscopy in knee osteoarthritis patients with suspected internal derangements? A pilot study. *J Clin Rheumatol* 2000;6:123-7.
22. Loeuille D, Sauliere N, Champigneulle J, et al. Comparing non-enhanced and enhanced sequences in the assessment of effusion and synovitis in knee OA: associations with clinical, macroscopic and microscopic features. *Osteoarthritis Cartilage* 2011;19:1433-9.
23. Saadat E, Jobke B, Chu B, et al. Diagnostic performance of in vivo 3-T MRI for articular cartilage abnormalities in human osteoarthritic knees using histology as standard of reference. *Euro Radiol* 2008;18:2292-302.
24. Takayama Y, Hatakenaka M, Tsushima H, et al. T1rho is superior to T2 mapping for the evaluation of articular cartilage denaturalization with osteoarthritis: radiological-pathological correlation after total knee arthroplasty. *Eur J Radiol* 2013;82:e192-8.
25. von Engelhardt LV, Lahner M, Klusmann A, et al. Arthroscopy vs. MRI for a detailed assessment of cartilage disease in osteoarthritis: diagnostic value of MRI in clinical practice. *BMC Musculoskelet Disord*. 2010;11:75.
26. Wong CS, Yan CH, Gong NJ, et al. Imaging biomarker with T1rho and T2 mappings in osteoarthritis - in vivo human articular cartilage study. *Eur J Radiol* 2013;82:647-50.
27. Yoshioka H, Stevens K, Hargreaves BA, et al. Magnetic resonance imaging of articular cartilage of the knee: Comparison between fat-suppressed three-dimensional SPGR imaging, fat-suppressed FSE imaging, and fat-suppressed three-dimensional DEFT imaging, and correlation with arthroscopy. *J Magn Reson Imaging*. 2004;20:857-64.
28. Zanetti M, Bruder E, Romero J, et al. Bone marrow edema pattern in osteoarthritic knees: correlation between MR imaging and histologic findings. *Radiology* 2000;215:835-40.
29. Graichen H, von Eisenhart-Rothe R, Vogl T, Englmeier KH, Eckstein F. Quantitative assessment of cartilage status in osteoarthritis by quantitative magnetic resonance imaging: technical validation for use in analysis of cartilage volume and further morphologic parameters. *Arthritis Rheum* 2004;50:811-6.
30. Brealey SD. Influence of magnetic resonance imaging of the knee on GPs' decisions: A randomised trial. *Br Journal Gen Pract* 2007;57:622-9.
31. Ritchie JF, Al-Sarawan M, Worth R, et al. A parallel approach: the impact of schuss radiography of the degenerate knee on clinical management. *Knee* 2004;11:283-7.
32. Barker K, Lamb SE, Toye F, et al. Association between radiographic joint space narrowing, function, pain and muscle power in severe osteoarthritis of the knee. *Clin Rehabil* 2004;18:793-800.
33. Cubukcu D, Sarsan A, Alkan H. Relationships between pain, function and radiographic findings in osteoarthritis of the knee: A cross-sectional study. *Arthritis*. 2012;1.
34. Bevers K, Bijlsma JW, Vriezekolk JE, et al. Ultrasonographic features in symptomatic osteoarthritis of the knee and relation with pain. *Rheumatology (Oxford)*. 2014;53:1625-9.
35. Neogi T, Nevitt MC, Yang M, et al. Consistency of knee pain: Correlates and association with function. *Osteoarthritis Cartilage* 2010;18:1250-5.

36. Sanghi D, Avasthi S, Mishra A, et al. Is radiology a determinant of pain, stiffness, and functional disability in knee osteoarthritis? A cross-sectional study. *J of Orthop Sci* 2011;16:719-25.
37. Madsen O BH, Egsmose C, Sylvest J. Isometric and isokinetic quadriceps strength in gonarthrosis; inter-relations between quadriceps strength, walking ability, radiology, subchondral bone density and pain. *Clinical Rheumatol* 1995;14:308-14.
38. Cicuttini FM, Baker J, Hart DJ, et al. Association of pain with radiological changes in different compartments and views of the knee joint. *Osteoarthritis Cartilage*. 1996;4(2):143-7.
39. Hernandez-Vaquero D, Fernandez-Carreira JM. Relationship between radiological grading and clinical status in knee osteoarthritis. a multicentric study. *BMC Musculoskel Disord* 2012;13
40. Lim BW, Hinman RS, Wrigley TV, et al. Varus malalignment and its association with impairments and functional limitations in medial knee osteoarthritis. *Arthritis Care Res* 2008;59:935-42.
41. Illingworth KD, El Bitar Y, Siewert K, et al. Correlation of WOMAC and KOOS scores to tibiofemoral cartilage loss on plain radiography and 3 Tesla MRI: data from the osteoarthritis initiative. *Knee Surg Sports Traumatol Arthrosc* 2014;22:1649-58.
42. Knoop J, Dekker J, Klein JP, et al. Biomechanical factors and physical examination findings in osteoarthritis of the knee: Associations with tissue abnormalities assessed by conventional radiography and high-resolution 3.0 Tesla magnetic resonance imaging. *Arthritis Res Ther* 2012;14(5).
43. Link TM SL, Ghosh MS, Ries M, Lu Y, Lane N, Majumdar S. Osteoarthritis: MR imaging findings in different stages of disease and correlation with clinical findings. *Radiology* 2003;226:373-81.
44. Hernigou P, Garabedian JM. Intercondylar notch width and the risk for anterior cruciate ligament rupture in the osteoarthritic knee: evaluation by plain radiography and CT scan. *Knee* 2002;9:313-6.
45. Janakiraman N, Teichtahl AJ, Wluka AE, Ding C, Jones G, Davis SR, et al. Static knee alignment is associated with the risk of unicompartmental knee cartilage defects. *J Orthop Res* 2008;26:225-30.
46. Issa SN, Dunlop D, Chang A, et al. Full-limb and knee radiography assessments of varus-valgus alignment and their relationship to osteoarthritis disease features by magnetic resonance imaging. *Arthritis Rheum* 2007;57:398-406.
47. Khan FA, Koff MF, Noisieux NO, et al. Effect of local alignment on compartmental patterns of knee osteoarthritis. *J Bone Joint Surg Am* 2008;90:1961-9.
48. Wang L, Vieira RL, Rybak LD, et al. Relationship between knee alignment and T1rho values of articular cartilage and menisci in patients with knee osteoarthritis. *Eur J Radiol* 2013;82:1946-52.
49. Friedrich KM, Shepard T, Chang G, et al. Does joint alignment affect the T2 values of cartilage in patients with knee osteoarthritis? *Eur Radiol* 2010;20:1532-8.
50. Teichtahl AJ, Wluka AE, Morris ME, et al. The relationship between the knee adduction moment and knee pain in middle-aged women without radiographic osteoarthritis. *J Rheumatol* 2006;33:1845-8.
51. von Eisenhart-Rothe R, Graichen H, et al. Femorotibial and patellar cartilage loss in patients prior to total knee arthroplasty, heterogeneity, and correlation with alignment of the knee. *Ann Rheum Dis* 2006;65:69-73.
52. Boissonneault A, Lynch JA, Wise BL, et al. Association of hip and pelvic geometry with tibiofemoral osteoarthritis: Multicenter Osteoarthritis Study (MOST). *Osteoarthritis Cartilage*. 2014;22:1129-35.
53. Ozgul A, Gunendi Z, Kesikburun S, et al. The association between patellar alignments features and tibiofemoral joint osteoarthritis. *Clin Rheumatol* 2013;32:1017-20.
54. van der Esch M, Knol DL, Schaffers IC, et al. Osteoarthritis of the knee: Multicompartmental or compartmental disease? *Rheumatology (Oxford)* 2014;53:540-6.

55. Cotofana S, Buck R, Wirth W, Roemer F, Duryea J, Nevitt M, et al. Cartilage thickening in early radiographic knee osteoarthritis: a within-person, between-knee comparison. *Arthritis care & research*. 2012;64:1681-90.
56. Ferraro B, Wilder FV, Leaveron PE. Site specific osteoarthritis and the index to ring finger length ratio. *Osteoarthritis Cartilage* 2010;18:354-7.
57. Holla JFM, Steultjens MPM, van der Leeden M, et al. Determinants of range of joint motion in patients with early symptomatic osteoarthritis of the hip and/or knee: An exploratory study in the CHECK cohort. *Osteoarthritis Cartilage*. 2011;19:411-9.
58. McDaniel G, Renner JB, Sloane R, et al. Association of knee and ankle osteoarthritis with physical performance. *Osteoarthritis Cartilage* 2011;19:634-8.
59. Ai F, Yu C, Zhang W, et al MR imaging of knee osteoarthritis and correlation of findings with reported patient pain. *J Univ Sci Technolog Med Sci* 2010;30:248-54.
60. Ballegaard C, Riis RGC, Bliddal H, Christensen R, Henriksen M, Bartels EM, et al. Knee pain and inflammation in the infrapatellar fat pad estimated by conventional and dynamic contrast-enhanced magnetic resonance imaging in obese patients with osteoarthritis: A cross-sectional study. *Osteoarthritis Cartilage*. 2014;22:933-40.
61. Lo GH, McAlindon TE, Niu J, et al. Bone marrow lesions and joint effusion are strongly and independently associated with weight-bearing pain in knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2009;17:1562-9.
62. Kornaat PR, Bloem JL, Ceulemans RY, et al. Osteoarthritis of the knee: association between clinical features and MR imaging findings. *Radiology* 2006;239:811-7.
63. Torres BA DD, Peterfy C, Guermazi A, et al. The relationship between specific tissue lesions and pain severity in persons with knee osteoarthritis. *Osteoarthritis Cartilage* 2006;14:1033-40.
64. Lange AK, Fiatarone Singh MA, Smith RM, Foroughi N, Baker MK, Shnier R, et al. Degenerative meniscus tears and mobility impairment in women with knee osteoarthritis. *Osteoarthritis Cartilage* 2007;15:701-8.
65. Bhattacharyya T, Gale D, Dewire P, et al. The clinical importance of meniscal tears demonstrated by magnetic resonance imaging in osteoarthritis of the knee. *J Bone Joint Surg Am* 2003;85-A:4-9.
66. Wenger A, Englund M, Wirth W, Hudelmaier M, Kwok K, Eckstein F. Relationship of 3D meniscal morphology and position with knee pain in subjects with knee osteoarthritis: a pilot study. *Eur Radiol* 2012;22:211-20.
67. Seah S, Wheaton D, Li L, Dyke JP, Talmo C, Harvey WF, et al. The relationship of tibial bone perfusion to pain in knee osteoarthritis. *Osteoarthritis Cartilage* 2012;20:1527-33.
68. Felson DT, Chaisson CE, Hill CL, Totterman SM, Gale ME, Skinner KM, et al. The association of bone marrow lesions with pain in knee osteoarthritis. *Ann Intern Med* 2001;134:541-9.
69. Sower MF, Hayes C, Jamadar D, Capul D, Lachance L, Jannausch M, et al. Magnetic resonance-detected subchondral bone marrow and cartilage defect characteristics associated with pain and X-ray-defined knee osteoarthritis. *Osteoarthritis Cartilage* 2003;11:387-93.
70. Sengupta M, Zhang YQ, Niu JB, et al. High signal in knee osteophytes is not associated with knee pain. *Osteoarthritis Cartilage* 2006;14:413-7.
71. Wluka AE, Wolfe R, Stuckey S, et al. How does tibial cartilage volume relate to symptoms in subjects with knee osteoarthritis? *Annals of the rheumatic diseases*. 2004;63:264-8.
72. Hernandez-Molina G, Neogi T, et al. The association of bone attrition with knee pain and other MRI features of osteoarthritis. *Ann Rheum Dis* 2008;67:43-7.
73. Baker K, Grainger A, Niu J, Clancy M, Guermazi A, Crema M, et al. Relation of synovitis to knee pain using contrast-enhanced MRIs. *Ann Rheum Dis* 2010;69:1779-83.

74. Hill CL, Gale DG, Chaisson CE, et al. Knee effusions, popliteal cysts, and synovial thickening: association with knee pain in osteoarthritis. *J Rheumatol* 2001;28:1330-7.
75. Hill CL, Seo GS, Gale D, Totterman S, Gale ME, Felson DT. Cruciate ligament integrity in osteoarthritis of the knee. *Arthritis Rheum* 2005;52:794-9.
76. Lindsey CT, Narasimhan A, Adolfo JM, Jin H, Steinbach LS, Link T, et al. Magnetic resonance evaluation of the interrelationship between articular cartilage and trabecular bone of the osteoarthritic knee. *Osteoarthritis Cartilage* 2004;12(2):86-96.
77. Cotofana S, Wyman BT, Benichou O, et al. Relationship between knee pain and the presence, location, size and phenotype of femorotibial denuded areas of subchondral bone as visualized by MRI. *Osteoarthritis Cartilage* 2013;21:1214-22.
78. Gale DR, Chaisson CE, Totterman SM, et al. Meniscal subluxation: association with osteoarthritis and joint space narrowing. *Osteoarthritis Cartilage* 1999;7:526-32.
79. Friedrich KM, Shepard T, de Oliveira VS, et al. T2 measurements of cartilage in osteoarthritis patients with meniscal tears. *AJR Am J Roentgenol.* 2009;193:W411-5.
80. Hovis KK, Stehling C, Souza RB, et al. Physical activity is associated with magnetic resonance imaging-based knee cartilage T2 measurements in asymptomatic subjects with and those without osteoarthritis risk factors. *Arthritis Rheum* 2011;63:2248-56.
81. Jungmann PM, Tham SC, Liebl H, et al. Association of trochlear dysplasia with degenerative abnormalities in the knee: data from the Osteoarthritis Initiative. *Skeletal Radiol* 2013;42:1383-92.
82. Kalichman L, Zhang Y, Niu J, et al. The association between patellar alignment and patellofemoral joint osteoarthritis features--an MRI study. *Rheumatology (Oxford).* 2007;46:1303-8.
83. Kornaat PR, Watt I, Riyazi N, Kloppenburg M, Bloem JL. The relationship between the MRI features of mild osteoarthritis in the patellofemoral and tibiofemoral compartments of the knee. *Eur Radiol* 2005;15:1538-43.
84. Krasnokutsky S, Belitskaya-Levy I, Bencardino J, et al. Quantitative magnetic resonance imaging evidence of synovial proliferation is associated with radiographic severity of knee osteoarthritis. *Arthritis Rheum* 2011;63:2983-91.
85. Lo GH, Hunter DJ, Nevitt M, et al. Strong association of MRI meniscal derangement and bone marrow lesions in knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2009;17:743-7.
86. Oda H, Igarashi M, Sase H, et al. Bone bruise in magnetic resonance imaging strongly correlates with the production of joint effusion and with knee osteoarthritis. *J Orthop Sci* 2008;13:7-15.
87. Stein V, Li L, Guermazi A, et al. The relation of femoral notch stenosis to ACL tears in persons with knee osteoarthritis. *Osteoarthritis Cartilage* 2010;18(2):192-9.
88. Naredo E, Cabero F, Palop MJ, et al. Ultrasonographic findings in knee osteoarthritis: a comparative study with clinical and radiographic assessment. *Osteoarthritis Cartilage* 2005;13:568-74.
89. D'Agostino MA, Conaghan P, Le Bars M, et al. EULAR report on the use of ultrasonography in painful knee osteoarthritis. Part 1: prevalence of inflammation in osteoarthritis. *Ann of Rheum Dis* 2005;64:1703-9.
90. de Miguel Mendieta E, Cobo Ibanez T, Uson Jaeger J, et al. Clinical and ultrasonographic findings related to knee pain in osteoarthritis. *Osteoarthritis Cartilage* 2006;14:540-4.
91. Hall M, Doherty S, Courtney P, et al. Synovial pathology detected on ultrasound correlates with the severity of radiographic knee osteoarthritis more than with symptoms. *Osteoarthritis Cartilage.* 2014;22:1627-33.

92. Iagnocco A, Meenagh G, Riente L, et al. Ultrasound imaging for the rheumatologist XXIX. Sonographic assessment of the knee in patients with osteoarthritis. *Clin Exp Rheumatol* 2010;28:643-6.
93. Wu PT, Shao CJ, Wu KC, et al. Pain in patients with equal radiographic grades of osteoarthritis in both knees: the value of gray scale ultrasound. *Osteoarthritis Cartilage* 2012;20:1507-13.
94. Kim HR, So Y, Moon SG, Lee IS, Lee SH. Clinical value of (99m)Tc-methylene diphosphonate (MDP) bone single photon emission computed tomography (SPECT) in patients with knee osteoarthritis. *Osteoarthritis Cartilage* 2008;16:212-8.
95. Kraus VB, McDaniel G, Worrell TW, et al. Association of bone scintigraphic abnormalities with knee malalignment and pain. *Ann Rheum Dis* 2009;68:1673-9.
96. McCrae F, Shouls J, Dieppe P, et al. Scintigraphic assessment of osteoarthritis of the knee joint. *Ann Rheum Dis* 1992;51:938-42.
97. Bruyere O, Honore A, Rovati LC, et al. Radiologic features poorly predict clinical outcomes in knee osteoarthritis. *Scand J Rheumatol* 2002;31:13-6.
98. Collins JE, Katz JN, Dervan EE, et al. Trajectories and risk profiles of pain in persons with radiographic, symptomatic knee osteoarthritis: Data from the osteoarthritis initiative. *Osteoarthritis Cartilage*. 2014;22:622-30.
99. Conaghan PG, D'Agostino MA, Le Bars M, et al. Clinical and ultrasonographic predictors of joint replacement for knee osteoarthritis: results from a large, 3-year, prospective EULAR study. *Annals of the rheumatic diseases*. 2010;69:644-7.
100. Podsiadlo P, Cicuttini FM, Wolski M, et al. Trabecular bone texture detected by plain radiography is associated with an increased risk of knee replacement in patients with osteoarthritis: a 6 year prospective follow up study. *Osteoarthritis Cartilage* 2014;22:71-5.
101. Cahue S, Dunlop D, Hayes K, et al. Varus-valgus alignment in the progression of patellofemoral osteoarthritis. *Arthritis Rheum* 2004;50:2184-90.
102. Cerejo R, Dunlop DD, Cahue S, et al. The influence of alignment on risk of knee osteoarthritis progression according to baseline stage of disease. *Arthritis Rheum* 2002;46:2632-6.
103. Brouwer GM, van Tol AW, Bergink AP, et al. Association between valgus and varus alignment and the development and progression of radiographic osteoarthritis of the knee. *Arthritis Rheum* 2007;56:1204-11.
104. Felson DT, Cooke TD, Niu J, et al. Can anatomic alignment measured from a knee radiograph substitute for mechanical alignment from full limb films? *Osteoarthritis Cartilage* 2009;17:1448-52.
105. Felson DT, Gale M, Niu J, et al. Osteophytes and progression of knee osteoarthritis. *Rheumatology (Oxford)* 2005;44:100-4.
106. Miyazaki T, Uchida K, Sato M, et al. Knee laxity after staircase exercise predicts radiographic disease progression in medial compartment knee osteoarthritis. *Arthritis Rheum* 2012;64:3908-16.
107. Moio K, Chang A, Eckstein F, et al. Varus-valgus alignment: reduced risk of subsequent cartilage loss in the less loaded compartment. *Arthritis and rheumatism*. 2011;63(4):1002-9.
108. Sharma L, Song J, Felson DT, et al. The role of knee alignment in disease progression and functional decline in knee osteoarthritis. *JAMA* 2001;286:188-95.
109. Cicuttini F, Wluka A, Hankin J, et al. Longitudinal study of the relationship between knee angle and tibiofemoral cartilage volume in subjects with knee osteoarthritis. *Rheumatology (Oxford)*. 2004;43:321-4.
110. Davies-Tuck M, Teichtahl AJ, Wluka AE, et al. Femoral sulcus angle and increased patella facet cartilage volume in an osteoarthritic population. *Osteoarthritis Cartilage* 2008;16:131-5.

111. Eckstein F, Le Graverand MP, Charles HC, et al. Clinical, radiographic, molecular and MRI-based predictors of cartilage loss in knee osteoarthritis. *Ann Rheum Dis* 2011;70:1223-30.
112. Bruyere O, Honore A, Ethgen O, et al. Correlation between radiographic severity of knee osteoarthritis and future disease progression. Results from a 3-year prospective, placebo-controlled study evaluating the effect of glucosamine sulfate. *Osteoarthritis Cartilage* 2003;11:1-5.
113. Felson D, Niu J, Sack B, et al. Progression of osteoarthritis as a state of inertia. *Ann Rheum Dis* 2013;72:924-9.
114. Woloszynski T, Podsiadlo P, Stachowiak GW, et al. Prediction of progression of radiographic knee osteoarthritis using tibial trabecular bone texture. *Arthritis Rheum* 2012;64:688-95.
115. Kraus VB, Feng S, Wang S, et al. Trabecular morphometry by fractal signature analysis is a novel marker of osteoarthritis progression. *Arthritis Rheum* 2009;60:3711-22.
116. Mazzuca SA, Brandt KD, Schauwecker DS, et al. Severity of joint pain and Kellgren-Lawrence grade at baseline are better predictors of joint space narrowing than bone scintigraphy in obese women with knee osteoarthritis. *J Rheumatol* 2005;32:1540-6.
117. Mazzuca SA, Brandt KD, Katz BP, et al. Risk factors for progression of tibiofemoral osteoarthritis: An analysis based on fluoroscopically standardised knee radiography. *Ann Rheum Dis* 2006;65:515-9.
118. Bijsterbosch J, Meulenbelt I, Watt I, et al. Clustering of hand osteoarthritis progression and its relationship to progression of osteoarthritis at the knee. *Ann Rheum Dis* 2014;73:567-72.
119. Neogi T, Nevitt M, Niu J, et al. Lack of association between chondrocalcinosis and increased risk of cartilage loss in knees with osteoarthritis: results of two prospective longitudinal magnetic resonance imaging studies. *Arthritis Rheumatism*. 2006;54:1822-8.
120. Buck RJ, Wirth W, Dreher D, et al. Frequency and spatial distribution of cartilage thickness change in knee osteoarthritis and its relation to clinical and radiographic covariates - data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2013;21:102-9.
121. Wirth W, Nevitt M, Hellio Le Graverand MP, et al. Lateral and medial joint space narrowing predict subsequent cartilage loss in the narrowed, but not in the non-narrowed femorotibial compartment--data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage* 2014;22:63-70.
122. Pelletier JP, Raynauld JP, Berthiaume MJ, et al. Risk factors associated with the loss of cartilage volume on weight-bearing areas in knee osteoarthritis patients assessed by quantitative magnetic resonance imaging: A longitudinal study. *Arthritis Res Ther* 2007;9.
123. Sharma L, Eckstein F, Song J, et al. Relationship of meniscal damage, meniscal extrusion, malalignment, and joint laxity to subsequent cartilage loss in osteoarthritic knees. *Arthritis Rheum* 2008;58:1716-26.
124. Wang Y, Wluka AE, Cicuttini FM. The determinants of change in tibial plateau bone area in osteoarthritic knees: a cohort study. *Arthritis Res Ther* 2005;7:R687-93.
125. Cicuttini F, Hankin J, Jones G, et al. Comparison of conventional standing knee radiographs and magnetic resonance imaging in assessing progression of tibiofemoral joint osteoarthritis. *Osteoarthritis Cartilage* 2005;13:722-7.
126. Tanamas S, Hanna FS, Cicuttini FM, et al. Does knee malalignment increase the risk of development and progression of knee osteoarthritis? A systematic review. *Arthritis Rheum* 2009;61:459-67.
127. Cicuttini FM, Jones G, Forbes A, et al. Rate of cartilage loss at two years predicts subsequent total knee arthroplasty: a prospective study. *Ann Rheum Dis* 2004;63:1124-7.

128. Eckstein F, Kwok CK, Boudreau RM, et al. Quantitative MRI measures of cartilage predict knee replacement: a case-control study from the Osteoarthritis Initiative. *Ann Rheum Dis* 2013;72:707-14.
129. Raynauld JP, Martel-Pelletier J, Haraoui B, et al. Risk factors predictive of joint replacement in a 2-year multicentre clinical trial in knee osteoarthritis using MRI: results from over 6 years of observation. *Ann Rheum Dis* 2011;70:1382-8.
130. Wluka AE, Ding C, Jones G, et al. The clinical correlates of articular cartilage defects in symptomatic knee osteoarthritis: a prospective study. *Rheumatology (Oxford)* 2005;44:1311-6.
131. Wang Y, Wluka AE, Berry PA, et al. Increase in vastus medialis cross-sectional area is associated with reduced pain, cartilage loss, and joint replacement risk in knee osteoarthritis. *Arthritis Rheum* 2012;64:3917-25.
132. Tanamas SK, Wluka AE, Pelletier JP, et al. The association between subchondral bone cysts and tibial cartilage volume and risk of joint replacement in people with knee osteoarthritis: a longitudinal study. *Arthritis Res Ther* 2010;12:R58.
133. Scher C, Craig J, Nelson F. Bone marrow edema in the knee in osteoarthrosis and association with total knee arthroplasty within a three-year follow-up. *Skeletal Radiol* 2008;37:609-17.
134. Eckstein F, Maschek S, Wirth W, et al. One year change of knee cartilage morphology in the first release of participants from the Osteoarthritis Initiative progression subcohort: association with sex, body mass index, symptoms and radiographic osteoarthritis status. *Ann Rheum Dis* 2009;68:674-9.
135. Amin S, Guermazi A, Lavalley MP, et al. Complete anterior cruciate ligament tear and the risk for cartilage loss and progression of symptoms in men and women with knee osteoarthritis. *Osteoarthritis Cartilage* 2008;16:897-902.
136. Everhart JS, Siston RA, Flanigan DC. Tibiofemoral subchondral surface ratio (SSR) is a predictor of osteoarthritis symptoms and radiographic progression: data from the Osteoarthritis Initiative (OAI). *Osteoarthritis Cartilage* 2014;22:771-8.
137. Moio K, Eckstein F, Chmiel JS, et al. Denuded subchondral bone and knee pain in persons with knee osteoarthritis. *Arthritis Rheum* 2009;60:3703-10.
138. Pelletier JP, Cooper C, Peterfy C, et al. What is the predictive value of MRI for the occurrence of knee replacement surgery in knee osteoarthritis? *Ann Rheum Dis* 2013;72:1594-604.
139. Hill CL, Hunter DJ, Niu J, et al. Synovitis detected on magnetic resonance imaging and its relation to pain and cartilage loss in knee osteoarthritis. *Ann Rheum Dis* 2007;66:1599-603.
140. Berthiaume MJ, Raynauld JP, Martel-Pelletier J, et al. Meniscal tear and extrusion are strongly associated with progression of symptomatic knee osteoarthritis as assessed by quantitative magnetic resonance imaging. *Ann Rheum Dis* 2005;64:556-63.
141. Chang A, Moio K, Chmiel JS, et al. Subregional effects of meniscal tears on cartilage loss over 2 years in knee osteoarthritis. *Ann Rheum Dis* 2011;70:74-9.
142. Crema MD, Hunter DJ, Burstein D, et al. Delayed gadolinium-enhanced magnetic resonance imaging of medial tibiofemoral cartilage and its relationship with meniscal pathology: a longitudinal study using 3.0T magnetic resonance imaging. *Arthritis Rheum* 2014;66:1517-24.
143. Wang Y, Wluka AE, Pelletier JP, et al. Meniscal extrusion predicts increases in subchondral bone marrow lesions and bone cysts and expansion of subchondral bone in osteoarthritic knees. *Rheumatology (Oxford)* 2010;49:997-1004.
144. Raynauld JP, Martel-Pelletier J, Berthiaume MJ, et al. Long term evaluation of disease progression through the quantitative magnetic resonance imaging of symptomatic knee

- osteoarthritis patients: correlation with clinical symptoms and radiographic changes. *Arthritis Res Ther* 2006;8:R21.
145. Hunter DJ, Zhang YQ, Niu JB, et al. The association of meniscal pathologic changes with cartilage loss in symptomatic knee osteoarthritis. *Arthritis Rheum* 2006;54:795-801.
 146. Crema MD, Cibere J, Sayre EC, et al. The relationship between subchondral sclerosis detected with MRI and cartilage loss in a cohort of subjects with knee pain: The knee osteoarthritis progression (KOAP) study. *Osteoarthritis Cartilage*. 2014;22:540-6.
 147. Eckstein F, Wirth W, Hudelmaier MI, et al. Relationship of compartment-specific structural knee status at baseline with change in cartilage morphology: a prospective observational study using data from the osteoarthritis initiative. *Arthritis Res Ther* 2009;11:R90.
 148. Driban JB, Lo GH, Lee J, et al. Quantitative bone marrow lesion size in osteoarthritic knees correlates with cartilage damage and predicts longitudinal cartilage loss. *BMC musculoskeletal disord* 2011;12
 149. Carrino JA, Blum J, Parellada JA, et al. MRI of bone marrow edema-like signal in the pathogenesis of subchondral cysts. *Osteoarthritis Cartilage* 2006;14:1081-5.
 150. Hunter DJ, Zhang Y, Niu J, et al. Increase in bone marrow lesions associated with cartilage loss: a longitudinal magnetic resonance imaging study of knee osteoarthritis. *Arthritis Rheum* 2006;54:1529-35.
 151. Hernandez-Molina G, Guermazi A, Niu J, et al. Central bone marrow lesions in symptomatic knee osteoarthritis and their relationship to anterior cruciate ligament tears and cartilage loss. *Arthritis Rheum* 2008;58:130-6.
 152. Felson DT, McLaughlin S, Goggins J, et al. Bone marrow edema and its relation to progression of knee osteoarthritis. *Ann Intern Med* 2003;139:330-6.
 153. Kothari A, Guermazi A, Chmiel JS, et al. Within-subregion relationship between bone marrow lesions and subsequent cartilage loss in knee osteoarthritis. *Arthritis Care Res* 2010;62:198-203.
 154. Felson DT, Lynch J, Guermazi A, et al. Comparison of BLOKS and WOMBS scoring systems part II. Longitudinal assessment of knee MRIs for osteoarthritis and suggested approach based on their performance: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage* 2010;18(1):1402-7.
 155. Wluka AE, Stuckey S, Snaddon J, et al. The determinants of change in tibial cartilage volume in osteoarthritic knees. *Arthritis Rheum* 2002;46:2065-72.
 156. Madan-Sharma R, Kloppenburg M, Kornaat PR, et al. Do MRI features at baseline predict radiographic joint space narrowing in the medial compartment of the osteoarthritic knee 2 years later? *Skeletal Radiol* 2008;37:805-11.
 157. Yusuf E, Kortekaas MC, Watt I, et al. Do knee abnormalities visualised on MRI explain knee pain in knee osteoarthritis? A systematic review. *Ann Rheum Dis* 2011;70:60-7.
 158. Dieppe P, Cushnaghan J, Young P, et al. Prediction of the progression of joint space narrowing in osteoarthritis of the knee by bone scintigraphy. *Ann Rheum Dis* 1993;52:557-63.
 159. Barrett JP, Jr., Rashkoff E, Sirna EC, et al. Correlation of roentgenographic patterns and clinical manifestations of symptomatic idiopathic osteoarthritis of the knee. *Clin Orthop Relat Res* 1990:179-83.
 160. Gaffney K, Ledingham J, Perry JD. Intra-articular triamcinolone hexacetonide in knee osteoarthritis: Factors influencing the clinical response. *Ann Rheum Dis* 1995;54:379-81.
 161. Gudbergsen H, Boesen M, Lohmander LS, et al. Weight loss is effective for symptomatic relief in obese subjects with knee osteoarthritis independently of joint damage severity assessed by high-field MRI and radiography. *Osteoarthritis Cartilage* 2012;20:495-502.

162. Gudbergesen H, Boesen M, Christensen R, et al. Radiographs and low field MRI (0.2T) as predictors of efficacy in a weight loss trial in obese women with knee osteoarthritis. *BMC Musculoskel Disord*. 2011;12:56.
163. Hellio le Graverand MP, Clemmer RS, et al. A 2-year randomised, double-blind, placebo-controlled, multicentre study of oral selective iNOS inhibitor, cindunistat (SD-6010), in patients with symptomatic osteoarthritis of the knee. *Ann Rheum Dis*. 2013;72:187-95.
164. Keurentjes JC, Fiocco M, So-Osman C, et al. Patients with Severe Radiographic Osteoarthritis Have a Better Prognosis in Physical Functioning after Hip and Knee Replacement: A Cohort-Study. *PLoS one* 2013;8.
165. Case JP, Baliunas AJ, Block JA. Lack of efficacy of acetaminophen in treating symptomatic knee osteoarthritis: A randomized, double-blind, placebo-controlled comparison trial with diclofenac sodium. *Arch Intern Med* 2003;163:169-78.
166. Sawitzke AD, Shi H, Finco MF, et al. The effect of glucosamine and/or chondroitin sulfate on the progression of knee osteoarthritis: A report from the glucosamine/chondroitin arthritis intervention trial. *Arthritis Rheum* 2008;58:3183-91.
167. Mazzuca SA, Brandt KD, Chakr R, et al. Varus malalignment negates the structure-modifying benefits of doxycycline in obese women with knee osteoarthritis. *Osteoarthritis Cartilage*. 2010;18:1008-11.
168. Toh EM, Prasad PS, Teanby D. Correlating the efficacy of knee viscosupplementation with osteoarthritic changes on roentgenological examination. *Knee*. 2002;9:321-30.
169. Chao J, Wu C, Sun B, Hose MK, et al. Inflammatory characteristics on ultrasound predict poorer longterm response to intraarticular corticosteroid injections in knee osteoarthritis. *J Rheumatol* 2010;37:650-5.
170. Hirsch G, Kitas G, Klocke R. Intra-articular corticosteroid injection in osteoarthritis of the knee and hip: factors predicting pain relief--a systematic review. *Semin Arthritis Rheum* 2013;42:451-73.
171. Anandacoomarasamy A, Bagga H, Ding C, et al. Predictors of clinical response to intraarticular hylan injections - A prospective study using synovial fluid measures, clinical outcomes, and magnetic resonance imaging. *J Rheumatol* 2008;35:685-90.
172. Knoop J, Dekker J, van der Leeden M, et al. Is the severity of knee osteoarthritis on magnetic resonance imaging associated with outcome of exercise therapy? *Arthritis Care Res* 2014;66(1):63-8.
173. Pessis E, Drape JL, Ravaud P, et al. Assessment of progression in knee osteoarthritis: results of a 1 year study comparing arthroscopy and MRI. *Osteoarthritis Cartilage* 2003;11:361-9.
174. Conrozier T, Mathieu P, Piperno M, et al. Selection of knee radiographs for trials of structure-modifying drugs in patients with knee osteoarthritis: A prospective, longitudinal study of Lyon schuss knee radiographs with the definition of adequate alignment of the medial tibial plateau. *Arthritis Rheum* 2005;52:1411-7.
175. Hellio Le Graverand MP, Vignon EP, et al. Head-to-head comparison of the Lyon Schuss and fixed flexion radiographic techniques. Long-term reproducibility in normal knees and sensitivity to change in osteoarthritic knees. *Ann Rheum Dis* 2008;67:1562-6.
176. Mazzuca SA, Hellio Le Graverand MP, Vignon E, et al. Performance of a non-fluoroscopically assisted substitute for the Lyon schuss knee radiograph: quality and reproducibility of positioning and sensitivity to joint space narrowing in osteoarthritic knees. *Osteoarthritis Cartilage* 2008;16(12):1555-9.

177. Piperno M, Graverand MPHL, Conrozier T, et al. Quantitative evaluation of joint space width in femorotibial osteoarthritis: Comparison of three radiographic views. *Osteoarthritis Cartilage* 1998;6:252-9.
178. Spector TD CP, Buckland-Wright JC, Garner P, et al. Effect of risedronate on joint structure and symptoms of knee osteoarthritis: results of the BRISK randomized, controlled trial. *Arthritis Res Ther* 2005;7:R625-33.
179. Mazzuca SA, Brandt KD, Buckwalter KA. Detection of radiographic joint space narrowing in subjects with knee osteoarthritis: Longitudinal comparison of the metatarsophalangeal and semiflexed anteroposterior views. *Arthritis Rheum* 2003;48:385-90.
180. Botha-Scheepers S, Kloppenburg M, Kroon HM, et al. Fixed-flexion knee radiography: the sensitivity to detect knee joint space narrowing in osteoarthritis. *Osteoarthritis Cartilage*. 2007;15:350-3.
181. Cicuttini FM, Wluka AE, Hankin J, et al. Comparison of patella cartilage volume and radiography in the assessment of longitudinal joint change at the patellofemoral joint. *J Rheumatol* 2004;31:1369-72.
182. Hellio Le Graverand MP, Buck RJ, et al. Change in regional cartilage morphology and joint space width in osteoarthritis participants versus healthy controls: a multicentre study using 3.0 Tesla MRI and Lyon-Schuss radiography. *Ann Rheum Dis* 2010;69:155-62.
183. Mazzuca SA, Brandt KD, Dieppe PA, et al. Effect of alignment of the medial tibial plateau and x-ray beam on apparent progression of osteoarthritis in the standing anteroposterior knee radiograph. *Arthritis Rheum* 2001;44:1786-94.
184. Pavelka K FS, Olejarova M, Gatterova J, et al. Hyaluronic acid levels may have predictive value for the progression of knee osteoarthritis. *Osteoarthritis Cartilage*. 2004;12:277-83.
185. Vignon E, Piperno M, Le Graverand MPH, et al. Measurement of radiographic joint space width in the tibiofemoral compartment of the osteoarthritic knee: Comparison of standing anteroposterior and Lyon schuss views. *Arthritis Rheum* 2003;48:378-84.
186. Wirth W, Duryea J, Hellio Le Graverand MP, et al. Direct comparison of fixed flexion, radiography and MRI in knee osteoarthritis: responsiveness data from the Osteoarthritis Initiative. *OsteoarthritisCartilage* 2013;21:117-25.
187. Boegard TL, Rudling O, Petersson IF, et al. Distribution of MR-detected cartilage defects of the patellofemoral joint in chronic knee pain. *Osteoarthritis Cartilage* 2003;11:494-8.
188. Mazzuca SA, Brandt KD, Buckwalter KA, et al. Pitfalls in the accurate measurement of joint space narrowing in semiflexed, anteroposterior radiographic imaging of the knee. *Arthritis Rheum* 2004;50:2508-15.
189. Miyazaki T WM, Kawahara H, Sato M, et al. Dynamic load at baseline can predict radiographic disease progression in medial compartment knee osteoarthritis. *Ann Rheum Dis* 2002;61:617-22.
190. Bruyere O, Henrotin YE, Honore A, et al. Impact of the joint space width measurement method on the design of knee osteoarthritis studies. *Aging Clin Exp Res* 2003;15:136-41.
191. Gossec L, Jordan JM, Mazzuca SA, et al. Comparative evaluation of three semi-quantitative radiographic grading techniques for knee osteoarthritis in terms of validity and reproducibility in 1759 X-rays: report of the OARSI-OMERACT task force. Extended report. *Osteoarthritis Cartilage*. 2008;16:742-8.
192. Lanyon P, Jones A, Doherty. Assessing progression of patello femoral osteoarthritis: A comparison between two radiographic methods. *Ann Rheum Dis* 1996;55:875-9.

193. LaValley MP, McLaughlin S, Goggins J, et al. The lateral view radiograph for assessment of the tibiofemoral joint space in knee osteoarthritis: its reliability, sensitivity to change, and longitudinal validity. *Arthritis Rheum* 2005;52:3542-7.
194. Nevitt MC, Peterfy C, Guermazi A, et al. Longitudinal performance evaluation and validation of fixed-flexion radiography of the knee for detection of joint space loss. *Arthritis Rheum* 2007;56:1512-20.
195. Reginster JY, Deroisy R, Rovati LC, et al. Long-term effects of glucosamine sulphate on osteoarthritis progression: A randomised, placebo-controlled clinical trial. *Lancet* 2001;357:251-6.
196. Sugiyama S IM, Suzuki Y, Shimizu K. procollagen II C propeptide level in the synovial fluid as a predictor of radiographic progression in early knee osteoarthritis. *Ann Rheum Dis* 2003;62:27-32.
197. Reichmann WM, Katz JN, Losina E. Differences in self-reported health in the osteoarthritis Initiative (OAI) and Third national health and nutrition Examination survey (NHANES-III). *PloS one*. 2011;6.
198. Duryea J, Neumann G, Niu J, Totterman S, et al. Comparison of radiographic joint space width with magnetic resonance imaging cartilage morphometry: Analysis of longitudinal data from the osteoarthritis initiative. *Arthritis Care Res* 2010;62:932-7.
199. Bartlett SJ, Ling SM, Mayo NE, et al. Identifying common trajectories of joint space narrowing over two years in knee osteoarthritis. *Arthritis Care Res* 2011;63:1722-8.
200. Bruyere O, Genant H, Kothari M, Zaim S, White D, Peterfy C, et al. Longitudinal study of magnetic resonance imaging and standard X-rays to assess disease progression in osteoarthritis. *Osteoarthritis Cartilage* 2007;15:98-103.
201. Crema MD, Nevitt MC, Guermazi A, et al. Progression of cartilage damage and meniscal pathology over 30 months is associated with an increase in radiographic tibiofemoral joint space narrowing in persons with knee OA - the MOST study. *Osteoarthritis Cartilage* 2014;22:1743-7.
202. Fukui N, Yamane S, Ishida S, et al. Relationship between radiographic changes and symptoms or physical examination findings in subjects with symptomatic medial knee osteoarthritis: A three-year prospective study. *BMC Musculoskel Disord* 2010;11
203. Teichtahl AJ, Wluka AE, Wang Y, et al. Obesity and adiposity are associated with the rate of patella cartilage volume loss over 2 years in adults without knee osteoarthritis. *AnnaRheum Dis* 2009;68:909-13.
204. Teichtahl AJ, Wluka AE, Cicuttini FM. Frontal plane knee alignment is associated with a longitudinal reduction in patella cartilage volume in people with knee osteoarthritis. *Osteoarthritis Cartilage* 2008;16:851-4.
205. Eckstein F, Buck RJ, Burstein D, et al. Precision of 3.0 Tesla quantitative magnetic resonance imaging of cartilage morphology in a multicentre clinical trial. *Ann Rheum Dis* 2008;67:1683-8.
206. Eckstein F, Benichou O, Wirth W, et al. Magnetic resonance imaging-based cartilage loss in painful contralateral knees with and without radiographic joint space narrowing: Data from the Osteoarthritis Initiative. *Arthritis Rheum* 2009;61:1218-25.
207. Blumenkrantz G, Lindsey CT, Dunn TC, et al. A pilot, two-year longitudinal study of the interrelationship between trabecular bone and articular cartilage in the osteoarthritic knee. *Osteoarthritis Cartilage* 2004;12:997-1005.
208. Eckstein F, Nevitt M, Gimona A, et al. Rates of change and sensitivity to change in cartilage morphology in healthy knees and in knees with mild, moderate, and end-stage radiographic osteoarthritis: results from 831 participants from the Osteoarthritis Initiative. *Arthritis Care Res* 2011;63:311-9.

209. Hunter DJ, Niu J, Zhang Y, Totterman S, Tamez J, Dabrowski C, et al. Change in cartilage morphometry: a sample of the progression cohort of the Osteoarthritis Initiative. *Ann Rheum Dis* 2009;68:349-56.
210. Hunter DJ, Li L, Zhang YQ, et al. Region of interest analysis: by selecting regions with denuded areas can we detect greater amounts of change? *Osteoarthritis Cartilage* 2010;18:175-83.
211. Maschek S, Wirth W, Ladel C, et al. Rates and sensitivity of knee cartilage thickness loss in specific central reading radiographic strata from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2014;22:1550-3.
212. Cromer MS, Bourne RM, Fransen M, et al. Responsiveness of quantitative cartilage measures over one year in knee osteoarthritis: comparison of radiography and MRI assessments. *J magn reson imaging* 2014;39:103-9.
213. Buck RJ, Wyman BT, Le Graverand MP, et al. Osteoarthritis may not be a one-way-road of cartilage loss--comparison of spatial patterns of cartilage change between osteoarthritic and healthy knees. *Osteoarthritis Cartilage* 2010;18:329-35.
214. Eckstein F, Kunz M, Schutzer M, et al. Two year longitudinal change and test-retest-precision of knee cartilage morphology in a pilot study for the osteoarthritis initiative. *Osteoarthritis Cartilage* 2007;15:1326-32.
215. Hudelmaier M, Wirth W, Wehr B, et al. Femorotibial cartilage morphology: reproducibility of different metrics and femoral regions, and sensitivity to change in disease. *Cells Tissues Organs* 2010;192:340-50.
216. Iranpour-Boroujeni T, Watanabe A, Bashtar R, et al. Quantification of cartilage loss in local regions of knee joints using semi-automated segmentation software: analysis of longitudinal data from the Osteoarthritis Initiative (OAI). *Osteoarthritis Cartilage*. 2011;19:309-14.
217. Raynauld JP, Martel-Pelletier J, Berthiaume MJ, et al. Quantitative magnetic resonance imaging evaluation of knee osteoarthritis progression over two years and correlation with clinical symptoms and radiologic changes. *Arthritis and rheumatism*. 2004;50:476-87.
218. Raynauld JP, Martel-Pelletier J, Berthiaume MJ, et al. Correlation between bone lesion changes and cartilage volume loss in patients with osteoarthritis of the knee as assessed by quantitative magnetic resonance imaging over a 24-month period. *Ann Rheum Dis* 2008;67:683-8.
219. Raynauld JP, Martel-Pelletier J, Abram F, et al. Analysis of the precision and sensitivity to change of different approaches to assess cartilage loss by quantitative MRI in a longitudinal multicentre clinical trial in patients with knee osteoarthritis. *Arthritis Res Ther* 2008;10:R129.
220. Raynauld JP M-pJ, Bias P, Laufer S, et al. Protective effects of licofelone, a 5-lipoxygenase and cyclo-oxygenase inhibitor, versus naproxen on cartilage loss in knee osteoarthritis: a first multicentre clinical trial using quantitative MRI. *Ann Rheum Dis* 2009;68:938-47.
221. Eckstein F, McCulloch CE, Lynch JA, et al. How do short-term rates of femorotibial cartilage change compare to long-term changes? Four year follow-up data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2012;20:1250-7.
222. Amin S, LaValley MP, Guermazi A, et al. The relationship between cartilage loss on magnetic resonance imaging and radiographic progression in men and women with knee osteoarthritis. *Arthritis Rheum* 2005;52:3152-9.
223. Gandy SJ, Dieppe PA, Keen MC, et al. No loss of cartilage volume over three years in patients with knee osteoarthritis as assessed by magnetic resonance imaging. *Osteoarthritis and cartilage* 2002;10:929-37.

224. Pelletier JP, Raynauld JP, Abram F, et al. A new non-invasive method to assess synovitis severity in relation to symptoms and cartilage volume loss in knee osteoarthritis patients using MRI. *Osteoarthritis Cartilage* 2008;16 Suppl 3:S8-13.
225. Brandt KD, Mazucca SA, Buckwalter KA. Acetaminophen, like conventional NSAIDs, may reduce synovitis in osteoarthritic knees. *Rheumatology (Oxford)* 2006;45:1389-94.
226. Driban JB, Lo GH, Price L, et al. Bone marrow lesion volume reduction is not associated with improvement of other periarticular bone measures: data from the Osteoarthritis Initiative. *Arthritis Res Ther* 2013;15:R153.
227. Eckstein F, Boudreau RM, Wang Z, et al. Trajectory of cartilage loss within 4 years of knee replacement - a nested case-control study from the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2014;22:1542-9.
228. Felson DT, Parkes MJ, Marjanovic EJ, et al. Bone marrow lesions in knee osteoarthritis change in 6-12 weeks. *Osteoarthritis Cartilage* 2012;20(12):1514-8.
229. Hunter DJ, Conaghan PG, Peterfy CG, et al. Responsiveness, effect size, and smallest detectable difference of Magnetic Resonance Imaging in knee osteoarthritis. *Osteoarthritis Cartilage* 2006;14 Suppl A:A112-5.
230. Hunter DJ, Bowes MA, Eaton CB, et al. Can cartilage loss be detected in knee osteoarthritis (OA) patients with 3-6 months' observation using advanced image analysis of 3T MRI? *Osteoarthritis Cartilage* 2010;18:677-83.
231. Hunter DJ, Zhang W, Conaghan PG, et al. Responsiveness and reliability of MRI in knee osteoarthritis: a meta-analysis of published evidence. *Osteoarthritis Cartilage* 2011;19:589-605.
232. Kubota M, Ishijima M, Kurosawa H, et al. A longitudinal study of the relationship between the status of bone marrow abnormalities and progression of knee osteoarthritis. *J Orthop Sci* 2010;15:641-6.
233. Crema MD, Hunter DJ, Burstein D, et al. Association of changes in delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) with changes in cartilage thickness in the medial tibiofemoral compartment of the knee: a 2 year follow-up study using 3.0 T MRI. *Ann Rheum Dis* 2014;73:1935-41.
234. Kornaat PR, Kloppenburg M, Sharma R, et al. Bone marrow edema-like lesions change in volume in the majority of patients with osteoarthritis; associations with clinical features. *Eur Radiol* 2007;17:3073-8.
235. Phan CM, Link TM, Blumenkrantz G, et al. MR imaging findings in the follow-up of patients with different stages of knee osteoarthritis and the correlation with clinical symptoms. *Eur radiol* 2006;16:608-18.
236. Cubukcu D, Ardic F, Karabulut N, et al. Hylan G-F 20 efficacy on articular cartilage quality in patients with knee osteoarthritis: clinical and MRI assessment. *Clin Rheumatol* 2005;24:336-41.
237. Stahl R, Blumenkrantz G, Carballido-Gamio J, et al. MRI-derived T2 relaxation times and cartilage morphometry of the tibio-femoral joint in subjects with and without osteoarthritis during a 1-year follow-up. *Osteoarthritis Cartilage*. 2007;15:1225-34.
238. Stahl R, Jain SK, Lutz J, Wyman BT, et al. Osteoarthritis of the knee at 3.0 T: comparison of a quantitative and a semi-quantitative score for the assessment of the extent of cartilage lesion and bone marrow edema pattern in a 24-month longitudinal study. *Skeletal radiol* 2011;40:1315-27.
239. Wirth W, Hellio Le Graverand MP, Wyman BT, et al. Regional analysis of femorotibial cartilage loss in a subsample from the Osteoarthritis Initiative progression subcohort. *Osteoarthritis Cartilage* 2009;17:291-7.

240. Wirth W, Larroque S, Davies RY, et al. Comparison of 1-year vs 2-year change in regional cartilage thickness in osteoarthritis results from 346 participants from the Osteoarthritis Initiative. *Osteoarthritis Cartilage* 2011;19:74-83.
241. Wirth W, Buck R, Nevitt M, Le Graverand MP, et al. MRI-based extended ordered values more efficiently differentiate cartilage loss in knees with and without joint space narrowing than region-specific approaches using MRI or radiography--data from the OA initiative. *Osteoarthritis Cartilage* 2011;19:689-99.
242. Wirth W, Benichou O, Kwok CK, et al. Spatial patterns of cartilage loss in the medial femoral condyle in osteoarthritic knees: data from the Osteoarthritis Initiative. *Magn Reson Med* 2010;63:574-81.
243. Zhang Y, Nevitt M, Niu J, et al. Fluctuation of knee pain and changes in bone marrow lesions, effusions, and synovitis on magnetic resonance imaging. *Arthritis Rheum* 2011;63:691-9.
244. Creamer P, Sharif M, George E, et al. Intra-articular hyaluronic acid in osteoarthritis of the knee: an investigation into mechanisms of action. *Osteoarthritis Cartilage* 1994;2:133-40.
245. Hall M, Doherty S, Courtney P, et al. Ultrasound detected synovial change and pain response following intra-articular injection of corticosteroid and a placebo in symptomatic osteoarthritic knees: a pilot study. *Ann Rheum Dis* 2014;73:1590-1.
246. Acebes JC, Sanchez-Pernaute O, Diaz-Oca A, et al. Ultrasonographic assessment of Baker's cysts after intra-articular corticosteroid injection in knee osteoarthritis. *J Clin Ultrasound* 2006;34:113-7.
247. Bandinelli F, Fedi R, Generini S, et al. Longitudinal ultrasound and clinical follow-up of Baker's cysts injection with steroids in knee osteoarthritis. *Clin Rheumatol* 2012;31:727-31.
248. Hall M, Doherty S, Courtney P, et al. Synovial pathology detected on ultrasound correlates with the severity of radiographic knee osteoarthritis more than with symptoms. *Osteoarthritis Cartilage*. 2014;22:1627-33.
249. Jan MH, Chai HM, Wang CL, et al. Effects of repetitive shortwave diathermy for reducing synovitis in patients with knee osteoarthritis: an ultrasonographic study. *Phys Ther* 2006;86:236-44.
250. Kawaguchi K, Enokida M, Otsuki R, et al. Ultrasonographic evaluation of medial radial displacement of the medial meniscus in knee osteoarthritis. *Arthritis Rheum* 2012;64:173-80.
251. Song IH, Althoff CE, Hermann KG, et al. Contrast-enhanced ultrasound in monitoring the efficacy of a bradykinin receptor 2 antagonist in painful knee osteoarthritis compared with MRI. *Ann Rheum Dis* 2009;68:75-83.
252. Keen HI, Mease PJ, Bingham CO, 3rd, et al. Systematic review of MRI, ultrasound, and scintigraphy as outcome measures for structural pathology in interventional therapeutic studies of knee arthritis: focus on responsiveness. *Journal Rheumatol* 2011;38:142-54.
253. Bum Park Y, Ah Choi W, Kim YK, et al. Accuracy of blind versus ultrasound-guided suprapatellar bursal injection. *J Clin Ultrasound* 2012;40:20-5.
254. Im SH, Lee SC, Park YB, et al. Feasibility of sonography for intra-articular injections in the knee through a medial patellar portal. *J Ultrasound Med* 2009;28:1465-70.
255. Jang SH, Lee SC, Lee JH, et al. Comparison of ultrasound (US)-guided intra-articular injections by in-plain and out-of-plain on medial portal of the knee. *Rheumatol Int* 2013;33:1951-9.
256. Sibbitt Jr WL, Band PA, Kettwich LG, et al. A randomized controlled trial evaluating the cost-effectiveness of sonographic guidance for intra-articular injection of the osteoarthritic knee. *J Clin Rheumatol* 2011;17:409-15.
257. Maricar N, Parkes MJ, Callaghan MJ, et al. Where and how to inject the knee--a systematic review. *Semin Arthritis Rheum* 2013;43:195-203.

258. Matsos M, Harish S, Zia P, et al. Ultrasound of the hands and feet for rheumatological disorders: Influence on clinical diagnostic confidence and patient management. *Skeletal Radiol* 2009;38:1049-54.
259. Moon JS, Lee K, Lee HS, Lee WC. Cartilage lesions in anterior bony impingement of the ankle. *Arthroscopy* 2010;26:984-9.
260. Tol JL, Verhagen RAW, Krips R, et al. The anterior ankle impingement syndrome: Diagnostic value of oblique radiographs. *Foot Ankle Int* 2004;25:63-8.
261. Bouaicha S, Ehrmann C, Moor BK, et al. Radiographic analysis of metatarsus primus elevatus and hallux rigidus. *Foot Ankle Int* 2010;31:807-14.
262. Iagnocco A, Filippucci E, Riente L, et al. Ultrasound imaging for the rheumatologist XXXV. Sonographic assessment of the foot in patients with osteoarthritis. *Clinical Exp Rheumatol* 2011;29:757-62.
263. Keen HI, Redmond A, Wakefield RJ, et al. An ultrasonographic study of metatarsophalangeal joint pain: synovitis, structural pathology and their relationship to symptoms and function. *Ann Rheum Dis* 2011;70:2140-3.
264. Menz HB, Munteanu SE, Landorf KB, et al. Radiographic evaluation of foot osteoarthritis: sensitivity of radiographic variables and relationship to symptoms. *Osteoarthritis Cartilage* 2009;17:298-303.
265. Drakonaki EE, Kho JS, Sharp RJ, et al. Efficacy of ultrasound-guided steroid injections for pain management of midfoot joint degenerative disease. *Skeletal Radiol* 2011;40:1001-6.
266. Han SH, Park DY, Kim TH. Prognostic factors after intra-articular hyaluronic acid injection in ankle osteoarthritis. *Yonsei Med J* 2014;55:1080-6.
267. Sun SF, Hsu CW, Sun HP, et al. The effect of three weekly intra-articular injections of hyaluronate on pain, function, and balance in patients with unilateral ankle arthritis. *J Bone Joint Surg Am.* 2011;93:1720-6.
268. Cicuttini FM, Baker J, Hart DJ, et al. Relation between Heberden's nodes and distal interphalangeal joint osteophytes and their role as markers of generalised disease. *Ann Rheum Dis* 1998;57:246-8.
269. Grainger AJ, Farrant JM, O'Connor PJ, Tan AL, Tanner S, Emery P, et al. MR imaging of erosions in interphalangeal joint osteoarthritis: is all osteoarthritis erosive? *Skeletal Radiol* 2007;36:737-45.
270. Haims AH MA, Schweitzer ME, Morrison WB, et al. MRI in the diagnosis of cartilage injury in the wrist. *AJR Am J Roentgenol* 2004;182:1267-70.
271. Haugen IK, Boyesen P, Slatkowsky-Christensen B, et al. Comparison of features by MRI and radiographs of the interphalangeal finger joints in patients with hand osteoarthritis. *Ann Rheum Dis* 2012;71:345-50.
272. Mutimer J, Green J, Field J. Comparison of MRI and wrist arthroscopy for assessment of wrist cartilage. *J Hand Surg Eur Vol* 2008;33:380-2.
273. Keen HI, Wakefield RJ, Grainger AJ, et al. Can ultrasonography improve on radiographic assessment in osteoarthritis of the hands? A comparison between radiographic and ultrasonographic detected pathology. *Ann Rheum Dis* 2008;67:1116-20.
274. Koutroumpas AC, Alexiou IS, Vlychou M, et al. Comparison between clinical and ultrasonographic assessment in patients with erosive osteoarthritis of the hands. *Clin Rheumatol* 2010;29:511-6.

275. Mathiessen A, Haugen IK, Slatkowsky-Christensen B, et al. Ultrasonographic assessment of osteophytes in 127 patients with hand osteoarthritis: exploring reliability and associations with MRI, radiographs and clinical joint findings. *Ann Rheum Dis* 2013;72:51-6.
276. Vlychou M, Koutroumpas A, Alexiou I, Fezoulidis I, Sakkas LI. High-resolution ultrasonography and 3.0 T magnetic resonance imaging in erosive and nodal hand osteoarthritis: high frequency of erosions in nodal osteoarthritis. *Clin Rheumatol* 2013;32:755-62.
277. Wittoek R, Carron P, Verbruggen G. Structural and inflammatory sonographic findings in erosive and non-erosive osteoarthritis of the interphalangeal finger joints. *Ann Rheum Dis* 2010;69:2173-6.
278. Iagnocco A, Filippucci E, Ossandon A, et al. High resolution ultrasonography in detection of bone erosions in patients with hand osteoarthritis. *J Rheumatol* 2005;32:2381-3.
279. Vlychou M, Koutroumpas A, Malizos K, et al. Ultrasonographic evidence of inflammation is frequent in hands of patients with erosive osteoarthritis. *Osteoarthritis Cartilage* 2009;17:1283-7.
280. Wittoek R, Jans L, Lambrecht V, et al. Reliability and construct validity of ultrasonography of soft tissue and destructive changes in erosive osteoarthritis of the interphalangeal finger joints: a comparison with MRI. *Ann Rheum Dis* 2011;70:278-83.
281. Iagnocco A, Coari G. Usefulness of high resolution US in the evaluation of effusion in osteoarthritis first carpometacarpal joint. *Scand J Rheumatol* 2000;29:170-3.
282. Jones G, Cooley HM, Bellamy N. A cross-sectional study of the association between Heberden's nodes, radiographic osteoarthritis of the hands, grip strength, disability and pain. *Osteoarthritis Cartilage* 2001;9:606-11.
283. Bijsterbosch J WI, Meulenbelt I, Rosendaal FR, et al. Clinical burden of erosive hand osteoarthritis and its relationship to nodes. *Ann Rheum Dis* 2010;69:1784-8.
284. Kortekaas MC KW, Reijnierse, Huizinga TWJ, et al. osteophytes and joint space narrowing are independently associated with pain in finger joints in hand osteoarthritis. *Ann Rheum Dis* 2011;70:1835-7.
285. Kwok WY, Kloppenburg M, Marshall M, et al. Comparison of clinical burden between patients with erosive hand osteoarthritis and inflammatory arthritis in symptomatic community-dwelling adults: The Keele clinical assessment studies. *Rheumatology (Oxford)* 2013;52:2260-7.
286. Marshall M, van der Windt D, Nicholls E, et al. Radiographic hand osteoarthritis: patterns and associations with hand pain and function in a community-dwelling sample. *Osteoarthritis Cartilage* 2009;17:1440-7.
287. Wittoek R, Cruyssen BV, Verbruggen G. Predictors of functional impairment and pain in erosive osteoarthritis of the interphalangeal joints: Comparison with controlled inflammatory arthritis. *Arthritis Rheum* 2012;64:1430-6.
288. Keen HI WR, Grainger AJ, Hensor EMA, et al. An ultrasonographic study of osteoarthritis of the hand: synovitis and its relationship to structural pathology and symptoms. *Arthritis Care Res* 2008;2008:1756-63.
289. Kortekaas MC, Kwok WY, Reijnierse M, et al. Pain in hand osteoarthritis is associated with inflammation: The value of ultrasound. *Ann Rheum Dis* 2010;69:1367-9.
290. Jonsson H, Eliasson GJ, Petursson E. Scintigraphic hand osteoarthritis (OA)--prevalence, joint distribution, and association with OA at other sites. *J rheumatol* 1999;26:1550-6.
291. Dahagin S B-ZS, Hazes JMW, Koes BW. Clinical burden of radiographic hand osteoarthritis: a systematic appraisal. *Arthritis Care Res* 2006;55:636-47.
292. Kwok WY, Plevier JW, Rosendaal FR, et al. Risk factors for progression in hand osteoarthritis: a systematic review. *Arthritis Care Res* 2013;65:552-62.

293. Haugen IK, Boyesen P, Slatkowsky-Christensen B, et al. Associations between MRI-defined synovitis, bone marrow lesions and structural features and measures of pain and physical function in hand osteoarthritis. *Ann Rheum Dis* 2012;71:899-904.
294. Wenham CY, Hensor EM, Grainger AJ, et al. A randomized, double-blind, placebo-controlled trial of low-dose oral prednisolone for treating painful hand osteoarthritis. *Rheumatology (Oxford)* 2012;51:2286-94. E
295. Haugen IK, Slatkowsky-Christensen B, Boyesen P, et al. Cross-sectional and longitudinal associations between radiographic features and measures of pain and physical function in hand osteoarthritis. *Osteoarthritis Cartilage*. 2013;21:1191-8.
296. Ding H, Solovieva S, Leino-Arjas P. Determinants of incident and persistent finger joint pain during a five-year followup among female dentists and teachers. *Arthritis Care Res* 2011;63:702-10.
297. Bijsterbosch J, Haugen IK, Malines C, et al. Reliability, sensitivity to change and feasibility of three radiographic scoring methods for hand osteoarthritis. *Ann Rheum Dis* 2011;70:1465-7.
298. Dahaghin S, Bierma-Zeinstra SMA, Reijman M, et al. Does hand osteoarthritis predict future hip or knee osteoarthritis? *Arthritis and rheumatism*. 2005;52:3520-7.
299. Filkova M, Senolt L, Braun M, et al. Serum hyaluronic acid as a potential marker with a predictive value for further radiographic progression of hand osteoarthritis. *Osteoarthritis Cartilage* 2009;17:1615-9.
300. Balblanc JC, Mathieu P, Mathieu L, et al. Progression of digital osteoarthritis: a sequential scintigraphic and radiographic study. *Osteoarthritis Cartilage* 1995;3:181-6.
301. Hutton CW, Higgs ER, Jackson PC, et al. 99mTc HMDP bone scanning in generalised nodal osteoarthritis. II. The four hour bone scan image predicts radiographic change. *Ann Rheum Dis* 1986;45:622-6.
302. McCarthy C, Cushnaghan J, Dieppe P. The predictive role of scintigraphy in radiographic osteoarthritis of the hand. *Osteoarthritis Cartilage* 1994;2:25-8.
303. Kortekaas MC, Kwok WY, Reijnierse M, et al. Follow-up study of inflammatory ultrasound features in hand osteoarthritis over a period of 3 months: variable as well as constant. *Osteoarthritis Cartilage* 2014;22:40-3.
304. Zhang W, Robertson J, Doherty S, et al. Index to ring finger length ratio and the risk of osteoarthritis. *Arthritis Rheum* 2008;58:137-44.
305. Mallinson PI, Tun JK, Farnell RD, et al. Osteoarthritis of the thumb carpometacarpal joint: correlation of ultrasound appearances to disability and treatment response. *Clin Radiol* 2013;68:461-5.
306. Botha-Scheepers S, Riyazi N, et al. Progression of hand osteoarthritis over 2 years: A clinical and radiological follow-up study. *Ann Rheum Dis* 2009;68:1260-4.
307. Botha-Scheepers S, Watt I, Breedveld FC, et al. Reading radiographs in pairs or in chronological order influences radiological progression in osteoarthritis. *Rheumatology (Oxford)* 2005;44:1452-5.
308. Maheu E, Cadet C, Gueneugues S, et al. Reproducibility and sensitivity to change of four scoring methods for the radiological assessment of osteoarthritis of the hand. *Ann Rheum Dis* 2007;66:464-9.
309. Macfarlane DG, Buckland-Wright JC, Lynch JA. Osteophytes in the osteoarthritic hand: their incidence, size, distribution, and progression. *Ann Rheum Dis* 1991;50:627-30.
310. Jans L, De Coninck T, Wittoek R, et al. 3 T DCE-MRI assessment of synovitis of the interphalangeal joints in patients with erosive osteoarthritis for treatment response monitoring. *Skeletal Radiol* 2013;42:255-60.

311. Keen HI, Wakefield RJ, Hensor EMA, et al. Response of symptoms and synovitis to intra-muscular methylprednisolone in osteoarthritis of the hand: An ultrasonographic study. *Rheumatology (Oxford)* 2010;49:1093-100.
312. Klauser AS FR, Kupferthaler K, Feuchtner G, et al. Sonographic criteria for therapy follow-up in the course of ultrasound-guided intra-articular injections of hyaluronic acid in hand osteoarthritis. *Eur J Radiol* 2012;81:1607-11.
313. Saltzherr MS, Selles RW, Bierma-Zeinstra SM, et al. Metric properties of advanced imaging methods in osteoarthritis of the hand: a systematic review. *Ann Rheum Dis* 2014;73:365-75.
314. Karalezli N, Ogun TC, Kartal S, et al. The pain associated with intraarticular hyaluronic acid injections for trapeziometacarpal osteoarthritis. *Clin Rheumatol* 2007;26:569-71.
315. Leydet-Quilici H, Le Corroller T, Bouvier C, et al. Advanced hip osteoarthritis: magnetic resonance imaging aspects and histopathology correlations. *Osteoarthritis Cartilage* 2010;18:1429-35.
316. Taljanovic MS, Graham AR, Benjamin JB, et al. Bone marrow edema pattern in advanced hip osteoarthritis: quantitative assessment with magnetic resonance imaging and correlation with clinical examination, radiographic findings, and histopathology. *Skeletal radiol* 2008;37:423-31..
317. Xu L, Hayashi D, Guermazi A, et al. The diagnostic performance of radiography for detection of osteoarthritis-associated features compared with MRI in hip joints with chronic pain. *Skeletal Radiol* 2013;42:1421-8.
318. Dolin SJ, De CWAC, Ashford N, et al. Factors affecting medical decision-making in patients with osteoarthritis of the hip: Allocation of surgical priority. *Disabil Rehabil* 2003;25:771-7.
319. Iagnocco A, Filippucci E, Riente L, et al. Ultrasound imaging for the rheumatologist XLI. Sonographic assessment of the hip in OA patients. *Clin Exp Rheumatol* 2012;30:652-7.
320. Recnik G, Vengust R, Kralj-Iglic V, et al. Association between sub-clinical acetabular dysplasia and a younger age at hip arthroplasty in idiopathic osteoarthritis. *J Int Med Res* 2009;37:1620-5.
321. Lieveense AM, Bierma-Zeinstra SMA, Verhagen AP, et al. Influence of obesity on the development of osteoarthritis of the hip: A systematic review. *Rheumatology (Oxford)* 2002;41:1155-62.
322. Wright AA CC, Abbott JH. Variables Associated With the Progression of Hip Osteoarthritis: A Systematic Review. *Arthritis Care Res* 2009;61:925-36.
323. Agricola R, Heijboer MP, Roze RH, et al. Pincer deformity does not lead to osteoarthritis of the hip whereas acetabular dysplasia does: Acetabular coverage and development of osteoarthritis in a nationwide prospective cohort study (CHECK). *Osteoarthritis Cartilage*. 2013;21:1514-21.
324. Agricola R, Reijman M, Bierma-Zeinstra SMA, et al. Total hip replacement but not clinical osteoarthritis can be predicted by the shape of the hip: A prospective cohort study (CHECK). *Osteoarthritis Cartilage*. 2013;21:559-64.
325. Arden NK, Lane NE, Parimi N, et al. Defining incident radiographic hip osteoarthritis for epidemiologic studies in women. *Arthritis Rheum* 2009;60:1052-9.
326. Gossec L, Jordan JM, Lam MA, et al. Comparative evaluation of three semi-quantitative radiographic grading techniques for hip osteoarthritis in terms of validity and reproducibility in 1404 radiographs: report of the OARSI-OMERACT Task Force. *Osteoarthritis Cartilage* 2009;17:182-7.
327. Gossec L, Tubach F, Baron G, et al. Predictive factors of total hip replacement due to primary osteoarthritis: A prospective 2 year study of 505 patients. *Ann Rheum Dis* 2005;64:1028-32.

328. Lieveense AM, Koes BW, Verhaar JAN, et al. Prognosis of hip pain in general practice: A prospective followup study. *Arthritis Care Res* 2007;57:1368-74.
329. Maillefert JF, Gueguen A, Nguyen M, et al. Relevant change in radiological progression in patients with hip osteoarthritis. I. Determination using predictive validity for total hip arthroplasty. *Rheumatology (Oxford)* 2002;41:142-7.
330. Lane NE NM, Hochberg MC, Hung Y, et al. Progression of Radiographic Hip Osteoarthritis Over Eight Years in a Community Sample of Elderly White Women. *Arthritis and Rheumatology (Oxford)* 2004;50:1477-86.
331. Ledingham J DS, Preston B, Milligan G, et al. Radiographic progression of hospital referred osteoarthritis of the hip. *Ann Rheum Dis* 1993;263-7.
332. Conrozier T, Jousseau CA, Mathieu P, et al. Quantitative measurement of joint space narrowing progression in hip osteoarthritis: A longitudinal retrospective study of patients treated by total hip arthroplasty. *Br J Rheumatol* 1998;37:961-8.
333. Reijman M, Hazes JMW, Pols H, et al. Role of radiography in predicting progression of osteoarthritis of the hip: Prospective cohort study. *BMJ* 2005;330:1183-5.
334. Barr RJ, Gregory JS, Reid DM, et al. Predicting OA progression to total hip replacement: Can we do better than risk factors alone using active shape modelling as an imaging biomarker? *Rheumatology (Oxford)* 2012;51:562-70.
335. Agricola R, Heijboer MP, Bierma-Zeinstra SMA, et al. Cam impingement causes osteoarthritis of the hip: A nationwide prospective cohort study (CHECK). *Ann Rheum Dis* 2013;72:918-23.
336. Dougados M GA, Nguyen M, Berdah L, et al. Radiological progression of hip osteoarthritis: definition, risk factors and correlations with clinical status. *Ann Rheum Dis* 1996;55:356-62.
337. Ahedi H, Aitken D, Blizzard L, et al. A population-based study of the association between hip bone marrow lesions, high cartilage signal, and hip and knee pain. *Clin Rheumatol* 2014;33:369-76.
338. Atchia I, Kane D, Reed MR, et al. Efficacy of a single ultrasound-guided injection for the treatment of hip osteoarthritis. *Annals Rheum Dis* 2011;70:110-6.
339. Lequesne M, Maheu E, Cadet C, et al. Structural effect of avocado/soybean unsaponifiables on joint space loss in osteoarthritis of the hip. *Arthritis Care Res* 2002;47:50-8.
340. Rennesson-Rey B, Rat AC, Chary-Valckenaere I, et al. Does joint effusion influence the clinical response to a single Hylan GF-20 injection for hip osteoarthritis? *Joint Bone Spine* 2008;75:182-8.
341. Rozendaal RM, Uitterlinden EJ, van Osch GJVM, et al. Effect of glucosamine sulphate on joint space narrowing, pain and function in patients with hip osteoarthritis; subgroup analyses of a randomized controlled trial. *Osteoarthritis Cartilage* 2009;17:427-32.
342. Hoeksma HL DJ, ROnday HK, Breedveld FC, et al. Manual therapy in osteoarthritis of the hip: outcome in subgroups of patients. *Rheumatology (Oxford)* 2005;44:461-4.
343. Deshmukh AJ, Panagopoulos G, Alizadeh A, et al. Intra-articular hip injection: does pain relief correlate with radiographic severity of osteoarthritis? *Skeletal Radiol* 2011;40:1449-54.
344. Robinson P, Keenan AM, Conaghan PG. Clinical effectiveness and dose response of image-guided intra-articular corticosteroid injection for hip osteoarthritis. *Rheumatology (Oxford)* 2007;46:285-91.
345. Auleley GR, Giraudeau B, Dougados M, et al. Radiographic assessment of hip osteoarthritis progression: Impact of reading procedures for longitudinal studies. *Ann Rheum Dis* 2000;59:422-7.

346. Botha-Scheepers S, Watt I, Rosendaal FR, et al. Changes in outcome measures for impairment, activity limitation, and participation restriction over two years in osteoarthritis of the lower extremities. *Arthritis Care Res* 2008;59:1750-5.
347. Conrozier T, Brandt K, Piperno M, et al. Reproducibility and sensitivity to change of a new method of computer measurement of joint space width in hip osteoarthritis. Performance of three radiographic views obtained at a 3-year interval. *Osteoarthritis Cartilage* 2009;17(7):864-70.
348. Conrozier T, Saxne T, Fan CSS, et al. Serum concentrations of cartilage oligomeric matrix protein and bone sialoprotein in hip osteoarthritis: A one year prospective study. *Ann Rheum Dis* 1998;57:527-32.
349. Dougados M, Nguyen M, Berdah L, et al. Evaluation of the structure-modifying effects of diacerein in hip osteoarthritis: ECHODIAH, a three-year, placebo-controlled trial. *Arthritis Rheum* 2001;44:2539-47.
350. Maheu E, Cadet C, Marty M, et al. Reproducibility and sensitivity to change of various methods to measure joint space width in osteoarthritis of the hip: a double reading of three different radiographic views taken with a three-year interval. *Arthritis Res Ther* 2005;7:R1375-85.
351. Maillefert JF, Sharp JT, Aho LS, et al. Comparison of a computer based method and the classical manual method for radiographic joint space width assessment in hip osteoarthritis. *Journal Rheumatol* 2002;29:2592-6.
352. Papaloucas CD, Ward RJ, Tonkin CJ, et al. Cancellous bone changes in hip osteoarthritis: a short-term longitudinal study using fractal signature analysis. *Osteoarthritis Cartilage* 2005;13:998-1003.
353. Pavelka K, Gatterova J, Gollerova V, et al. A 5-year randomized controlled, double-blind study of glycosaminoglycan polysulphuric acid complex (Rumalon(registered trademark)) as a structure modifying therapy in osteoarthritis of the hip and knee. *Osteoarthritis Cartilage*. 2000;8:335-42.
354. Ratzlaff C, Van Wyngaarden C, Duryea J. Location-specific hip joint space width for progression of hip osteoarthritis - Data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2014;22:1481-7.
355. Jaremko JL, Lambert RG, Zubler V, et al. Methodologies for semiquantitative evaluation of hip osteoarthritis by magnetic resonance imaging: approaches based on the whole organ and focused on active lesions. *J Rheumatol* 2014;41:359-69.