

Difference between SLE classification and diagnosis and importance of attribution. Response to: 'Do the 2019 EULAR/ACR SLE classification criteria close the door on certain groups of SLE patients?' by Chi *et al*

In their letter, Dr Chi and colleagues¹ express two concerns regarding the new European League Against Rheumatism/American College of Rheumatology (EULAR/ACR) classification criteria. On the one hand, they correctly remark that patients with uncommon, isolated organ manifestations may be more difficult to classify with any set of classification criteria. On the other hand, they caution against mistaking malignancies for systemic lupus erythematosus (SLE).

The first sensitivity-related issue is certainly correct. However, it is important to reiterate that the new EULAR/ACR criteria are classification criteria. Our manuscript explicitly states that they should not be used as diagnostic criteria and that diagnosis remains with the appropriately trained physician.^{2,3} In order for a criteria set to remain feasible, that is, not to contain hundreds of items, uncommon isolated manifestations or laboratory tests cannot be accounted for. We had a focus on early symptoms,⁴ where only fever qualified as sufficiently performing for classification in early disease to be added to the criteria list. Still, it is important to be aware of non-criteria organ manifestations to fully understand the complexity of SLE, and we thank our colleagues for bringing this up. As SLE is a highly heterogeneous disease, there are probably hundreds of rare SLE manifestations that are not included in the new criteria set, yet, the new criteria still have excellent sensitivity when compared with prior criteria sets.

For the second specificity-related issue, it is also important to distinguish classification from diagnosis. In addition, this gives us the opportunity to reiterate two important features of the new criteria^{1,2} that should limit the risk of classifying patients with malignancies as patients with SLE. One, most of the malignancies that are mimicking SLE features are haematological. In this regard, going back to domains and counting the highest ranking item in each domain only is an advantage. This step, which followed an in-depth evaluation of associations between SLE items,⁵ will prevent counting several changes in blood counts.

Even more important is the general attribution rule of the new criteria: items are only to be counted for SLE, if there is no more likely alternative explanation.^{2,3} This approach requires both clinical understanding and diligence but should also safeguard against misclassifying malignancies.

Martin Aringer ¹, Karen H Costenbader,² Thomas Dörner ³, Sindhu R Johnson⁴

¹Division of Rheumatology, Department of Medicine III, University Medical Center and Faculty of Medicine, TU Dresden, Dresden, Germany

²Division of Rheumatology, Immunology and Allergy, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

³Department of Rheumatology and Clinical Immunology, Charité – Universitätsmedizin Berlin, Corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department of Rheumatology and Clinical Immunology, Berlin, Germany

⁴Division of Rheumatology, Department of Medicine, Toronto Western Hospital, Mount Sinai Hospital; Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada

Correspondence to Professor Martin Aringer, Internal Medicine III, Medical Faculty, Technical University of Dresden, Dresden 01069, Germany; martin.aringer@uniklinikum-dresden.de

Contributors All four authors drafted this response together and approved of the final version.

Funding The SLE classification criteria project was equally supported by EULAR and the ACR.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Aringer M, Costenbader KH, Dörner T, *et al*. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2019-216338

Received 18 September 2019

Accepted 21 September 2019



► <http://dx.doi.org/10.1136/annrheumdis-2019-216253>

Ann Rheum Dis 2019;**0**:1. doi:10.1136/annrheumdis-2019-216338

ORCID iDs

Martin Aringer <http://orcid.org/0000-0003-4471-8375>

Thomas Dörner <http://orcid.org/0000-0002-6478-7725>

REFERENCES

- Chi H, Teng J, Wang Z, *et al*. Do the 2019 EULAR/ACR SLE classification criteria close the door on certain groups of SLE patients? *Ann Rheum Dis* 2019;annrheumdis-2019-216253.
- Aringer M, Costenbader K, Daikh D, *et al*. 2019 European League against Rheumatism/American College of rheumatology classification criteria for systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:1151–9.
- Aringer M, Costenbader K, Daikh D, *et al*. 2019 European League against Rheumatism/American College of rheumatology classification criteria for systemic lupus erythematosus. *Arthritis Rheumatol* 2019;71:1400–12.
- Tedeschi SK, Johnson SR, Boumpas D, *et al*. Developing and refining new candidate criteria for systemic lupus erythematosus classification: an international collaboration. *Arthritis Care Res* 2018;70:571–81.
- Touma Z, Cervera R, Brinks R, *et al*. Associations among classification criteria items within systemic lupus erythematosus. *Arthritis Care Res* 2019.