Gut-derived CD8⁺ tissue-resident memory T cells are expanded in the peripheral blood and synovia of SpA patients

We read with interest the recently published paper from Qiayum *et al*¹ demonstrating a novel integrin-expressing mature Crohn's disease (CD)8⁺ T cell population defined as CD49a⁺CD103⁺β7⁺CD29⁺ cells in the synovial fluids of ankylosing spondylitis (AS) patients. Although the authors did not analyse gut samples from AS patients, they speculate that these cells might be gut-derived cells. Interestingly, as stated by authors, the transcriptional and phenotypic signature of these cells is reminiscent of human tissue-resident memory T cells (T_{RM}). T_{RM} are a subset of cells important as the first line of defence from infection in mucosal tissues, never studied in spondyloarthritis (SpA).² For clarifying whether these cells could be of intestinal original, we set up additional analyses, wondering if we could see similar results in paired samples of patients with SpA.

Paired gut and synovial samples and fluids and peripheral blood samples (PCMCs) were obtained from patients with SpA (HLA-B27 positive; n=6), never treated with biologic agents at the time of sample collection. Gut samples were also obtained from healthy controls (HCs) (HLA-B27 negative; n=6) and synovial tissues from osteoarthritis (OA) patients (HLA-B27 negative; n=6). Peripheral blood mononuclear cells (PBMCs) were also obtained from HCs. CD103 and CD8 expression were assessed by immunohistochemistry. The percentage of T_{RM} T cells (defined as CD8⁺CD69⁺CD103⁺ cells) among isolated lamina propria mononuclear cells (LPMCs) and PBMCs from SpA patients and controls were also analysed by flow cytometry.

Part of the results is shown in figure 1. In the gut tissues, the number of CD8⁺CD103⁺ cells was consistently increased in the inflammatory SpA samples compared with non-inflammatory samples (figure 1A–E). Tissue distribution confirmed their predominant localisation in the context of epithelial layer (figure 1A–D). Flow cytometric analysis of LPMCs confirmed the expansion of CD8⁺CD69⁺CD103⁺ T_{RM} T cells,

Figure 1 T_{RM}s in the gut, peripheral blood and synovia of SpA patients. (A–D) Representative imaging showing CD103 (A and C) and CD8 (B and D) expression in sequential gut sections of controls (A–B) and SpA patients (C–D). (E) Higher numbers of CD8/CD103 positive cells were observed in SpA patients compared with controls. (F) Representative dot plots showing gating strategy for T_{RM}s in the peripheral blood of SpA patients. (G–I) Percentages of T_{RM}s among LPMC (G), PBMC (H) and SFMC (I) in SpA patients and controls. (J–K) Representative imaging showing CD103 (J) and CD8 (K) expression in sequential synovial sections of SpA patients. (L) Higher numbers of CD8/CD103 positive cells were observed in SpA patients compared with controls. (A–D) and (J–K): Original magnification ×250. LPMC, lamina propria mononuclear cell; PBMC, peripheral blood mononuclear cell; T_{RM}, tissue-resident memory T cells.

Correspondence

mainly producing IFN γ , in SpA patients compared with HCs (figure 1F–G). The expansion of CD8⁺CD69⁺CD103⁺ T_{RM} T cells, mainly producing IFN γ , was also confirmed in SpA PBMCs (figure 1H) and synovial mononuclear cells (SFMC) (figure 1I), compared with HCs. The majority of circulating and synovial fluids CD8⁺CD69⁺CD103⁺ T_{RM} expressed the intestinal homing receptor α 4 β 7 (67% and 75%, respectively) suggesting their gut origin (data not shown). Finally, immunohistochemical analysis of sequential synovial samples confirmed tissue infiltration of CD8⁺CD103⁺ cells in the inflamed synovial tissues of SpA patients (figure 1J–L).

The existence of a gut–joint has been hypothesised in SpA patients.³ The inflamed gut could actively participate in the pathogenesis of SpA through the production of proinflammatory cytokines, such as IL-23p19⁴ and IL-9,⁵ and the differentiation of potentially pathogenic innate cells producing IL-22 and IL-17.³ T_{RM} are a critical component of mucosal immune defence by acting as peripheral sentinels capable of rapidly mobilising protective tissue immunity on pathogen recognition.² Our data confirm the expansion of T_{RM} in the synovial compartment of SpA patients, providing evidence of T_{RM} expansion in the peripheral blood and the gut. The expression of α 4 β 7 by circulating T_{RM} in SpA might support the re-circulation of these cells from the gut to the peripheral blood and inflamed joints.

Giuliana Guggino,¹ Aroldo Rizzo,² Daniele Mauro,³ Federica Macaluso,¹ Francesco Ciccia³

¹Department of Health Promotion, Mother and Child Care, Internal Medicine and Medical Specialties, Università degli Studi di Palermo, Palermo, Italy

²Dipartimento di Oncologia, Azienda Ospedaliera Villa Sofia-Cervello, Palermo, Italy

³Dipartimento di Medicina di Precisione, Università degli Studi della Campania Luigi Vanvitelli, Napoli, Italy

Correspondence to Dr Francesco Ciccia, Dipartimento di Medicina di Precisione, Università degli Studi della Campania Luigi Vanvitelli, Napoli 80131, Italy; francescociccia@tiscali.it

Handling editor Josef S Smolen

Contributors All the authors gave substantial contributions to the conception or design of the work, the acquisition, analysis or interpretation of data, drafting the work or revising it critically for important intellectual content, or final approval of the version published. All the authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding This study was in part supported by a grant of Ministero dell'Istruzione, dell'Università e della Ricerca Scientifica from Italy.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Guggino G, Rizzo A, Mauro D, et al. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2019-216456

Received 10 October 2019

Accepted 13 October 2019

Ann Rheum Dis 2019;0:1–2. doi:10.1136/annrheumdis-2019-216456

ORCID iD

Francesco Ciccia <http://orcid.org/0000-0002-9352-1264>

REFERENCES

- 1 Qaiyum Z, Gracey E, Yao Y, et al. Integrin and transcriptomic profiles identify a distinctive synovial CD8⁺ T cell subpopulation in spondyloarthritis. *Ann Rheum Dis* 2019. doi:10.1136/annrheumdis-2019-215349. [Epub ahead of print: 30 Aug 2019].
- 2 Schenkel JM, Masopust D. Tissue-Resident memory T cells. *Immunity* 2014;41:886–97.
- 3 Rizzo A, Ferrante A, Guggino G, et al. Gut inflammation in spondyloarthritis. *Best Pract Res Clin Rheumatol* 2017;31:863–76.
- 4 Ciccia F, Guggino G, Zeng M, et al. Proinflammatory CX3CR1+CD59+Tumor necrosis factor-like molecule 1A+Interleukin-23+ monocytes are expanded in patients with ankylosing spondylitis and modulate innate lymphoid cell 3 immune functions. *Arthritis Rheumatol* 2018;70:2003–13.
- 5 Ciccia F, Guggino G, Ferrante A, et al. Interleukin-9 overexpression and Th9 polarization characterize the inflamed gut, the synovial tissue, and the peripheral blood of patients with psoriatic arthritis. *Arthritis Rheumatol* 2016;68:1922–31.