Frequency of MRI changes suggestive of axial spondyloarthritis in the axial skeleton in a large population-based cohort of individuals aged <45 years

I read with great interest the recent report by Baraliakos *et al* published in the *Annals of Rheumatic Disease*, about a general population cohort study of healthy volunteers that examined the presence of bone marrow oedema and fatty lesions on MRI of the spine and sacroiliac joints.¹ Over the past decade, imaging techniques such as the MRI have revolutionised the application of radiographic findings in the early diagnosis of axial spondyloarthritis (axSpA).²

In this study, the authors elegantly identified an increased frequency of fatty lesions as well as bone marrow oedema in the vertebral corners of the spine, particularly in the lower part of the thoracic spine. I agree with the authors' comment that the vertebral MRI lesions found in the healthy volunteers could be induced by mechanical load or early osteoarthritis; however, given the distribution and the characteristics of the lesions in some patients, these findings could be reflective of diffuse idiopathic skeletal hyperostosis (DISH).

DISH is a common skeletal condition of unknown aetiology characterised by calcification of spinal ligaments and entheses, and radiographic evidence of flowing ossification along with the vertebral bodies, most commonly in the thoracic spine.³ From an epidemiologic perspective, DISH has an incidence rate of 4%–7% and could be associated with symptoms of inflammatory back pain and stiffness among patients, mimicking axSpA.^{4,5}

In the current study, the authors pointed out that older patients had increased frequency of the thoracic lesions on MRI, a characteristic radiographic feature of DISH, although the prevalence of the disease increases with age, DISH may affect patients under the age 45.^{6,7} Moreover, a recent study that evaluated the spine of 53 DISH patients with the use of MRI, demonstrated bone marrow oedema and fat deposition in at least one vertebral corner, in 76% and 67% of patients, respectively, similar results seen in axSpA.⁸

In conclusion, we believe caution should be taken regarding the interpretation of the MRI of the spine findings in order to avoid an erroneous diagnosis of axSpA, and consequently, the use of unnecessary, expensive and potentially harmful treatments. Future studies are warranted to elucidate the long-term progression of the inflammatory spinal lesions in healthy adults and to provide us with a better understanding of the natural history of conditions such as DISH.

Konstantinos Parperis ^{1,2}

¹Internal Medicine, Division of Rheumatology, University of Cyprus Medical School, Nicosia, Cyprus

²Department of Medicine, University of Arizona College of Medicine Phoenix, Phoenix, Arizona, USA

Correspondence to Dr Konstantinos Parperis, Internal Medicine, Division of Rheumatology, University of Cyprus Medical School, Nicosia 1678, Cyprus; kparpe02@ucy.ac.cy

Contributors I confirm that I have contributed to the planning, conduct, and reporting of the work described in the article

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer I confirm the manuscript has not been submitted or is not simultaneously being submitted elsewhere, and that no portion of the data has been or will be published in proceedings or transactions of meetings or symposium volumes.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Parperis K. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2019-216773

Received 5 December 2019

Accepted 8 December 2019

Ann Rheum Dis 2019;0:1. doi:10.1136/annrheumdis-2019-216773

ORCID iD

Konstantinos Parperis <http://orcid.org/0000-0001-6009-0130>

REFERENCES

- Baraliakos X, Richter A, Feldmann D, *et al*. Frequency of MRI changes suggestive of axial spondyloarthritis in the axial skeleton in a large population-based cohort of individuals aged <45 years. *Ann Rheum Dis* 2019;annrheumdis-2019-215553.
- Mandl P, Navarro-Compán V, Terslev L, *et al*. EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice. *Ann Rheum Dis* 2015;74:1327–39.
- Belanger TA, Rowe DE. Diffuse idiopathic skeletal hyperostosis: musculoskeletal manifestations. *J Am Acad Orthop Surg* 2001;9:258–67.
- Paley D, Schwartz M, Cooper P, *et al*. Fractures of the spine in diffuse idiopathic skeletal hyperostosis. *Clin Orthop Relat Res* 1991;22–32.
- Mata S, Fortin PR, Fitzcharles MA, *et al*. A controlled study of diffuse idiopathic skeletal hyperostosis. clinical features and functional status. *Medicine* 1997;76:104–17.
- Mader R, Verlaan J-J, Buskila D. Diffuse idiopathic skeletal hyperostosis: clinical features and pathogenic mechanisms. *Nat Rev Rheumatol* 2013;9:741–50.
- Sieper J, Rudwaleit M, Baraliakos X, *et al*. The assessment of spondyloarthritis International Society (ASAS) Handbook: a guide to assess spondyloarthritis. *Ann Rheum Dis* 2009;68:ii1–44.
- Latourte A, Charlon S, Etcheto A, *et al*. Imaging findings suggestive of axial spondyloarthritis in diffuse idiopathic skeletal hyperostosis. *Arthritis Care Res* 2018;70:145–52.