Relationship between rheumatoid arthritis and interstitial lung disease is a 'two-way street'. Response to: 'Autoantibodies and interstitial lung disease in rheumatoid arthritis: towards a 'mix-and-match' approach' by Alunno *et al*

We welcome the comments by Alunno *et al* on our article about the association between anticarbamylated protein antibody (anti-CarP) specificities and rheumatoid arthritis-associated interstitial lung disease (RA-ILD).¹ The authors proposed a 'mix and match approach' consisting of the assessment of various antibody specificities at RA diagnosis with the aim of predicting the development of ILD.² We believe this hypothesis is reasonable, considering that different antimodified protein antibodies (AMPA) including anti-CarP, anticitrullinated and antiacetylated protein antibodies (ACPA and anti-AceP, respectively) have been associated with RA-ILD.^{1 3 4} Furthermore, a greater number of coexisting specificities of a single AMPA have been found in patients with RA-ILD.^{3 5}

The prevalence of ILD and its risk factors fluctuate between RA cohorts, partially due to differences in the screening methods, the population examined, and the defining criteria used. Although these factors were controlled in our study, there were differences in some baseline features between the main population and the replication sample, as pointed out by Alunno *et al*.² This may be due to the small sample size of the replication sample or because ILD screening and diagnosis was ultimately based on physician's criteria based on the ILD committee dictates from two hospitals. However, our final model was fitted after adjusting for these features and so they did not affect the results. Recently, Zhu *et al* reported a higher proportion of anti-CarP in Chinese patients diagnosed with RA-ILD compared to RA controls without ILD (53% vs 16%).⁶ Their findings are consistent with and enhance the external validity of our observations. However, larger multi-ethnic studies are still required.

We consider the association between ILD and RA is a 'two-way street'. It should be considered that: (1) ILD may be present before or around RA onset in one in three patients⁷ and (2) ACPA have been found in more than 20% of patients with idiopathic pulmonary fibrosis (IPF),⁸ of whom approximately one-third subsequently may develop RA.⁹ Thus, the 'mix and match approach' should be considered in RA and IPF. We believed a broader view (eg, multidisciplinary ILD committees) on the issues implied in the relation between ILD and RA should be considered in the design of future prospective collaborative studies.

Raul Castellanos-Moreira ¹, Sebastian Cruz Rodríguez-García ¹, Isabel Haro,² Raimon Sanmarti ¹

¹Rheumatology Department, Hospital Clínic de Barcelona, Barcelona, Spain

²Unit of Synthesis and Biomedical Applications of Peptides, Institute of Advanced Chemistry of Catalonia (IQAC-CSIC), Consejo Superior de Investigaciones Científicas, Barcelona, Spain

Correspondence to Dr Raimon Sanmarti, Rheumatology Department, Hospital Clínic de Barcelona, Barcelona 08036, Spain; sanmarti@clinic.cat

Handling editor Josef S Smolen

Twitter Raul Castellanos-Moreira @raul_cast_morei and Sebastian Cruz Rodríguez-García @sdclrodriguez

Contributors All authors have collaborated, read and approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Castellanos-Moreira R, Rodríguez-García SC, Haro I, *et al*. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2020-217432

Received 6 May 2020

Revised 7 May 2020

Accepted 12 May 2020

► <https://doi.org/10.1136/annrheumdis-2020-217410>

Ann Rheum Dis 2020;**0**:1. doi:10.1136/annrheumdis-2020-217432

ORCID iDs

Raul Castellanos-Moreira <http://orcid.org/0000-0002-4104-4101>

Sebastian Cruz Rodríguez-García <http://orcid.org/0000-0002-7773-151X>

Raimon Sanmarti <http://orcid.org/0000-0002-8864-3806>

REFERENCES

- Castellanos-Moreira R, Rodríguez-García SC, Gomara MJ, *et al*. Anti-carbamylated proteins antibody repertoire in rheumatoid arthritis: evidence of a new autoantibody linked to interstitial lung disease. *Ann Rheum Dis* 2020;79:587–94.
- Alunno A, Bistoni O, Bonifacio AF, *et al*. Autoantibodies and interstitial lung disease in rheumatoid arthritis: towards a 'mix-and-match' approach? *Ann Rheum Dis* 2020;annrheumdis-2020-217410.
- Giles JT, Danoff SK, Sokolove J, *et al*. Association of fine specificity and repertoire expansion of anticitrullinated peptide antibodies with rheumatoid arthritis associated interstitial lung disease. *Ann Rheum Dis* 2014;73:1487–94.
- England BR, Duryee MJ, Roul P, *et al*. Malondialdehyde-Acetaldehyde adducts and antibody responses in rheumatoid arthritis-associated interstitial lung disease. *Arthritis Rheumatol* 2019;71:1483–93.
- Alunno A, Bistoni O, Pratesi F, *et al*. Anti-citrullinated alpha enolase antibodies, interstitial lung disease and bone erosion in rheumatoid arthritis. *Rheumatology* 2018;57:850–5.
- Zhu H, Zhao LJ, Zhou Y, *et al*. [Significance of anti-carbamylated protein antibodies in patients with rheumatoid arthritis-associated interstitial lung disease]. *Beijing Da Xue Xue Bao Yi Xue Ban* 2019;51:1003–7.
- Hyldegard C, Hilberg O, Pedersen AB, *et al*. A population-based cohort study of rheumatoid arthritis-associated interstitial lung disease: comorbidity and mortality. *Ann Rheum Dis* 2017;76:1700–6.
- Solomon JJ, Matson S, Kelmenson LB, *et al*. IgA antibodies directed against citrullinated protein antigens are elevated in patients with idiopathic pulmonary fibrosis. *Chest* 2019;S0012-3692(19)34454-X.
- Katsumata M, Hozumi H, Yasui H, *et al*. Frequency and clinical relevance of anti-cyclic citrullinated peptide antibody in idiopathic interstitial pneumonias. *Respir Med* 2019;154:102–8.