Response to: 'Comment on: 'EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update' by Gossec *et al*' by Wei *et al*

We thank Wei *et al*¹ for their correspondence on the recently published European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis (PsA).² The authors of this correspondence raise some interesting points.

The first point relates to the role of cyclosporine A (CsA) in the treatment of patients with psoriatic disease. The authors mention that this drug is often used in Asia, in particular in skin psoriasis. They suggest that CsA could be proposed for PsA. While CsA is recommended for the management of skin psoriasis,³ EULAR's systematic literature searches did not reveal a convincing efficacy in PsA based on two small randomised controlled trials^{4,5}; indeed, even a randomised trial on the addition of CsA in patients who had an insufficient response to methotrexate failed to meet its primary endpoint.⁶ Therefore, CsA is not put forward in the EULAR recommendations for the management of PsA,² and these recommendations do not deal with patients who only have psoriasis.

The second point of the authors' comments relates to the definition of oligoarticular disease and, more specifically, the necessity for a full 66/68 joint count. We fully support using the 66/68 joint counts in PsA, as studies have shown that more limited joint counts may miss significant proportions of inflammation.⁷ The authors mentioned the Disease Activity in Psoriatic Arthritis (DAPSA) score, which is also recommended in the updated EULAR recommendations and, indeed, uses the 66/68 counts.⁸

Wei *et al* support the new recommendation for tapering of biological DMARDs in patients reaching sustained remission, but suggest that an algorithm or a tapering guidance would be useful.^{1,2} Currently, given the lack of data on this subject, it is difficult to propose formal tapering strategies.⁴ In the absence of any data for PsA, Wei *et al* may be referred to the EULAR RA management recommendations⁹ for a potential orientation, given that in RA tapering schedules have been studied.¹⁰ Tapering should be further investigated in PsA. The last point regards elderly patients and we agree with the authors that research is warranted on the management of elderly patients with PsA.

Laure Gossec ^{1,2}, Xenofon Baraliakos,³ Iain McInnes,⁴ Andreas Kerschbaumer ⁵, Maarten de Wit ⁶, Maxime Dougados,⁷ Jette Primdahl ^{8,9}, Désirée van der Heijde ¹⁰, Josef S Smolen¹¹

¹Institut Pierre Louis d'Epidémiologie et de Santé Publique, INSERM, Sorbonne Université, Paris, France

²APHU, Rheumatology Department, Hôpital Universitaire Pitie Salpêtrière, Paris, France

³Rheumazentrum Ruhrgebiet, Ruhr-Universität Bochum, Herne, Germany

⁴Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK

⁵Division of Rheumatology, Department of Medicine 3, and 2nd Department of Medicine, Medical University of Vienna, Hietzing Hospital, Vienna, Austria

⁶EULAR past Vice President representing People with Arthritis/Rheumatism in Europe (PARE), EULAR, Amsterdam, The Netherlands

⁷Medicine Faculty, Paris Descartes University, Paris, France

⁸Danish Hospital for Rheumatic Diseases, University Hospital of Southern Denmark, Sønderborg, Denmark

⁹Department of Regional Health Research, University of Southern Denmark, Odense, Denmark

¹⁰Rheumatology, Leiden University Medical Center, Leiden, The Netherlands

¹¹Medical University of Vienna, Vienna, Austria

Correspondence to Professor Laure Gossec, Institut Pierre Louis d'Epidémiologie et de Santé Publique, INSERM, Sorbonne Université, Paris 75005, France; laure.gossec@gmail.com

Handling editor David S Pisetsky

Contributors LG, XB and JSS drafted the text. The other authors reviewed the text and commented. All authors approved the final version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement A patient was involved in this response, as part of the steering committee of the recommendations

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Gossec L, Baraliakos X, McInnes I, *et al*. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2020-218456

Received 14 August 2020

Accepted 15 August 2020

► <http://dx.doi.org/10.1136/annrheumdis-2020-218385>

Ann Rheum Dis 2020;**0**:1. doi:10.1136/annrheumdis-2020-218456

ORCID iDs

Laure Gossec <http://orcid.org/0000-0002-4528-310X>

Andreas Kerschbaumer <http://orcid.org/0000-0002-6685-8873>

Maarten de Wit <http://orcid.org/0000-0002-8428-6354>

Jette Primdahl <http://orcid.org/0000-0002-1049-4150>

Désirée van der Heijde <http://orcid.org/0000-0002-5781-158X>

REFERENCES

- Wei CY *et al*. Comment on: 'EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update' by Gossec, *et al*. *Ann Rheum Dis*. doi: 10.1136/annrheumdis-2020-218385.
- Gossec L, Baraliakos X, Kerschbaumer A, *et al*. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis* 2020;79:700–12.
- Stiff KM, Glines KR, Porter CL, *et al*. Current pharmacological treatment guidelines for psoriasis and psoriatic arthritis. *Expert Rev Clin Pharmacol* 2018;11:1209–18.
- Kerschbaumer A, Smolen JS, Dougados M, *et al*. Pharmacological treatment of psoriatic arthritis: a systematic literature research for the 2019 update of the EULAR recommendations for the management of psoriatic arthritis. *Ann Rheum Dis* 2020;79:annrheumdis-2020-217163–786.
- Ramiro S, Smolen JS, Landewé R, *et al*. Pharmacological treatment of psoriatic arthritis: a systematic literature review for the 2015 update of the EULAR recommendations for the management of psoriatic arthritis. *Ann Rheum Dis* 2016;75:490–8.
- Fraser AD, van Kuijk AWR, Westhovens R, *et al*. A randomised, double blind, placebo controlled, multicentre trial of combination therapy with methotrexate plus ciclosporin in patients with active psoriatic arthritis. *Ann Rheum Dis* 2005;64:859–64.
- Duarte-García A, Leung YY, Coates LC, *et al*. Endorsement of the 66/68 joint count for the measurement of musculoskeletal disease activity: OMERACT 2018 psoriatic arthritis workshop report. *J Rheumatol* 2019;46:996–1005.
- Schoels MM, Aletaha D, Alasti F, *et al*. Disease activity in psoriatic arthritis (PSA): defining remission and treatment success using the DAPSA score. *Ann Rheum Dis* 2016;75:811–8.
- Smolen JS, Landewé RBM, Bijlsma JWI, *et al*. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis* 2020;79:annrheumdis-2019-216655–699.
- Burmester GR, Buttgerit F, Bernasconi C, *et al*. Continuing versus tapering glucocorticoids after achievement of low disease activity or remission in rheumatoid arthritis (SEMIRA): a double-blind, multicentre, randomised controlled trial. *Lancet* 2020;396:267–76.