

Online supplementary table 1. Clinical symptoms of patients with polyarteritis nodosa treated with tocilizumab.

	Symptoms
1	Fever, painful erythematous nodules on her extremities, bilateral arthritis in her ankles and knees, digital ischemia, myalgia
2	Fever, subcutaneous nodules, livedo reticularis, myalgias (myositis), weight loss, sensorimotor polyneuropathy
3	Fever, myalgias (fasciitis), weight loss, arthritis, subcutaneous nodules, drop hand, abdominal and flank pain, hypertension, increase of creatinine
4	Fever, myalgias, fatigue, weight loss
5	Myalgias (myositis), arthritis, livedo reticularis, sensorimotor polyneuropathy
6	Fever, fatigue, myalgias, weight loss, loss of appetite, polyneuropathy, arthralgia, pain in upper and lower extremities, abdominal pain, headache, double vision, tinnitus, dizziness
7	Necrotic purpura, constitutional symptoms, myalgias, weight loss, kidney artery aneurysms
8	Necrotic purpura, arthritis, livedo reticularis, occlusion and aneurysm of arm arteries, neuropathy
9	Multiple arthritis, myalgia, tenosynovitis, subcutaneous nodules, paraesthesia in the dominant hand, weight loss
10	Crural skin ulcers, subcutaneous nodules, multiple intestinal perforations, arterial hypertension, myalgias, weight loss, testicular pain, adhesive ileus
11	Fever, periostitis, epididymoorchitis, arthritis, myositis, stiff neck, right scrotum swelling and tenderness, vertebral arteritis