Response to: 'Correspondence on 'Increased risk of systemic lupus erythematosus in patients with autoimmune haemolytic anaemia: a nationwide population-based cohort study' by Goulielmos and Zervou

We thank Goulielmos and Zervou¹ for their interest on our article entitled 'Increased risk of systemic lupus erythematosus in patients with autoimmune haemolytic anaemia: a nationwide population-based cohort study'.²

Goulielmos and Zervou¹ commented that the occurrence of systemic lupus erythematosus (SLE) and autoimmune haemolytic anaemia (AIHA) may be related to genetic and environmental factors, in addition to the abnormal immune system function. In the recently presented data, Goulielmos and Zervou¹ showed that various shared genetic polymorphisms were associated with the increased susceptibility of ITP and SLE. According to existing research, it has been reported that AIHA and SLE share common genetic susceptibility as well, including human leucocyte antigen (HLA)-B8 and HLA-DQ6 regions,^{3,4} and single-nucleotide polymorphisms of the *CTLA-4*,⁵ *NFATC1*,⁶ *LT-a*,⁷ *FcγRIIa*,⁸ *FOXP3*⁹ and *CARD11*¹⁰ genes. We strongly agree with the point put forward by Goulielmos and Zervou; as early as 1982, Lippman *et al*¹¹ had found that a variety of autoimmune diseases are genetically related, including AIHA and SLE.

In our article, we used claims data from the Taiwanese National Health Insurance Database (NHIRD) to explore the incidence of SLE in patients with AIHA and demonstrated that patients with AIHA are at a very high risk of incidental SLE.² Due to lack of gene-related data in the NHIRD, we may not be able to directly calculate the genetic correlation between these two diseases. However, given that the registry of beneficiaries provides information on the relationships between the insured individuals and his/her dependents and only spouses and blood relatives are qualified to be dependents of an insured individual, we are able to identify parent–offspring relationships directly.¹²

Therefore, we conducted another population based-study using the 2003–2017 NHIRD to compare the risk of AIHA in the first-degree relatives (FDRs) of patients with SLE with AIHA and that in the FDRs of patients with SLE without AIHA. We identified 29 859 patients with SLE during 2003–2017. Of them, 726 patients had at least one hospitalisation with a diagnosis of AIHA and 28 875 patients never had a diagnosis of AIHA in outpatient or inpatient visits during 2003–2017. From this

nationwide SLE cohort, we identified 451 FDRs of patients with SLE with AIHA and 12 756 FDRs of patients with SLE without AIHA. We further matched these two FDR groups (1:2) for age and sex and selected 417 FDRs of patients with SLE with AIHA and 1668 matched FDRs of patients with SLE without AIHA. We compared the risk of AIHA development during 2003–2017 between FDRs of patients with SLE with AIHA and FDRs of patients with SLE without AIHA in the prematched and post-matched FDR cohorts using logistic regression and conditional logistic regression, respectively. As shown in table 1, the risk of AIHA tended to be higher in FDRs of patients with SLE with AIHA than those of patients with SLE without AIHA in the prematched cohort (OR 3.80, 95% CI 0.86 to 16.74, $p=0.077$).

In conclusion, patients with AIHA had increased risk of SLE, which was correlated with the same gene loci of them. We recommend that while evaluating the laboratory characteristics of patients with AIHA, clinicians should also pay attention to the medical history of patients' relatives, avoid delayed diagnosis of SLE and give appropriate treatment in time. At the same time, we also need to further study and identify the specific shared gene loci of AIHA and SLE so as to provide appropriate treatment as soon as possible.

Han-You Mo,^{1,2} Xiao-Huan Chen^{2,3}, James Cheng Chung Wei^{2,4,5}
Hsin-Hua Chen^{6,7,8,9,10}

¹Department of Rheumatology, The Affiliated Hospital of Guilin Medical University, Guilin, China

²Institute of Medicine, College of Medicine, Chung Shan Medical University, Taichung, Taiwan

³Department of Rheumatology, Guilin Medical University, Guilin, China

⁴Department of Allergy, Immunology & Rheumatology, Chung Shan Medical University Hospital, Taichung, Taiwan

⁵Graduate Institute of Integrated Medicine, China Medical University, Taichung, Taiwan

⁶Department of Medical Research, Taichung Veterans General Hospital, Taichung, Taiwan

⁷Division of Allergy, Immunology and Rheumatology, Department of Internal Medicine, Taichung Veterans General Hospital, Taichung, Taiwan

⁸Department of Industrial Engineering and Enterprise Information, Tunghai University, Taichung, Taiwan

⁹School of Medicine, National Yang-Ming University, Taipei, Taiwan

¹⁰Institute of Biomedical Science and Rong Hsing Research Centre for Translational Medicine, Chung Hsing University, Taichung, Taiwan

Correspondence to Dr Hsin-Hua Chen; shc5555@hotmail.com

Handling editor Josef S Smolen

Contributors H-YM and JCCW conceptualised the research and drafted the manuscripts. X-HC interpreted the data and drafted the manuscript. H-HC contributed to the research design, performed data analysis and graph generation, and critically revised the manuscript. All authors have read and approved the final manuscript.

Table 1 Risk of AIHA in FDRs of patients with SLE with AIHA compared with that in FDRs of patients with SLE patients without AIHA

	Prematched FDR cohort*				Postmatched FDR cohort†	
	Univariable		Multivariable		OR (95% CI)	P value
	OR (95% CI)	P value	OR (95% CI)	P value		
FDR group						
FDRs of patients with SLE with AIHA	Reference		Reference		Reference	
FDRs of patients with SLE without AIHA	3.73 (0.85 to 16.36)	0.081	3.80 (0.86 to 16.74)	0.077	>999 (<0.001 to >999)	0.998
Gender						
Female	Reference		Reference			
Male	0.12 (0.03 to 0.53)	0.005	0.13 (0.03 to 0.56)	0.007		
Age (year)	0.97 (0.93 to 1.01)	0.083	0.97 (0.93 to 1.01)	0.140		

*Logistic regression analyses.

†Conditional logistic regression analyses and for estimation of OR of AIHA risk in FDRs of patients with SLE with AIHA compared with that in FDRs of patients with SLE without AIHA.

AIHA, autoimmune haemolytic anaemia; FDR, first-degree relative; SLE, systemic lupus erythematosus.

Funding This work was supported by funding from Chung Shan Medical University Hospital grant number CSH-2018-C-023 and the National Natural Science Foundation of China Grants [81760298].

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Obtained.

Ethics approval This study was approved by the Institutional Review Board of Taichung Veterans General Hospital in Taiwan (approval number: CE19038A).

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

H-YM and JCCW contributed equally.

To cite Mo H-Y, Chen X-H, Wei JCC, *et al.* *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2020-219328

Received 9 November 2020

Accepted 10 November 2020

► <http://dx.doi.org/10.1136/annrheumdis-2020-219321>

Ann Rheum Dis 2020;**0**:1–2. doi:10.1136/annrheumdis-2020-219328

ORCID iDs

Xiao-Huan Chen <http://orcid.org/0000-0002-5482-4756>

James Cheng Chung Wei <http://orcid.org/0000-0003-0310-2769>

Hsin-Hua Chen <http://orcid.org/0000-0002-7304-4587>

REFERENCES

- Goulielmos G, Zervou M. Correspondence on increased risk of systemic lupus erythematosus in patients with autoimmune haemolytic anaemia: a nationwide population-based cohort study. *Ann Rheum Dis* 2020.
- Mo H-Y, Wei JCC, Chen X-H, *et al.* Increased risk of systemic lupus erythematosus in patients with autoimmune haemolytic anaemia: a nationwide population-based cohort study. *Ann Rheum Dis* 2020. doi:10.1136/annrheumdis-2020-218886. [Epub ahead of print: 22 Sep 2020].
- Barcellini W, Giannotta J, Fattizzo B. Autoimmune hemolytic anemia in adults: primary risk factors and diagnostic procedures. *Expert Rev Hematol* 2020;13:585–97.
- Rezaieyazdi Z, Tavakkol-Afshari J, Esmaili E, *et al.* Association of HLA-DQB1 allelic sequence variation with susceptibility to systemic lupus erythematosus. *Iran J Allergy Asthma Immunol* 2008;7:91–5.
- Pavkovic M, Georgievski B, Cevreska L, *et al.* Ctl4-4 exon 1 polymorphism in patients with autoimmune blood disorders. *Am J Hematol* 2003;72:147–9.
- Lanata CM, Nititham J, Taylor KE, *et al.* Genetic contributions to lupus nephritis in a multi-ethnic cohort of systemic lupus erythematosus patients. *PLoS One* 2018;13:e0199003.
- D'Abronzio LS, Barros MMO, Bordin JO, *et al.* Analysis of polymorphisms of TNF- α , IL-1 α , IL-10, IL-12 and CTLA-4 in patients with warm autoimmune haemolytic anaemia. *Int J Lab Hematol* 2012;34:356–61.
- Khoa PD, Sugiyama T, Yokochi T. Fc gamma receptor II polymorphism in Vietnamese patients with systemic lupus erythematosus. *Lupus* 2003;12:704–6.
- Lee HJ, Li CW, Hammerstad SS, *et al.* Immunogenetics of autoimmune thyroid diseases: a comprehensive review. *J Autoimmun* 2015;64:82–90.
- Liu Z, Yu Y, Yue Y, Zheng L, Yangsheng Y Y, *et al.* Genetic alleles associated with SLE susceptibility and clinical manifestations in Hispanic patients from the Dominican Republic. *Curr Mol Med* 2019;19:164–71.
- Lippman SM, Arnett FC, Conley CL, *et al.* Genetic factors predisposing to autoimmune diseases. autoimmune hemolytic anemia, chronic thrombocytopenic purpura, and systemic lupus erythematosus. *Am J Med* 1982;73:827–40.
- Kuo C-F, Grainge MJ, See L-C, *et al.* Familial aggregation of gout and relative genetic and environmental contributions: a nationwide population study in Taiwan. *Ann Rheum Dis* 2015;74:369–74.