

Response to: 'Correspondence on 'Systemic sclerosis and the COVID-19 pandemic: World Scleroderma Foundation preliminary advice for patient management' by Snarskaya and Vasileva

We read with great interest the response from Snarskaya and Vasileva.¹ A second wave of SARS-CoV-2 infection is now a global reality²: since the publication of the World Scleroderma Foundation preliminary advice on the management on systemic sclerosis (SSc) patients during the first COVID-19 pandemic,³ growing evidence has accumulated on COVID-19 affected SSc patients.⁴ Up to 22 June, at the end of the first wave, 25 cases of SSc-COVID-19 were published in the literature, most of them undergoing immunosuppressant treatment with corticosteroids (CCS), conventional synthetic disease modifying antirheumatic drugs (csDMARDs) (methotrexate or mycophenolate mofetil) or biological DMARD (bDMARDs) (rituximab or tocilizumab) and with a variable outcome.⁴ This heterogeneity was a limitation in understanding the possible influence of ongoing immunosuppression and the effect of administered treatments. A recent meta-analysis, including 62 studies with more than 300 000 patients with autoimmune diseases, concluded that bDMARD monotherapy was a protective factor for COVID-19-related hospitalisation and death, in particular with antitumour necrosis factor alpha, while the use of CCS, conventional-synthetic DMARDs (csDMARD) and csDMARDs/bDMARD combination exposed the patients to a higher risk.⁵ A focused analysis on SSc population has described more SSc-COVID-19 cases: 1/64 patients in Crisafulli *et al*,⁶ 1/168 patients in Bellan *et al*⁷ and 2/526 patients in Del Papa *et al*,⁸ with one of them, previously exposed to methotrexate and rituximab, with a fatal outcome.

Other major rheumatological scientific societies have released their advice for the management of adult patients affected by rheumatic diseases during the COVID-19 pandemic.^{9, 10} The European League against Rheumatism (EULAR) and the American College of Rheumatology (ACR) suggested the continuation of CCS, csDMARD and bDMARD to all patients without suspected or confirmed COVID-19 infection. In addition to the same context, the ACR advised for the initiation of immunosuppressive treatment in patients with newly diagnosis or active rheumatic disease.¹⁰ Regarding COVID-19 positive cases among patients with rheumatic conditions, the EULAR suggested to continue chronic CCS also in case of symptomatic COVID-19

infection, to evaluate the continuation of other immunosuppressive drugs (IMS) on a case-by-case basis and to adhere with local recommendations for the treatment of COVID-19 disease.⁹ For SARS-CoV-2 exposed patients, the ACR recommends immunosuppressive treatments temporary interruption and to restart them on a negative test results or 2 weeks observation without symptoms. Similarly, it is suggested to stop or postpone the initiation of the immunosuppressants in case of documented or highly suspected COVID-19 disease.¹⁰ In the literature, a case-by-case evaluation was suggested for interleukin (IL)-6 inhibitors,¹⁰ possibly also considering Janus Kinase inhibitors.¹¹ Finally, the ACR suggested the restart of the immunosuppressive treatment in patients with non-complicated COVID-19 disease after 7–14 days of symptom resolution or 10–17 days after SARS-CoV-2 positive PCR test but without any symptom development.¹⁰ Regarding more severe cases of COVID-19 disease, again, a case-by-case evaluation is strongly advised.¹⁰

Today, the interaction between the immunosuppressive treatments and the COVID-19 is a matter of debate, and the main actors in this scenario are, on one side, the immunosuppressants to control SSc and, on the other side, the immunosuppressive treatments targeting the COVID-19-related hyperinflammatory syndrome (eg, CCS, anticytokines such as anti-IL-1 and anti-IL-6, and JAK inhibitors).¹²

In this second pandemic wave, it may be wise to shift the therapeutic focus from the rheumatic manifestations to the potential infectious complications. In figure 1, the different clinical scenarios are presented: the case of non-exposure allows the continuation, while the exposure and/or the test/serology positivity always suggest a treatment discontinuation and a restart in due time according to the test positivity/negativity. Following the precautionary advice of EULAR and ACR, we may propose that the interruption (and not the continuation) of the immunosuppressive treatment should be discussed case-by-case, in particular regarding patients exposed to B-depleting agents. Through its direct target on CD20 positive B cells, rituximab may potentially reduce the immune response against the virus and increase the risk of reinfection.¹³ Patients exposed to rituximab may benefit from a temporary interruption of other concomitant csDMARDs and/or the postponement of the next treatment cycle, with a more intensive monitoring of COVID-19-related and SSc-related manifestations. Despite the negative result of the data testing the therapeutic effect of convalescent plasma in severe COVID-19 pneumonia,¹⁴ CD-20 depleted patients with might benefit from the use of blood-derived




Figure 1 Algorithm for the management of immunosuppressive treatment according to SARS-CoV-2 exposure and COVID-19 disease. ab, antibodies; IMS, immunosuppressive drugs.

COVID-19 convalescent plasma, given the promising results shown in a small cohort of patients.¹⁵

We herein agree with the suggestions delivered from other societies and consider the possible role of PCR-retesting and anti-COVID-19 antibodies measuring before restarting IMS treatment in SSc-patients with initially positive PCR test. This preliminary consideration will be updated once data from larger initiatives, such as the EUSTAR COVID-19 registry, the EULAR COVID-19 database and the global alliance COVID-19 registry, will provide more robust evidences.

Cosimo Bruni ,¹ Marco Matucci-Cerinic ^{1,2}

¹Department of Experimental and Clinical Medicine, Division of Rheumatology, Università degli Studi di Firenze, Firenze, Italy

²Department of Geriatric Medicine, Division of Rheumatology, Azienda Ospedaliero Universitaria Careggi, Firenze, Italy

Correspondence to Dr Marco Matucci-Cerinic, Department of Experimental and Clinical Medicine, Division of Rheumatology, Università degli Studi di Firenze, Firenze 50121, Italy; marco.matuccicerinic@unifi.it

Handling editor Josef S Smolens

Acknowledgements We acknowledge Professor Yannick Allanore, Professor Massimo Clementi, Professor Lorenzo Dagna, Professor Nemanja Damjanov, Professor Amato De Paulis, Professor Christopher Denton, Professor Oliver Distler, Professor David Fox, Professor Daniel E Furst, Professor Dinesh Khanna, Professor Thomas Krieg, Professor Masataka Kuwana, Professor Eun Bong Lee, Professor Mengtao Li, Professor Shiv Pillai, Dr Yukai Wang, Professor Xiaofeng Zeng and Professor Gloria Taliani for their participation to the original manuscript.

Contributors CB and MM-C conceived, drafted and revised the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests MM-C reports grants and personal fees from Actelion, personal fees from Biogen, personal fees from Bayer, personal fees from Boehringer Ingelheim, personal fees from CSL Behring, personal fees from Eli-Lilly, outside the submitted work. CB reports personal fees from Actelion, personal fees from Eli Lilly, grants from Gruppo Italiano Lotta alla Sclerodermia (GILS), grants from New Horizon Fellowship, grants from European Scleroderma Trials and Research (EUSTAR), grants from Foundation for Research in Rheumatology (FOREUM), grants from Fondazione Italiana per la Ricerca sull'Artrite (FIRA), outside the submitted work.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Bruni C, Matucci-Cerinic M. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2020-219533

Received 24 November 2020

Revised 19 December 2020

Accepted 21 December 2020



► <http://dx.doi.org/10.1136/annrheumdis-2020-219489>

Ann Rheum Dis 2021;0:1–2. doi:10.1136/annrheumdis-2020-219533

ORCID iDs

Cosimo Bruni <http://orcid.org/0000-0003-2813-2083>

Marco Matucci-Cerinic <http://orcid.org/0000-0002-9324-3161>

REFERENCES

- Snarskaya ES, Vasileva KD. Correspondence on 'Systemic sclerosis and the COVID-19 pandemic: World Scleroderma Foundation preliminary advice for patient management'. *Ann Rheum Dis* 2020.
- Denton CP, Campochiaro C, Bruni C. COVID-19 and systemic sclerosis: rising to the challenge of a pandemic. *J Scleroderma Relat Disord* 2020;2397198320963393.
- Matucci-Cerinic M, Bruni C, Allanore Y, et al. Systemic sclerosis and the COVID-19 pandemic: world scleroderma Foundation preliminary advice for patient management. *Ann Rheum Dis* 2020;79:724–6.
- Russo B, Ben Shimol J, Matucci-Cerinic M. Systemic sclerosis and COVID-19: what's new in the literature. *Clin Exp Rheumatol* 2020.
- Akiyama S, Hamdeh S, Micic D, et al. Prevalence and clinical outcomes of COVID-19 in patients with autoimmune diseases: a systematic review and meta-analysis. *Ann Rheum Dis* 2020. doi:10.1136/annrheumdis-2020-218946. [Epub ahead of print: 13 Oct 2020].
- Crisafulli F, Lazzaroni M-G, Zingarelli S, et al. Impact of COVID-19 on outpatient therapy with iloprost for systemic sclerosis digital ulcers. *J Scleroderma Relat Disord* 2020;79:239719832095229. doi:10.1177/2397198320952299
- Bellan M, Parisi S, Stobbione P, et al. Impact of the COVID-19 outbreak on an Italian cohort of systemic sclerosis patients. *Scand J Rheumatol* 2020;49:505–6.
- Del Papa N, Sambataro G, Minniti A, et al. Impact of COVID-19 outbreak in an Italian cohort of patients with systemic sclerosis. *Ther Adv Musculoskelet Dis* 2020;12:1759720X20953356.
- Landewé RB, Machado PM, Kroon F, et al. EULAR provisional recommendations for the management of rheumatic and musculoskeletal diseases in the context of SARS-CoV-2. *Ann Rheum Dis* 2020;79:851–8.
- Mikulis TR, Johnson SR, Fraenkel L, et al. American College of rheumatology guidance for the management of rheumatic disease in adult patients during the COVID-19 pandemic: version 2. *Arthritis Rheumatol* 2020;72:e1–12.
- Orlandi M, Lepri G, Bruni C, et al. The systemic sclerosis patient in the COVID-19 era: the challenging crossroad between immunosuppression, differential diagnosis and long-term psychological distress. *Clin Rheumatol* 2020;39:2043–7.
- Gremese E, Brondani G, Apollonio L. Correspondence on 'Prevalence and clinical outcomes of COVID-19 in patients with autoimmune diseases: a systematic review and meta-analysis'. *Ann Rheum Dis* 2020.
- Mehta P, Porter JC, Chambers RC, et al. B-Cell depletion with rituximab in the COVID-19 pandemic: where do we stand? *Lancet Rheumatol* 2020;2:e589–90.
- Simonovich VA, Burgos Pratz LD, Scibona P, et al. A randomized trial of convalescent plasma in Covid-19 severe pneumonia. *N Engl J Med* 2020. doi:10.1056/NEJMoa2031304. [Epub ahead of print: 24 Nov 2020].
- Hueso T, Pouderoux C, Péré H, et al. Convalescent plasma therapy for B-cell-depleted patients with protracted COVID-19. *Blood* 2020;136:2290–5.