Response to: 'Correspondence on 'Interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial' by de Wolff *et al*

We read with interest the correspondence on our article 'Interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial' by Wolff *et al*¹ in which the use of a composite endpoint, combining disease activity, functional and serological parameters, the newly developed CRESS² in future primary Sjögren's syndrome (pSS) trials is discussed, to lower placebo response and facilitate inclusions in clinical trials.

As suggested by Wolff *et al*, a composite endpoint including multiple important pSS features might reveal more appropriate than only Euler Sjögren Syndrome Disease Activity index (ESSDAI) or only gland function as primary endpoint in this complex and heterogeneous disease to lower placebo response. We agree this is an important point to consider for future trials. As mentioned in our reply³ to the correspondence from Wang *et al*,⁴ NECESSITY, a European initiative, will combine data from our trial⁵ with those of previous randomised trials to determine new clinical composite outcomes in pSS, capturing both systemic and glandular features of the disease. An initiative from OMERACT on clinical outcomes in pSS is also ongoing on this crucial topic.

Wolff *et al* also suggest that another advantage of a composite endpoint might be to facilitate a broader inclusion of patients with pSS in clinical trials, that is, to allow the inclusion of patients with either an active systemic disease or a high burden of symptoms such as fatigue or dryness. However, in pSS, the pathogenesis of fatigue and dryness remains uncertain, with no key pathogenic cytokine or cell population identified to date. Targeting a unique goal to achieve a relevant improvement concomitantly in systemic activity, fatigue and dryness remain a great challenge in pSS. Indeed, even in the few recent positive phase II clinical trials, no drug was capable to improve all these three dimensions of the disease. Therefore, focusing on the evaluation of symptoms or systemic manifestations related to the pathway targeted might reveal more straightforward than a composite outcome. To increase the odds of finding effective drugs, biomarkers are also needed to stratify patients according to a molecular endotype amenable to a specific targeted therapy.

We therefore definitively hope that a relevant composite score will emerge from CRESS, NECESSITY and OMERACT initiatives and that the ongoing collective effort to clarify the pathogenesis of the disease will allow to make pSS enter the individualised medicine age.

Renaud Felten ^{1,2} Jacques-Eric Gottenberg ^{1,2}

¹Rheumatology Department, National Reference Center for Rare East South-West Autoimmune Systemic Diseases RESO, Strasbourg University Hospitals, Strasbourg, Alsace, France

²CNRS, UPR3572, Institute of Molecular and Cellular Biology (IBMC), Immunology Laboratory, Immunopathology and Therapeutic Chemistry, University of Strasbourg, Strasbourg, France

Correspondence to Dr Renaud Felten, Rheumatology Department, National Reference Center for Rare East South-West Autoimmune Systemic Diseases RESO, Strasbourg University Hospitals, 1 avenue Molière, 67098 Strasbourg, Alsace, France; renaud.felten@chru-strasbourg.fr

Handling editor Josef S Smolen

Contributors Both authors contributed to the concept, design and writing of the letter and have approved the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Felten R, Gottenberg J-E. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2021-220220

Received 21 March 2021

Accepted 22 March 2021

► <http://dx.doi.org/10.1136/annrheumdis-2021-220198>

Ann Rheum Dis 2021;0:1. doi:10.1136/annrheumdis-2021-220220

ORCID iDs

Renaud Felten <http://orcid.org/0000-0002-4951-4032>

Jacques-Eric Gottenberg <http://orcid.org/0000-0002-9469-946X>

REFERENCES

- 1 Wolff L, Arends S, Verstappen G. Correspondence on interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial. *Ann Rheum Dis* 2020.
- 2 Arends S, de WL, van NJ, *et al*. Composite of Relevant Endpoints for Sjögren's Syndrome (CRESS) [abstract]. *Arthritis Rheumatol* 2020;72.
- 3 Felten R, Gottenberg J-E. Response to: 'Correspondence on 'Interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial' by Wang *et al*. *Ann Rheum Dis* 2021. doi:10.1136/annrheumdis-2021-219882
- 4 Wang B, Chen S, Xuan J, *et al*. Correspondence on 'Interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial'. *Ann Rheum Dis*;18:annrheumdis-2020-219666.
- 5 Felten R, Devauchelle-Pensec V, Seror R, *et al*. Interleukin 6 receptor inhibition in primary Sjögren syndrome: a multicentre double-blind randomised placebo-controlled trial. *Ann Rheum Dis* 2020. doi:10.1136/annrheumdis-2020-218467