Correspondence on 'NCF1-339 polymorphism is associated with altered formation of neutrophil extracellular traps, high serum interferon activity and antiphospholipid syndrome in systemic lupus erythematosus' by Linge *et al*

We read with great interest the article by Linge *et al*¹ on 'NCF1-339 polymorphism is associated with altered formation of neutrophil extracellular traps, high serum interferon activity and antiphospholipid syndrome in systemic lupus erythematosus'¹ as well as the response of Linge and Bengtsson² to the comment by Joob and Wiwanitkit,³ who raised a question regarding the importance and the role of the NCF1-339 rs201802880 (p. Arg90His) polymorphism in systemic lupus erythematosus (SLE). Linge and Bengtsson³ nicely placed this polymorphism in a functional context, pointing out that the amino acid shift from arginine to histidine at a PX domain of the NCF1 protein (a domain of crucial importance in the membrane binding⁴ reduces reactive oxygen species (ROS) response.⁵ Of note, the rs201802880 p. Arg90His variant of the *NCF1* gene has been associated, apart from SLE, with rheumatoid arthritis and Sjögren's syndrome in adult patients⁶ as well as with early-onset interferonopathy in paediatric patients.⁷

Our aim with this letter was to extend the information given by Linge and Bengtsson³ by presenting data that elucidate further the significance of the rs201802880 Arg90His variant from another viewpoint, using data from a structural biology approach, thus analysing this variant in the various genetically controlled functions that is involved, including the neutrophil extracellular trap formation, the reduced extracellular ROS production in neutrophils and the decreased NADPH oxidase function.¹ To this end, we have recently evaluated the structural significance of the rs201802880 variant on the p47^{phox} PX domain of the NCF1 protein by constructing a three-dimensional structural model, focusing on the phosphate-binding pocket of NCF1 p47^{phox} domain with the electrostatic molecular surface.⁸ Our molecular investigation showed that in this variant, the functionality of the p47^{phox} PX cytosolic subunit of neutrophil NADPH oxidase has been modified, leading to affinity reduction to PtdIns(3,4)P2 caused by the loss of specific phosphoinositide head group interactions and affecting the p47^{phox} translocation to the plasma membrane.

The structural biological information reported here may help the interpretation of the findings of Linge *et al*¹ from the structural-functional point of view and, apparently, is in agreement with the statements in the comment of Linge and Bengtsson³ that emphasise on the crucial importance of the NCF1-339 polymorphism under discussion.

George N Goulielmos^{1,2}, Maria I Zervou,¹ Elias Eliopoulos³

¹Section of Molecular Pathology and Human Genetics, Department of Internal Medicine, School of Medicine, University of Crete, Heraklion, Greece
²Department of Internal Medicine, University Hospital of Heraklion, Heraklion, Greece
³Laboratory of Genetics, Department of Biotechnology, Agricultural University of Athens, Athens, Greece

Correspondence to Dr George N Goulielmos, Section of Molecular Pathology and Human Genetics, Department of Internal Medicine, School of Medicine, University of Crete, Heraklion 74100, Greece; goulielmos@med.uoc.gr

Handling editor Josef S Smolen

Contributors GNG, MIZ and EE prepared the manuscript and approved the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Goulielmos GN, Zervou MI, Eliopoulos E. *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2021-221871

Received 17 November 2021

Accepted 18 November 2021

Ann Rheum Dis 2022;0:1. doi:10.1136/annrheumdis-2021-221871

ORCID iD

George N Goulielmos <http://orcid.org/0000-0002-9797-2310>

REFERENCES

- 1 Linge P, Arve S, Olsson LM, *et al*. NCF1-339 polymorphism is associated with altered formation of neutrophil extracellular traps, high serum interferon activity and antiphospholipid syndrome in systemic lupus erythematosus. *Ann Rheum Dis* 2020;79:254–61.
- 2 Linge CP, Bengtsson A. Response to: 'NCF1-339 polymorphism and systemic lupus erythematosus' by Joob and Wiwanitkit. *Ann Rheum Dis* 2021;80:e195.
- 3 Joob B, Wiwanitkit V. NCF1-339 polymorphism and systemic lupus erythematosus. *Ann Rheum Dis* 2021;80:e194.
- 4 Zhong J, Olsson LM, Urbanaviciute V, *et al*. Association of Nox2 subunits genetic variants with autoimmune diseases. *Free Radic Biol Med* 2018;125:72–80.
- 5 Li XJ, Marchal CC, Stull ND, *et al*. P47Phox phox homology domain regulates plasma membrane but not phagosome neutrophil NADPH oxidase activation. *J Biol Chem* 2010;285:35169–79.
- 6 Yokoyama N, Kawasaki A, Matsushita T, *et al*. Association of NCF1 polymorphism with systemic lupus erythematosus and systemic sclerosis but not with ANCA-associated vasculitis in a Japanese population. *Sci Rep* 2019;9:16366.
- 7 Schnappauf O, Heale L, Dissanayake D, *et al*. Homozygous variant P. Arg90His in NCF1 is associated with early-onset Interferonopathy: a case report. *Pediatr Rheumatol Online J* 2021;19:54.
- 8 Goulielmos GN, Zervou MI, Eliopoulos E. Comment on: homozygous variant P. Arg90His in NCF1 is associated with early-onset interferonopathy: a case report. *Pediatr Rheumatol Online J* 2021;19:125.