EULAR recommendations for cardiovascular risk management in rheumatic and musculoskeletal diseases, including systemic lupus erythematosus and antiphospholipid syndrome

George C Drosos ¹, Daisy Vedder ², Eline Houben,³ Laura Boekel ², Fabiola Atzeni,⁴ Sara Badreh,⁵ Dimitrios T Boumpas ^{6,7}, Nina Brodin,^{8,9} Ian N Bruce,^{10,11} Miguel Ángel González-Gay ¹², Søren Jacobsen ^{13,14}, György Kerekes,¹⁵ Francesca Marchiori,¹⁶ Chetan Mukhtyar ¹⁷, Manuel Ramos-Casals,¹⁸ Naveed Sattar,¹⁹ Karen Schreiber,²⁰ Savino Sciascia ²¹, Elisabet Svenungsson ²², Zoltan Szekanecz ²³, Anne-Kathrin Tausche,²⁴ Alan Tyndall,²⁵ Vokko van Halm,²⁶ Alexandre Voskuyl,²⁷ Gary J Macfarlane ²⁸, Michael M Ward ²⁹, Michael T Nurmohamed,^{2,30} Maria G Tektonidou ^{1,7}

Handling editor Josef S Smolen

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2021-221733>).

For numbered affiliations see end of article.

Correspondence to

Professor Maria G Tektonidou, First Department of Propaedeutic Internal Medicine, National and Kapodistrian University of Athens, Athens, Attica, Greece; mtektionidou@gmail.com

GCD, DV and EH contributed equally.
MTN and MGT contributed equally.

GCD, DV and EH are joint first authors.
MTN and MGT are joint senior authors.

Received 23 October 2021
Accepted 5 January 2022

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Drosos GC, Vedder D, Houben E, *et al.* *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2021-221733

ABSTRACT

Objective To develop recommendations for cardiovascular risk (CVR) management in gout, vasculitis, systemic sclerosis (SSc), myositis, mixed connective tissue disease (MCTD), Sjögren's syndrome (SS), systemic lupus erythematosus (SLE) and antiphospholipid syndrome (APS).

Methods Following European League against Rheumatism (EULAR) standardised procedures, a multidisciplinary task force formulated recommendations for CVR prediction and management based on systematic literature reviews and expert opinion.

Results Four overarching principles emphasising the need of regular screening and management of modifiable CVR factors and patient education were endorsed. Nineteen recommendations (eleven for gout, vasculitis, SSc, MCTD, myositis, SS; eight for SLE, APS) were developed covering three topics: (1) CVR prediction tools; (2) interventions on traditional CVR factors and (3) interventions on disease-related CVR factors. Several statements relied on expert opinion because high-quality evidence was lacking. Use of generic CVR prediction tools is recommended due to lack of validated rheumatic diseases-specific tools. Diuretics should be avoided in gout and beta-blockers in SSc, and a blood pressure target <130/80 mm Hg should be considered in SLE. Lipid management should follow general population guidelines, and antiplatelet use in SLE, APS and large-vessel vasculitis should follow prior EULAR recommendations. A serum uric acid level <0.36 mmol/L (<6 mg/dL) in gout, and disease activity control and glucocorticoid dose minimisation in SLE and vasculitis, are recommended. Hydroxychloroquine is recommended in SLE because it may also reduce CVR, while no particular immunosuppressive treatment in SLE or urate-lowering therapy in gout has been associated with CVR lowering.

Conclusion These recommendations can guide clinical practice and future research for improving CVR management in rheumatic and musculoskeletal diseases.

INTRODUCTION

Patients with inflammatory rheumatic diseases have an increased risk of cardiovascular disease,¹ in comparison to the general population, which prompted the development (2010) and update (2015/16) of European League against Rheumatism (EULAR) recommendations for cardiovascular risk (CVR) management in patients with rheumatoid arthritis (RA), ankylosing spondylitis and psoriatic arthritis.² Accumulating evidence has shown elevated cardiovascular morbidity and mortality in other rheumatic and musculoskeletal diseases (RMDs) including gout, vasculitis, systemic sclerosis (SSc), myositis, mixed connective tissue disease (MCTD), Sjögren's syndrome (SS), systemic lupus erythematosus (SLE) and the antiphospholipid syndrome (APS).^{3–13} Estimations of the incidence of cardiovascular events vary among the different disease groups (Supplementary systematic literature review (SLR) report, section II).

The higher CVR in patients with rheumatic diseases is not sufficiently explained by differences in the prevalence of traditional CVR factors,^{14–18} suggesting that specific treatment recommendations tailored to patients with these conditions are needed. Chronic inflammation has been considered a key feature in cardiovascular disease pathogenesis in RMDs,¹⁹ demonstrated also in the general population by associations with serum C-reactive protein (CRP) levels^{20 21} and the efficacy of medications targeting inflammatory pathways,^{22–24} while new links between inflammation, immunity and cardiometabolic factors are being researched.²⁵ Furthermore, patients with RMDs are often exposed to immunomodulators and glucocorticoids. Although better control of inflammation may reduce CVR in individual patients,^{23 24} it is not known if some side effects of these medications might outweigh any anti-inflammatory benefit, thereby increasing the CVR.

Recommendation

Therefore, a EULAR Task Force was formed to develop recommendations for the management of CVR in patients with SLE, APS, gout, vasculitis, SSc, myositis, MCTD and SS based on an evidence-based approach and experts' consensus.

METHODS

Task force

Two convenors (MTN and MGT) guided the task force together with two methodologists (GJM and MMW) and four fellows (DV, GCD, EH and LB), responsible for the SLRs. Furthermore, the task force included 20 members from 11 European countries: 12 rheumatologists, 2 cardiologists, 1 metabolic medicine physician, 1 healthcare professional, 2 patient representatives and 2 EMerging EULAR NETwork members (KS and SS). The process followed the updated EULAR standardised operating procedures²⁶ and the Appraisal of Guidelines for Research and Evaluation II instrument.²⁷

At the initial task force meeting, a first set of research questions, prepared by the convenors, was discussed with the panel and formulated on four major topics: use of cardiovascular prediction tools; interventions targeting traditional CVR factors; interventions targeting disease-related CVR factors and prevalence/incidence of cardiovascular disease. Thereafter, final research questions were developed using the PICO format (P, population; I, intervention; C, comparator; O, outcomes).

Collection of evidence

A comprehensive SLR was performed by two groups working in parallel: the gout, vasculitis, SSc/myositis/MCTD/SS group (convenor: MTN; methodologist: GJM; fellows: DV, EH and LB), and the SLE and APS group (convenor: MGT; methodologist: MMW; fellow: GCD). The protocol for the literature search was based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement.²⁸ Search terms were developed with the help of experienced librarians of the VU Amsterdam, Northwest Clinics Alkmaar (for gout, vasculitis, SSc, myositis, MCTD and SS SLRs) and the National Institutes of Health, USA (for SLE and APS SLRs). PubMed, Embase and the Cochrane Library were searched for full-length English-language published articles from their inception to March 2020, while searches for incidence and prevalence of cardiovascular events were extended up to November 2020. Exclusion criteria and the search terms for each disease separately are presented in the Supplementary SLR report (section IA). The outcome was cardiovascular events rather than surrogate markers of cardiovascular disease.

Data abstraction is described in Supplementary SLR report (section IB). Retrieved studies were screened by title and abstract and articles selected for full text review were then examined independently by two persons for each group (DV, EH, LB, MN, CM, and GCD, MGT and MMW) with consultation of other task force members. A number of individually searched articles (one for gout,²⁹ three for SLE/APS^{30–32} published after the initial search periods were included due to their importance. Data extraction was performed by the fellows (DV, EH and LB) and CM under supervision of MN and GJM in the gout, vasculitis, SSc, myositis, MCTD and SS group, and by GCD, MGT and MMW in the SLE and APS group. Quality assessment was performed using the Cochrane risk-of-bias tool³³ for randomised clinical trials and the Newcastle-Ottawa Scale³⁴ for observational studies. Formal pooling and meta-analysis of risks could not be performed due to the diversity of outcomes, exposures and measures of association reported in the primary studies.

Evidence summaries and draft recommendations were formulated for review by all task force members before the second meeting.

Consensus on statements

The virtual second task force meeting included the presentation of SLR results and discussion and editing of the first draft of recommendations. Recommendations were accepted when $\geq 75\%$ of the task force members voted agreement. After additional discussions on wording changes and voting on text, a final set of recommendations and overarching principles was prepared, including the level of evidence (LoE) and grade of recommendation (GoR) according to the Oxford Centre for Evidence Based Medicine system.³⁵ All task force members indicated their level of agreement (LoA) for each recommendation (0, no agreement at all; 10, full agreement), and results were averaged. The manuscript was reviewed and approved by all task force members and the EULAR Executive Committee before submission.

RESULTS

For gout, vasculitis, SSc, myositis, MCTD and SS, 105 articles were included in the SLR, while for SLE and APS, 75 articles were included (figures 1 and 2). SLR results including the flow chart and evidence tables for each PICO are presented in Supplementary SLR report (section II); all articles included in the SLRs are shown in section III.

Overarching principles

The task force developed four overarching principles emphasising the need for increased awareness of elevated CVR in RMDs, regular CVR screening, assessment and management of modifiable CVR factors, and patient education about CVR, treatment adherence and lifestyle changes (table 1).

Recommendations

Gout, vasculitis, SSc, myositis, MCTD and SS

CVR prediction tools

1. In patients with gout, vasculitis, SSc, myositis, MCTD and SS, we recommend thorough assessment of traditional CVR factors. The use of cardiovascular prediction tools as for the general population is recommended. (LoE: 5, GoR: D)

No studies have investigated the accuracy of cardiovascular prediction tools in patients with gout, vasculitis, SSc, myositis, MCTD and SS. It is currently uncertain to what extent the elevated risk for cardiovascular disease is driven by an increased prevalence of traditional or disease-specific risk factors. Existing tools, such as the Framingham Risk Score (FRS), QRISK3 or Systematic Coronary Risk Evaluation (SCORE) have been based on large general population cohorts with long follow-ups.^{36–38} Therefore, for gout, vasculitis, SSc, myositis, MCTD and SS, we recommend the use of prediction tools developed in the general population.

2. For ANCA-associated vasculitis the Framingham score may underestimate the CVR. Information from the European Vasculitis Society (EUVAS) model may supplement modifiable Framingham risk factors and is recommended to take into account. (LoE: 2b, GoR: D)

In patients with ANCA-associated vasculitis the observed incidence of cardiovascular events exceeded Framingham predicted incidence in two studies.^{39 40} Furthermore, one study on CVR in ANCA-associated vasculitis found a higher area under the curve (AUC) for the EUVAS model (AUC 0.73) based on age, diastolic

Figure 1 Flow chart of systematic literature review for cardiovascular risk management in gout, vasculitis, systemic sclerosis, myositis, mixed connective tissue disease and Sjögren's syndrome. Articles on cardiovascular incidence and prevalence are also included.

hypertension, and PR3 ANCA status in comparison with the Framingham model (AUC 0.65).⁴¹ Although this study was not designed for the evaluation of CVR, these disease-specific factors could be used for risk assessment in addition to Framingham risk factors but further work is needed to validate these findings.

Interventions targeting traditional CVR factors

3. *In patients with gout, vasculitis, SSc, myositis, MCTD, and SS, blood pressure (BP) management should follow recommendations used in the general population. (LoE: 5, GoR: D)*

We found no trials that assessed the use of antihypertensive treatment in these patients. One small retrospective cohort study found an increase of severe cranial ischaemic events in patients with giant-cell arteritis (GCA) treated with beta blockers.⁴² One large prospective cohort study in SSc found a protective effect of calcium channel blockers (CCB), ACE inhibitors (ACEI), and angiotensin receptor blockers (ARB) with ventricular arrhythmias.⁴³ Both studies did not control for confounding by indication. Altogether, currently, there is no evidence to modify the hypertension treatment target levels in patients with gout,

vasculitis, SSc, myositis, MCTD and SS from those used in the general population.

4. *In patients with gout, diuretics should be avoided. (LoE: 5, GoR: D)*

Following the EULAR recommendations on management of gout, use of thiazide and loop diuretics should be avoided, if possible, because of their effect to increase serum uric acid (SUA) levels.⁴⁴ Instead, the use of CCB or losartan could be considered. This topic was not updated as part of this guideline as the literature search focused on the effect of antihypertensives on cardiovascular outcomes and not on potential effect on SUA levels.

5. *In patients with SSc beta blockers should be avoided. (LoE: 5, GoR: D)*

Although large trials are lacking and therefore based on expert opinion, beta blockers are considered contraindicated due to their effect on Raynaud's phenomenon.

6. *In patients with gout, vasculitis, SSc, myositis, MCTD, and SS, lipid management should follow recommendations used in the general population. (LoE: 5, GoR: D)*

Figure 2 Flow chart of systematic literature review for cardiovascular risk management in systemic lupus erythematosus and the antiphospholipid syndrome.

In gout patients, no studies evaluated the effect of statins on cardiovascular disease or mortality in comparison with the general population. Two retrospective cohort studies suggested a protective effect of statins on mortality in patients with gout after 5 and 10 years, relative to patients not using statins.^{45 46} Because of the limited evidence, we recommend following guidelines on lipid management for the general population. Furthermore, myotoxicity as side effect of the combination of a statin and prophylactic colchicine (0.5 mg/day) is rare and routine discontinuation of the statin is not recommended.⁴⁷

Three studies in patients with GCA did not find an association between statins and cardiovascular events,^{42 48 49} but a fourth study of 103 patients with GCA, 28 of whom were treated with statins, reported a lower risk of cardiovascular hospitalisations with a longer cumulative duration of statin treatment (HR 0.993 per one additional daily dose).⁵⁰ No studies controlled for confounding by indication.

7. In patients with gout, vasculitis, SSc, myositis, MCTD, and SS, standard use of low-dose aspirin for primary prevention is not recommended. Treatment with platelet inhibitors should follow

recommendations used in the general population. (LoE: 2b/5, GoR: D)

In 2009 EULAR recommended the use of aspirin for prevention of cardiovascular and cerebrovascular events in individuals with large vessel vasculitis (LoE: 3, GoR: C).⁵¹ More recently the American College of Rheumatology (ACR) has used the same literature base to conditionally recommend the use of aspirin in flow critical large vessel vasculitis.⁵² However, in 2020 an update of the 2009 EULAR recommendations reappraised this evidence and concluded that the risk-benefit analysis was not favourable, and blanket use of antiplatelets was not essential unless indicated for other reasons.⁵³ Based on newly published studies, we agree with the 2020 iteration.^{41 48 49} In patients with gout, ANCA-associated vasculitis, SSc, myositis, MCTD and SS we did not find studies on this topic.

8. In patients with gout, we recommend a SUA level below 0.36 mmol/L (6 mg/dL) to potentially lower the risk of cardiovascular events and cardiovascular mortality. (LoE: 2b, GoR: C)

Retrospective cohort studies in patients with gout showed an association between an elevated SUA (per 0.06 mmol/L (1 mg/

Table 1 EULAR overarching principles and recommendations for the management of CVR in gout, vasculitis, SSc, myositis, MCTD, SS, SLE, and APS

Overarching principles	LoA* (SD)
A. Clinicians should be aware of increased CVR in patients with RMDs including gout, vasculitis, SSc, myositis, MCTD, SS, SLE and APS. For all RMDs, reduction of disease activity is likely to lessen CVR.	9.92 (0.39)
B. Rheumatologists are responsible for CVR assessment and management in collaboration with primary care providers, internists or cardiologists and other healthcare providers.	9.55 (1.12)
C. CVR factor screening should be performed regularly in all individuals with RMDs. Risk management should include screening for and strict control of CVR factors (smoking cessation, management of blood pressure, lipids and diabetes). CVR assessment is recommended within 6 months of diagnosis and repeated based on individual patient characteristics and risk levels.	9.55 (0.84)
D. Patient education and counselling on CVR, treatment adherence and lifestyle modifications, such as healthy diet and regular physical activity, are important in the management of CVR in these patients.	9.88 (0.42)
Recommendations for gout, vasculitis, SSc, myositis, MCTD and SS	
1. In patients with gout, vasculitis, SSc, myositis, MCTD and SS, we recommend thorough assessment of traditional CVR factors. The use of cardiovascular prediction tools for the general population is recommended. (LoE: 5, GoR†: D)	9.48 (0.84)
2. For ANCA-associated vasculitis the Framingham score may underestimate the CVR. Information from the EUVAS model may supplement modifiable Framingham risk factors and is recommended to take into account. (LoE: 2b, GoR: D)	8.59 (1.50)
3. In patients with gout, vasculitis, SSc, myositis, MCTD and SS, blood pressure management should follow recommendations used in the general population. (LoE: 5, GoR: D)	9.66 (0.62)
4. In patients with gout, diuretics should be avoided. (LoE: 5, GoR: D)	8.88 (2.06)
5. In patients with SSc beta blockers should be avoided. (LoE: 5, GoR: D)	8.92 (2.11)
6. In patients with gout, vasculitis, SSc, myositis, MCTD and SS, lipid management should follow recommendations used in the general population. (LoE: 5, GoR: D)	9.48 (1.08)
7. In patients with gout, vasculitis, SSc, myositis, MCTD and SS, standard use of platelet inhibitors for primary prevention is not recommended. Treatment with platelet inhibitors should follow recommendations used in the general population. (LoE: 2b/5, GoR: D)	9.37 (1.14)
8. In patients with gout, we recommend a serum uric acid level below 0.36 mmol/L (6 mg/dL) to potentially lower the risk on cardiovascular events and cardiovascular mortality. (LoE: 2b, GoR: C)	9.03 (1.34)
9. In patients with gout there is no preference for a particular urate-lowering therapy from the cardiovascular point of view. (LoE: 1b, GoR: B)	9.14 (1.35)
10. In patients with ANCA-associated vasculitis, remission induction and remission maintenance will also reduce CVR. (LoE: 2b, GoR: D)	9.07 (1.35)
11. In patients with giant-cell arteritis an optimal glucocorticoid regimen that balances the risk of relapse and glucocorticoid use side effects may also reduce CVR. (LoE: 2b, GoR: D)	9.14 (1.06)
Recommendations for SLE and the APS	
1. In patients with SLE and/or APS, a thorough assessment of traditional CVR factors and disease-related risk factors is recommended to guide risk factor modification. (LoE: 2b, GoR: D)	9.88 (0.32)
2A. In patients with SLE, lower levels of blood pressures are associated with lower rates of cardiovascular events and a blood pressure target of <130/80 mm Hg should be considered. (LoE: 2b, GoR: C)	9.70 (0.54)
2B. In patients with lupus nephritis, ACE inhibitors or angiotensin receptor blockers are recommended for all patients with urine protein-to-creatinine ratio >500 mg/g or arterial hypertension. (LoE: 5, GoR: D)	9.51 (0.64)
2C. In patients with APS, blood pressure management should follow recommendations used in the general population. (LoE: 5, GoR: D)	9.81 (0.39)
3. In patients with SLE and/or APS, lipid treatment should follow recommendations used in the general population. (LoE: 5, GoR: D)	9.70 (0.54)
4A. Patients with SLE may be candidates for preventative strategies as in the general population, including low-dose aspirin, based on their individual CVR profile. (LoE: 2b, GoR: D)	9.29 (1.37)
4B. In asymptomatic aPL carriers (not fulfilling any vascular or obstetric APS classification criteria) with a high-risk aPL profile with or without traditional risk factors, prophylactic treatment with low-dose aspirin (75–100 mg daily) is recommended. (LoE: 2a, GoR: B) In patients with SLE and no history of thrombosis or pregnancy complications: (1) with high-risk aPL profile, prophylactic treatment with low-dose aspirin is recommended (LoE: 2a, GoR: B); (2) with low-risk aPL profile, prophylactic treatment with low-dose aspirin may be considered. (LoE: 2b, GoR: C)	9.44 (0.97)
5. In patients with SLE, low disease activity should be maintained to also reduce CVR. (LoE: 2b, GoR: B)	9.59 (1.11)
6. In patients with SLE, treatment with the lowest possible corticosteroid dose is recommended to minimise any potential cardiovascular harm. (LoE: 2b, GoR: C)	9.59 (0.79)
7. In patients with SLE, no specific immunosuppressive medication can be recommended for the purpose of lowering the risk of cardiovascular events. (LoE: 2b, GoR: C)	9.44 (0.89)
8. In patients with SLE, treatment with hydroxychloroquine (which is recommended for all patients unless contraindicated) should be considered to also reduce the risk of cardiovascular events. (LoE: 2b, GoR: B)	9.66 (0.73)

*LoA, level of agreement; numbers in column indicate the mean (SD) of the LoA among task force members.

†LoE, level of evidence: 1a: systematic review of RCTs; 1b: individual RCT; 2a: systematic review of cohort studies; 2b: individual cohort study (and low-quality RCT); 3a: systematic review of case-control studies; 3b: individual case-control study; 4: case series and poor-quality cohort and case-control studies; 5: expert opinion without explicit critical appraisal, or based on physiology, bench research or 'first principles'.

#GoR, grade of recommendation: A: consistent level 1 studies; B: consistent level 2 or 3 studies, or extrapolations from level 1 studies; C: level 4 studies or extrapolations from level 2 or 3 studies; D: level 5 evidence or troublingly inconsistent or inconclusive studies of any level.

ACE, angiotensin-converting enzyme; ANCA, antineutrophil cytoplasmic antibodies; aPL, antiphospholipid antibodies; APS, antiphospholipid syndrome; CVR, cardiovascular risk; EULAR, European League against Rheumatism; EUVAS, European Vasculitis Society; MCTD, mixed connective tissue disease; RMDs, rheumatic and musculoskeletal diseases; SLE, systemic lupus erythematosus; SS, Sjögren's syndrome; SSc, systemic sclerosis.

dL)) and cardiovascular events.^{54–55} The association might be stronger in patients with SUA levels above 0.48 mmol/L (8 mg/dL),⁵⁶ than in patients with SUA levels higher than 0.36 mmol/L (6 mg/dL).⁵⁷ Studies on the effect of urate-lowering therapy

(ULT) showed conflicting results. Evidence originates predominantly from observational studies and often lacked data on treatment adherence and SUA levels during treatment. One study showed a linear dose response relation with a decline in the CVR

Recommendation

in the group with the highest defined daily dose.⁵⁸ This suggests that adequate ULT possibly lowers the CVR. This possibility was supported by two studies that showed a protective association of respectively 'high dose' allopurinol and ULT resulting in SUA <0.36 mmol/L (<6 mg/dL) on cardiovascular events and cardiovascular mortality.^{59 60} Altogether, although numbers of events were often low and associations were stronger for the highest SUA quartiles and higher dose ULT, it is possible that achieving lower SUA level decreases the risk on CV events. A cut-off value of 0.36 mmol/L (6 mg/dL) is used in the management of gout activity and could also benefit the risk of cardiovascular events. There is not sufficient evidence to support a threshold lower than 0.36 mmol/L (6 mg/dL) for CVR management.

9. In patients with gout there is no preference for a particular ULT from the cardiovascular point of view. (LoE: 1b, GoR: B)

Current guidelines recommend allopurinol as the first choice of ULT followed by febuxostat. Most studies on CVR compared these two xanthine oxidase inhibitors. Overall, regardless of the used dosage and duration of treatment, no difference was seen in number of cardiovascular events.^{61–63} In 2018, the CARES trial reported a higher risk of cardiovascular mortality with febuxostat than allopurinol.⁶² However, no difference was seen in the primary composite cardiovascular disease endpoint. Recently, the FAST trial showed no difference in CVR between patients using allopurinol or febuxostat.²⁹ Because of the limitations of the CARES trial (high number drop-outs, no difference in primary outcome, most events occurred after discontinuation of study) and the non-inferiority results of the FAST trial, we do not recommend the use of a specific ULT regarding cardiovascular outcomes.

Interventions targeting disease-related CVR factors

10. In patients with ANCA-associated vasculitis, remission induction and remission maintenance will also reduce CVR. (LoE: 2b, GoR: D)

In three of four included studies an association was found between high disease activity scores (Birmingham Vasculitis Activity Scores version 3) and a higher risk for cardiovascular events.^{64–66}

11. In patients with GCA an optimal glucocorticoid regimen that balances the risk of relapse and glucocorticoid use side effects may be considered to also reduce CVR. (LoE: 2b, GoR: D)

In patients with vasculitis, SSc, myositis, MCTD, and SS the primary goal is disease control with the lowest possible dose of glucocorticoids. In GCA two studies found a higher CVR in patients with a higher (daily/cumulative) prednisone dose. One study found that the use of an immunosuppressant in addition to glucocorticoid was a protective factor against new cardiovascular events.^{67 68} The increased CVR associated with glucocorticoids has to be balanced with the risk of relapse. Special attention and frequent evaluation of risks and benefits are warranted for patients with ongoing low dose glucocorticoids.

SLE and/or APS

CVR prediction tools

1. In patients with SLE and/or APS, a thorough assessment of traditional CVR factors and disease-related risk factors is recommended to guide risk factor modification. (LoE: 2b, GoR: D)

The FRS underestimates CVR in SLE patients^{18 69–71} with stroke, more often than myocardial infarction (MI), accounting for excess 'missed' risk by the FRS.^{69 70} A modified version of the FRS that used a 2.0 multiplier was found, retrospectively,

to improve the measure's sensitivity from 0.13 to 0.31 while maintaining good specificity to identify patients with a moderate/high risk of coronary artery disease.⁷² A study examining cardiovascular mortality in middle-aged patients with SLE found that SCORE predicted less than half the observed fatal cardiovascular events.⁷³ The QRISK3 tool included weights for SLE,³⁸ but validation studies in SLE populations have not yet been performed. Direct comparison of the performance of most commonly used generic risk assessment tools in SLE is currently lacking. A new SLE-specific risk score that included disease-related variables (SLEDAI, lupus anticoagulant and low C3) along with traditional risk factors found higher estimated risks than the American College of Cardiology/American Heart Association risk equation, except among patients whose risk was already moderate/high from traditional risk factors.⁷⁴ This prediction equation requires more testing and independent validation. Given the limitations of the current evidence, the task force did not endorse use of any particular CVR assessment tool, but instead recommended a thorough assessment of traditional and disease-related risk factors to guide cardiovascular prevention interventions.

No studies were identified that examined generic CVR prediction scores in APS. The adjusted Global APS Score (aGAPSS), a clinical score including the three major antiphospholipid antibodies (aPL), hypertension and lipidaemia, was developed to predict thrombosis, though data on cardiovascular events were not reported separately.⁷⁵ Modification of the aGAPSS by adding points for diabetes mellitus, smoking, and obesity to create a score specific for cardiovascular disease, the aGAPSS_{CVD} score, increased its discriminative ability and accuracy for CVR prediction in one study,⁷⁶ but further testing is needed.

Interventions targeting traditional CVR factors

2A. In patients with SLE, lower levels of BP are associated with lower rates of cardiovascular events and a BP target of <130/80 mm Hg should be considered. (LoE: 2b, GoR: C)

2B. In patients with lupus nephritis, ACEi or ARBs are recommended for all patients with urine protein-to-creatinine ratio >500 mg/g or arterial hypertension. (LoE: 5, GoR: D)

2C. In patients with APS, hypertension management should follow recommendations used in the general population. (LoE: 5, GoR: D)

A. SLE. Hypertension is associated with a higher risk of both coronary artery disease events⁷⁷ and first ischaemic stroke⁷⁸ in SLE. It, therefore, follows that BP control with antihypertensive medications should reduce the risk of cardiovascular events.⁷⁹ Recent mean systolic BP ≥ 132 mm Hg was identified as a determinant of a higher risk of cardiovascular events, and systolic BP had a stronger association than diastolic BP.⁸⁰ A recent study of patients with SLE examining three BP categories (normotensive; systolic BP 130–139/diastolic BP 80–89; systolic BP ≥ 140 /diastolic BP ≥ 90 mm Hg) reported an increased risk of cardiovascular events in both hypertensive groups compared with the normotensive group,³⁰ suggesting that a target BP of less than 130/80 should be used.

B. Lupus nephritis. Evidence specifically addressing the impact of antihypertensive treatment on cardiovascular events in lupus nephritis is scarce. In a retrospective cohort analysis,⁸¹ risk of a cardiovascular event was not associated with treatment with ACEI/ARB, but 18% in the ACEI/ARB group had end-stage renal disease compared with 2.4% in the comparison group and this imbalance would be expected to affect the comparison of CVRs. The panel endorsed the current EULAR/ERA-EDTA

recommendation on the use of ACEI/ARB for patients with lupus nephritis with concomitant hypertension or high-level proteinuria.³²

C. APS. No studies were identified on the use of specific anti-hypertensives for cardiovascular prevention in patients with APS. These patients should be managed according to recommendations for the general population.⁸²

3. *In patients with SLE and/or APS, hyperlipidaemia treatment should follow recommendations used in the general population.* (LoE: 5, GoR: D)

Higher levels of total cholesterol and low-density lipoprotein cholesterol have been associated with a higher risk of MI and stroke in SLE.^{74 78 83} One study using national administrative data found that patients with SLE treated with lipid-lowering agents had a significantly lower risk of coronary artery disease during follow-up (mean 8.4 years) than those not treated, while short-duration or long-duration statin use were both associated with a lower risk of stroke.⁸⁴ Several other observational studies included statin use as a covariate in prediction of cardiovascular events, and identified statin use as a risk factor for events, likely representing confounding by indication.^{71 85–88} Diagnosis of SLE is not sufficient per se for prescribing lipid-lowering treatment for primary cardiovascular prevention.⁸⁹ In APS, no study was identified that examined the effect of lipid-lowering agents on cardiovascular events. The task force judged that hyperlipidaemia treatment should follow the recommendations used in the general population.⁸⁹

4A. *Patients with SLE may be candidates for preventive strategies as in the general population, including low-dose aspirin, based on their individual CVR profile.* (LoE: 2b, GoR: D)

4B. *In asymptomatic aPL carriers with a high-risk profile with or without traditional risk factors, prophylactic treatment with low-dose aspirin (75–100 mg daily) is recommended.* (LoE: 2 a, GoR: B) *In patients with SLE and no history of thrombosis or pregnancy complications, prophylactic treatment with low-dose aspirin is recommended for those with a high-risk aPL profile* (LoE: 2a, GoR: B) *and may be considered for those with a low risk APL profile.* (LoE: 2b; GoR: C)

The panel agreed to include the corresponding statements (and LoE and GoR) about the prophylactic use of antiplatelets in SLE and APS from the recent EULAR recommendations for the management of SLE⁹⁰ and APS,⁹¹ respectively. The LoA from our task force group is shown in [table 1](#). Use of low-dose aspirin for cardiovascular prevention in patients with SLE or APS should be individualised (particularly in the presence of a high-risk aPL profile) according to EULAR recommendations.

Interventions targeting disease-related CVR factors

5. *In patients with SLE, low disease activity should be maintained to also reduce CVR.* (LoE: 2b, GoR: B)

SLE activity has often been reported as a predictor of cardiovascular events. With the exception of two studies,^{86 92} higher time-integrated SLEDAI levels were associated with an increased risk of cardiovascular events,^{69 77 79 93} more so than baseline or single measurements.^{78 94 95} In three studies,^{71 96 97} baseline SLEDAI was found to be higher in patients with cardiovascular events, although it was not carried to multivariable analysis. Associations of SLEDAI with cardiovascular events was found to be stronger when considering categories of activity compared with per-unit increases,⁶⁹ suggesting a non-linear association of disease activity with cardiovascular events.

Many studies did not consider simultaneously the association of measures of disease activity and SLE medication use;

therefore, results may be confounded. In an analysis that adjusted for current prednisone dose, a 1-point increase in SLEDAI was marginally associated with an increased risk of cardiovascular events (relative risk 1.05, 95% CI 1.00 to 1.11).⁶⁹ Available evidence indicates that higher disease activity may be associated with a higher risk of cardiovascular events. Thus, in addition to its importance in general patient management,⁹⁰ a low-disease activity state may also have a beneficial effect on cardiovascular health.

6. *In patients with SLE, treatment with the lowest possible glucocorticoid dose is recommended to minimise any potential cardiovascular harm.* (LoE: 2b, GoR: C)

Mean dosage, cumulative exposure and duration of glucocorticoid treatment have all been investigated with reference to cardiovascular events in SLE. Higher current glucocorticoid dose was associated with a higher risk of atherothrombotic events, ischaemic heart disease, and/or stroke in two studies,^{69 98} but was protective in one study⁷⁹ and not associated with stroke in the SLICC inception cohort.⁹⁹ Higher mean daily doses, greater cumulative doses, and ever-use of prednisone 30 mg/day or more were more consistently associated with increased risks of cardiovascular events in both cohort and case-control studies,^{71 92 100 101} although glucocorticoid use was not significantly associated with cardiovascular events in two analyses of the Toronto cohort.^{95 97} Not all studies adjusted for SLE activity. A retrospective study that adjusted for SLE activity⁹⁸ found that higher daily doses (prednisone >10 mg) administered continuously were significantly associated with both MI and stroke. In a retrospective and non-randomised study, patients treated at clinics following a glucocorticoid dose-minimisation strategy had lower prednisone exposures and markedly lower risks of cardiovascular damage by the SLICC measure, particularly for stroke.¹⁰² Most evidence suggests that higher glucocorticoid exposure (cumulative and mean daily dose) increases CVR in SLE. The task force recommended treatment with the lowest possible corticosteroid dose to minimise risks of cardiovascular harm.

7. *In patients with SLE, no specific immunosuppressive medication can be recommended for the purpose of lowering the risk of cardiovascular events.* (LoE: 2b, GoR: C)

Use of immunosuppressants as a class in SLE have had largely null or conflicting associations with cardiovascular events.^{79 99 103} Three studies from the Toronto lupus cohort reported either a protective⁹⁶ or null association,^{93 97} while one study found that patients treated with immunosuppressants vs those not treated were more likely to develop a cardiovascular event in univariate but not multivariate analyses.⁹⁵ Immunosuppressive therapy was also associated to higher odds of ischaemic heart disease and cardiovascular events in the LUMINA¹⁰⁴ and Hopkins lupus cohort.⁶⁹

Studies of individual medications suggest that use of methotrexate, mycophenolate, cyclosporine, or rituximab had neutral associations with cardiovascular events.^{88 92 105} Conflicting results have been reported for cyclophosphamide^{71 106} and azathioprine.^{71 88 106}

A common limitation in many studies was the examination of ever use vs never use of immunosuppressants, which may be too crude an exposure. No studies considered issues of confounding by indication, and positive associations with cardiovascular disease may reflect risks due to associated disease activity or severity, or concomitant glucocorticoid use. Based on current evidence, the task force concluded that no specific immunosuppressive medication can be recommended for reducing the risk of cardiovascular events. Furthermore, the committee call for better quality pharmacoepidemiologic studies in future, using recent advances in this field.

Recommendation

8. In patients with SLE, treatment with hydroxychloroquine (which is recommended for all SLE patients, unless contraindicated) should be considered to also reduce the risk of cardiovascular events. (LoE: 2b, GoR: B)

A large body of evidence has addressed the role of antimalarials in cardiovascular prevention in SLE. In six cohort studies, antimalarial use was associated with lower risk of either atherothrombotic events or coronary artery disease,^{69 77 79 88 94 107} although in one study protection was only associated with current long-term use.⁶⁹ Several other studies reported null associations.^{85 87 92 93 95 103 106} Two of seven case-control studies also reported less use of hydroxychloroquine or antimalarials among cases with cardiovascular events than controls,^{100 108} with only one study reporting increased risk.⁹⁷ No associations with risk of stroke specifically have been reported.^{99 109} Importantly, patients with less active disease are more often treated with antimalarials, while SLE activity may be the risk factor for cardiovascular disease; this possible selection bias was not addressed. Additionally, studies did not report results stratified by the presence of APS or aPL, therefore, it is unclear if any reduced risk is limited to patients with SLE and aPL. The task force endorsed treatment with hydroxychloroquine, as should be provided to all patients with SLE, as it may also reduce the risk of cardiovascular events.

DISCUSSION

The 2021 EULAR recommendations for CVR management in RMDs comprise overarching principles and guidance informed by the currently available evidence on several potential interventions aiming to improve cardiovascular outcomes in these disorders. The LoA for most statements was high, indicating a coherent perspective on behalf of health professionals from different areas of care and patients alike for CVR reduction efforts.

The majority of the included RMDs are uncommon diseases limiting the ability to perform large observational studies to assess the impact of traditional and disease-specific risk factors on cardiovascular disease burden and clinical trials on the long-term cardiovascular effects of preventive treatments. One of the main challenges of these recommendations was the low LoE due to few studies on many of the research questions. Confounding by indication and lack of propensity adjustment was a common limitation in the included studies and therefore several statements relied on expert opinion. Future studies that better identify exposures and outcomes may help overcome these methodological issues.

There are several additional issues that need to be addressed in the future efforts for CVR management in RMDs. Systemic RMDs are complex diseases with a wide range of clinical manifestations of various severity that may affect cardiovascular health in diverse ways. Considering personalised patient care, the potential impact of individual patient clinical phenotype on cardiovascular prognosis also merits further investigation. In guidelines for cardiovascular prevention in the general population, risk stratification represents a prerequisite for CVR management (eg, BP targets or lipid-lowering therapy).^{82 89} In this context, it is important to recognise that underperformance of clinical CVR prediction tools used in the general population may hamper CVR prevention and management in RMDs. The use of prediction tools that incorporate CRP¹¹⁰ (eg, Reynolds risk score¹¹¹), the presence of specific RMDs (RA, SLE) or anti-inflammatory agents (eg, QRISK3)³⁸ or multipliers of baseline risk (eg, modified SCORE)¹¹² has been suggested by some guideline committees for CVR stratification in the general population but their use in RMDs needs to be further tested and validated. Thus, studies on disease-specific tools for CVR assessment including disease-specific in addition to traditional CVR factors, as well as

Box 1 Research agenda and future perspectives

1. Validation of existing generic and modified CVR prediction tools in large prospective studies, and development of new disease-specific equations.
2. Additive value of vascular imaging and/or circulating biomarkers in CVR assessment in RMDs.
3. Identification of patient subgroups with higher CVR.
4. Long-term effects of current and new drugs for RMDs on CVR factors and cardiovascular events.
5. Role of antithrombotic agents used in some RMDs (eg, aspirin, LMWH in SLE/APS) to reduce the overall CVR in these patients.
6. Need for large educational campaigns within the rheumatological and other medical specialties and patient associations to increase CVR awareness.
7. Best implementation methods for the CVR recommendations. APS, antiphospholipid syndrome; CVR, cardiovascular risk; LMWH, low-molecular weight heparin; RMDs, rheumatic and musculoskeletal diseases; SLE, systemic lupus erythematosus.

risk qualifiers including the evaluation of the predictive value of nonclinical tools, are warranted. These issues, along with other relevant questions such as the pragmatic use of any risk score (simplicity often aids use) will hopefully inspire future research increasing the quality of evidence in CVR management in RMDs, are presented in the Research Agenda (box 1). One of future challenges is the better identification of patient subgroups at higher CVR including for example those with longer disease duration, and number of flares/relapses (eg, in SLE, vasculitis, gout)^{55 66 113–115} or those with certain demographic (age, gender, race/ethnicity)¹¹⁶ and disease characteristics (eg, aPL positivity in SLE, polyarticular or tophaceous phenotype in gout).^{55 113 117}

Long-term effects of current and new drugs for RMDs on CVR need further investigation. The deleterious cardiometabolic effects of the excessive exposure to glucocorticoids are well known.¹¹⁸ Current recommendations by the ACR¹¹⁹ and the EULAR^{53 90 120 121} for the management of RMDs emphasise the adverse effects and the need of the limited dose of glucocorticoids. Limiting glucocorticoid exposure to the lowest effective dose to control active disease for the shortest duration possible and eventually discontinuation, as well as weighting the benefits and risks before starting systemic glucocorticoids, can help reduce cardiovascular harm. Several anti-inflammatory agents (eg, colchicine,¹²² anti-IL1b¹²³) have been shown to lower cardiovascular outcomes in randomised controlled trials for secondary prevention of cardiovascular disease in the general population and other trials are ongoing (eg, hydroxychloroquine¹²⁴) but further evidence is needed on the cardiovascular outcomes and safety of such immunoregulatory agents in RMDs. Although the role of hydroxychloroquine in APS, and of non-steroidal anti-inflammatory drugs (NSAIDs) in SLE, was examined in our SLR (Supplementary SLR report, section II), the panel agreed that any statement on the use of these medications should be deferred until more robust evidence is available. More evidence is needed about the effect of glucocorticoids, NSAIDs and IL-1 antagonists, the dosage and duration of colchicine treatment, and the risk and benefits of the concomitant use of colchicine and statins in patients with gout.

Most of the recommendations of established low-cost clinical interventions may apply to both high-resource and low-resource countries worldwide. Implementation strategies for promoting

CVR management in RMDs include interactive educational workshops involving health professionals, patients and stakeholders with the support of healthcare professional societies and patient associations, social media dissemination and strategies customised to local and national policies such as academic detailing, audits and feedback techniques.

The panel believes that these recommendations will enable healthcare providers and patients to mutually engage in a long-term care pathway tailored to patients' needs and expectations for improving cardiovascular health in RMDs. As new data accumulate, this first set of 'best available' evidence on cardiovascular prevention in gout, vasculitis, SSc, myositis, MCTD, SS, SLE and APS will be timely updated.

Author affiliations

¹First Department of Propaedeutic Internal Medicine, Laiko Hospital, National and Kapodistrian University of Athens, Athens, Greece

²Reade, Amsterdam Rheumatology and Immunology Center, Amsterdam, The Netherlands

³Department of Internal Medicine, Northwest Clinics, Alkmaar, The Netherlands

⁴Rheumatology Unit, Department of Internal Medicine, University of Messina, Messina, Italy

⁵EULAR Patient Research Partner, Brussels, Belgium

⁶4th Department of Internal Medicine, "Attikon" University Hospital, Athens, Greece

⁷Joint Academic Rheumatology Program, Medical School, National and Kapodistrian University of Athens, Athens, Greece

⁸Division of Physiotherapy, Department of Neurobiology, Care Sciences and Society, Karolinska Institute, Huddinge, Sweden

⁹Department of Orthopaedics, Danderyd Hospital Corp, Stockholm, Sweden

¹⁰Centre for Epidemiology Versus Arthritis, Centre for Musculoskeletal Research, University of Manchester, Manchester, UK

¹¹NIHR Manchester Biomedical Research Centre, Manchester University Hospitals NHS Trust, Manchester Academic Health Science Centre, Manchester, UK

¹²Rheumatology Division, Hospital Universitario Marqués de Valdecilla and University of Cantabria, Santander, Spain

¹³Copenhagen Lupus and Vasculitis Clinic, Rigshospitalet, Copenhagen, Denmark

¹⁴Department of Clinical Medicine, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark

¹⁵Intensive Care Unit, Department of Medicine, University of Debrecen, Debrecen, Hungary

¹⁶EULAR Patient Research Partner, Rome, Italy

¹⁷Rheumatology Department, Norfolk and Norwich University Hospital, Colney Lane, UK

¹⁸Department of Autoimmune Diseases, ICMiD, University of Barcelona, Hospital Clínic, Barcelona, Spain

¹⁹Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK

²⁰EMEUNET member, Danish Hospital for Rheumatic Diseases, Sønderburg, Denmark

²¹EMEUNET member, CMiD-Nephrology, San Giovanni Bosco Hospital, University of Torino, Torino, Italy

²²Department of Medicine, Rheumatology Unit, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden

²³Department of Rheumatology, University of Debrecen Faculty of Medicine, Debrecen, Hungary

²⁴Department of Rheumatology, University Clinic Carl Gustav Carus at the TU Dresden, Dresden, Germany

²⁵Department of Rheumatology, University of Basel, Basel, Switzerland

²⁶Department of Cardiology, Amsterdam University Medical Center, location VU University medical center, Amsterdam, The Netherlands

²⁷Department of Rheumatology and Clinical Immunology, Amsterdam UMC, Vrije Universiteit, Amsterdam, The Netherlands

²⁸Department of Public Health, NHS Grampian, Aberdeen, UK

²⁹Intramural Research Program, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, Maryland, USA

³⁰Amsterdam University Medical Center, location VU University Medical Centre, Amsterdam, The Netherlands

Twitter Chetan Mukhtyar @cmukhtyar, Gary J Macfarlane @UAberdeenEpi and Maria G Tektonidou @mariatektonidou

Acknowledgements The committee wishes to acknowledge the support of the EULAR Standing Committee on Clinical Affairs and the EULAR Secretariat.

Contributors GCD, DV and EH drafted the manuscript. MN and MGT convened and supervised the project and edited the manuscript. GCD, DV, EH and LB performed the systematic literature review under the supervision of the

methodologists and convenors. GJM and MMW supervised the methodology of the systematic literature review and the Delphi process. All authors contributed with comments and revisions and approved the final version.

Funding This study was funded by European Alliance of Associations for Rheumatology, EULAR. Project number: CL112.

Competing interests GCD, DV, EH, LB, SB, DTB, NB, GK, FM, CM, MR-C, KS, SS, VPvH, GJM, MMW and MN have nothing to declare. FA: research grants from BMS, Celgene, Novartis and Sandoz and consulting fees from AbbVie, Biogen, BMS, Celgene, Janssen, Lilly, Novartis, Pfizer and Sanofi-Aventis, all unrelated to this manuscript; INB: grant from GSK paid to institution, consulting fees from Astra Zeneca, GSK, Eli Lilly, UC, MSD paid to institution, support for attending meetings and/or travel from GSK, participation on a data safety monitoring board/or advisory board from Aurinia, Astra Zeneca and ILTOO paid to institution, all unrelated to this manuscript; MAG-G: grants/research support from AbbVie, MSD, Jansen and Roche paid to institution and personal consulting fees/participation in company sponsored speakers bureau from AbbVie, Pfizer, Roche, Celgene, MSD, Novartis, SOBI and Sanofi, all unrelated to this manuscript; SJ: grants from BMS paid to institution, personal consulting fees from Astra Zeneca, and personal fees from Danish Medicolegal Council, all unrelated to this manuscript; NS: grants paid to institution from Astrazeneca, Boehringer Ingelheim and Roche Diagnostics, personal consulting fees from Afimmune, Amgen, Astrazeneca, Boehringer Ingelheim, Eli-Lilly, Hanmi Pharmaceuticals, MSD, Novartis, Novo Nordisk, Pfizer and Sanofi, all unrelated to this manuscript; ES: grant from Merck and honoraria from Janssen, all unrelated to this manuscript; ZS: research grants from Pfizer paid to institution and personal consulting fees from Pfizer, MSD, Lilly, Novartis, Roche, Gedeon Richter, Boehringer Ingelheim, Abbvie, all unrelated to this manuscript; A-KT: speakers fee from Berlin Chemie Menarini, Novartis and personal fees and non-financial support from AstraZeneca and Grünenthal, all unrelated to this manuscript; AT: consulting fees from Magenta Therapeutics and personal fees from Novartis and Idorsia for participation on a Data Safety Monitoring Board or Advisory Board, all unrelated to this manuscript; AV: personal consulting fees from Astra Zeneca and GS, all unrelated to this manuscript; MGT: research grants from Genesis, GSK, MSD, Pfizer and UCB paid to institution, and personal consulting fees from Genesis, GSK and Novartis, all unrelated to this manuscript.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

George C Drosos <http://orcid.org/0000-0001-7987-9588>

Daisy Vedder <http://orcid.org/0000-0003-1715-9555>

Laura Boekel <http://orcid.org/0000-0001-5473-7786>

Dimitrios T Boumpas <http://orcid.org/0000-0002-9812-4671>

Miguel Ángel González-Gay <http://orcid.org/0000-0002-7924-7406>

Søren Jacobsen <http://orcid.org/0000-0002-5654-4993>

Chetan Mukhtyar <http://orcid.org/0000-0002-9771-6667>

Savino Sciascia <http://orcid.org/0000-0003-1266-9441>

Elisabet Svenungsson <http://orcid.org/0000-0003-3396-3244>

Zoltan Szekanecz <http://orcid.org/0000-0002-7794-6844>

Gary J Macfarlane <http://orcid.org/0000-0003-2322-3314>

Michael M Ward <http://orcid.org/0000-0003-1857-9367>

Maria G Tektonidou <http://orcid.org/0000-0003-2238-0975>

REFERENCES

- England BR, Thiele GM, Anderson DR, *et al*. Increased cardiovascular risk in rheumatoid arthritis: mechanisms and implications. *BMJ* 2018;361:k1036.
- Agca R, Heslinga SC, Rollefstad S, *et al*. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis* 2017;76:17–28.
- Clarson LE, Chandratte P, Hider SL, *et al*. Increased cardiovascular mortality associated with gout: a systematic review and meta-analysis. *Eur J Prev Cardiol* 2015;22:335–43.
- Kuo C-F, Yu K-H, See L-C, *et al*. Risk of myocardial infarction among patients with gout: a nationwide population-based study. *Rheumatology* 2013;52:111–7.
- Alenghat FJ. The prevalence of atherosclerosis in those with inflammatory connective tissue disease by race, age and traditional risk factors. *Sci Rep* 2016;6:20303.

- 6 Aouba A, Gonzalez Chiappe S, Eb M, *et al.* Mortality causes and trends associated with giant cell arteritis: analysis of the French national death certificate database (1980–2011). *Rheumatology* 2018;57:1047–55.
- 7 Houben E, Penne EL, Voskuyl AE, *et al.* Cardiovascular events in anti-neutrophil cytoplasmic antibody-associated vasculitis: a meta-analysis of observational studies. *Rheumatology* 2018;57:555–62.
- 8 Zöller B, Li X, Sundquist J, *et al.* Risk of subsequent ischemic and hemorrhagic stroke in patients hospitalized for immune-mediated diseases: a nationwide follow-up study from Sweden. *BMC Neurol* 2012;12:41.
- 9 Zöller B, Li X, Sundquist J, *et al.* Risk of subsequent coronary heart disease in patients hospitalized for immune-mediated diseases: a nationwide follow-up study from Sweden. *PLoS One* 2012;7:e33442.
- 10 Tektonidou MG, Lewandowski LB, Hu J, *et al.* Survival in adults and children with systemic lupus erythematosus: a systematic review and Bayesian meta-analysis of studies from 1950 to 2016. *Ann Rheum Dis* 2017;76:2009–16.
- 11 Cervera R, Serrano R, Pons-Estel GJ, *et al.* Morbidity and mortality in the antiphospholipid syndrome during a 10-year period: a multicentre prospective study of 1000 patients. *Ann Rheum Dis* 2015;74:1011–8.
- 12 Houben E, Mendel A, van der Heijden JW, *et al.* Prevalence and management of cardiovascular risk factors in ANCA-associated vasculitis. *Rheumatology* 2019;58:2333–5.
- 13 Cen X, Feng S, Wei S, *et al.* Systemic sclerosis and risk of cardiovascular disease: a PRISMA-compliant systematic review and meta-analysis of cohort studies. *Medicine* 2020;99:e23009.
- 14 Choi HK, Curhan G. Independent impact of gout on mortality and risk for coronary heart disease. *Circulation* 2007;116:894–900.
- 15 Clarson LE, Hider SL, Belcher J, *et al.* Increased risk of vascular disease associated with gout: a retrospective, matched cohort study in the UK clinical practice research Datalink. *Ann Rheum Dis* 2015;74:642–7.
- 16 Kurmann RD, Sandhu AS, Crowson CS, *et al.* Cardiovascular risk factors and atherosclerotic cardiovascular events among incident cases of systemic sclerosis: results from a population-based cohort (1980–2016). *Mayo Clin Proc* 2020;95:1369–78.
- 17 Geovanini GR, Libby P. Atherosclerosis and inflammation: overview and updates. *Clin Sci* 2018;132:1234–52.
- 18 Esdaile JM, Abrahamowicz M, Grodzicky T, *et al.* Traditional Framingham risk factors fail to fully account for accelerated atherosclerosis in systemic lupus erythematosus. *Arthritis Rheum* 2001;44:2331–7.
- 19 Manzi S, Wasko MC. Inflammation-Mediated rheumatic diseases and atherosclerosis. *Ann Rheum Dis* 2000;59:321–5.
- 20 Ridker PM, Hennekens CH, Buring JE, *et al.* C-Reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000;342:836–43.
- 21 The Emerging Risk Factors Collaboration, Kaptoge S, Di Angelantonio E, *et al.* C-Reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med* 2012;367:1310–20.
- 22 Zhao TX, Mallat Z. Targeting the immune system in atherosclerosis: JACC state-of-the-art review. *J Am Coll Cardiol* 2019;73:1691–706.
- 23 Ajala ON, Everett BM. Targeting inflammation to reduce residual cardiovascular risk. *Curr Atheroscler Rep* 2020;22:66.
- 24 Lawler PR, Bhatt DL, Godoy LC, *et al.* Targeting cardiovascular inflammation: next steps in clinical translation. *Eur Heart J* 2021;42:113–31.
- 25 Libby P. The changing landscape of atherosclerosis. *Nature* 2021;592:524–33.
- 26 van der Heijde D, Aletaha D, Carmona L, *et al.* 2014 update of the EULAR standardised operating procedures for EULAR-endorsed recommendations. *Ann Rheum Dis* 2015;74:8–13.
- 27 Brouwers MC, Kho ME, Brouman GP, *et al.* AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ* 2010;182:E839–42.
- 28 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
- 29 Mackenzie IS, Ford I, Nuki G, *et al.* Long-term cardiovascular safety of febuxostat compared with allopurinol in patients with gout (FAST): a multicentre, prospective, randomised, open-label, non-inferiority trial. *Lancet* 2020;396:1745–57.
- 30 Tselios K, Gladman DD, Su J, *et al.* Impact of the new American College of Cardiology/American Heart Association definition of hypertension on atherosclerotic vascular events in systemic lupus erythematosus. *Ann Rheum Dis* 2020;79:612–7.
- 31 Kravvariti E, Koutsogianni A, Samoli E, *et al.* The effect of hydroxychloroquine on thrombosis prevention and antiphospholipid antibody levels in primary antiphospholipid syndrome: a pilot open label randomized prospective study. *Autoimmun Rev* 2020;19:102491.
- 32 Fanouriakis A, Kostopoulou M, Cheema K, *et al.* 2019 update of the joint European League against rheumatism and European renal Association-European dialysis and transplant association (EULAR/ERA-EDTA) recommendations for the management of lupus nephritis. *Ann Rheum Dis* 2020;79:713–23.
- 33 Higgins JPT, Altman DG, Gotzsche PC, *et al.* The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- 34 Deeks JJ, Dinnes J, D'Amico R, *et al.* Evaluating non-randomised intervention studies. *Health Technol Assess* 2003;7;.:1–173. iii-x.
- 35 CEBM. Oxford centre for evidence-based Medicine—Levels of evidence (March 2009), 2009. Available: <https://www.cebm.ox.ac.uk/resources/levels-of-evidence/oxford-centre-for-evidence-based-medicine-levels-of-evidence-march-2009> [Accessed October 2021].
- 36 D'Agostino RB, Vasan RS, Pencina MJ, *et al.* General cardiovascular risk profile for use in primary care: the Framingham heart study. *Circulation* 2008;117:743–53.
- 37 Conroy RM, Pyörälä K, Fitzgerald AP, *et al.* Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;24:987–1003.
- 38 Hippisley-Cox J, Coupland C, Brindle P. Development and validation of QRISK3 risk prediction algorithms to estimate future risk of cardiovascular disease: prospective cohort study. *BMJ* 2017;357:j2099.
- 39 Berti A, Matteson EL, Crowson CS, *et al.* Risk of cardiovascular disease and venous thromboembolism among patients with incident ANCA-associated vasculitis: a 20-year population-based cohort study. *Mayo Clin Proc* 2018;93:597–606.
- 40 Terrier B, Chironi G, Pagnoux C, *et al.* Factors associated with major cardiovascular events in patients with systemic necrotizing vasculitides: results of a longterm followup study. *J Rheumatol* 2014;41:723–9.
- 41 Suppiah R, Judge A, Batra R, *et al.* A model to predict cardiovascular events in patients with newly diagnosed Wegener's granulomatosis and microscopic polyangiitis. *Arthritis Care Res* 2011;63:588–96.
- 42 Grossman C, Barshack I, Koren-Morag N, *et al.* Risk factors for severe cranial ischaemic events in patients with giant cell arteritis. *Clin Exp Rheumatol* 2017;35 Suppl 103:88–93.
- 43 Valentini G, Huscher D, Riccardi A, *et al.* Vasodilators and low-dose acetylsalicylic acid are associated with a lower incidence of distinct primary myocardial disease manifestations in systemic sclerosis: results of the DeSicipher inception cohort study. *Ann Rheum Dis* 2019;78:1576–82.
- 44 Richette P, Doherty M, Pascual E, *et al.* 2016 updated EULAR evidence-based recommendations for the management of gout. *Ann Rheum Dis* 2017;76:29–42.
- 45 Garcia-Gil M, Comas-Cufi M, Ramos R, *et al.* Effectiveness of statins as primary prevention in people with gout: a population-based cohort study. *J Cardiovasc Pharmacol Ther* 2019;24:542–50.
- 46 Keller SF, Rai SK, Lu N, *et al.* Statin use and mortality in gout: a general population-based cohort study. *Semin Arthritis Rheum* 2018;48:449–55.
- 47 Finkelstein Y, Aks SE, Hutson JR, *et al.* Colchicine poisoning: the dark side of an ancient drug. *Clin Toxicol* 2010;48:407–14.
- 48 Narváez J, Bernad B, Nolla JM, *et al.* Statin therapy does not seem to benefit giant cell arteritis. *Semin Arthritis Rheum* 2007;36:322–7.
- 49 Pariente A, Guéron A, Alamowitch S, *et al.* Ischemic stroke in giant-cell arteritis: French retrospective study. *J Autoimmun* 2019;99:48–51.
- 50 Pugnet G, Sailler L, Fournier J-P, *et al.* Predictors of cardiovascular hospitalization in giant cell arteritis: effect of statin exposure. A French population-based study. *J Rheumatol* 2016;43:2162–70.
- 51 Mukhtyar C, Guillevin L, Cid MC, *et al.* EULAR recommendations for the management of large vessel vasculitis. *Ann Rheum Dis* 2009;68:318–23.
- 52 Maz M, Chung SA, Abril A, *et al.* 2021 American College of Rheumatology/Vasculitis Foundation guideline for the management of giant cell arteritis and Takayasu arteritis. *Arthritis Rheumatol* 2021;73:1349–65.
- 53 Hellmich B, Agüeda A, Monti S, *et al.* 2018 update of the EULAR recommendations for the management of large vessel vasculitis. *Ann Rheum Dis* 2020;79:19–30.
- 54 Chen J-H, Lan J-L, Cheng C-F, *et al.* Effect of urate-lowering therapy on all-cause and cardiovascular mortality in hyperuricemic patients without gout: a case-matched cohort study. *PLoS One* 2015;10:e0145193.
- 55 Perez-Ruiz F, Martínez-Indart L, Carmona L, *et al.* Topheaceous gout and high level of hyperuricaemia are both associated with increased risk of mortality in patients with gout. *Ann Rheum Dis* 2014;73:177–82.
- 56 Essex MN, Hopps M, Bienen EJ, *et al.* Evaluation of the relationship between serum uric acid levels and cardiovascular events in patients with gout: a retrospective analysis using electronic medical record data. *J Clin Rheumatol* 2017;23:160–6.
- 57 Stack AG, Hanley A, Casserly LF, *et al.* Independent and conjoint associations of gout and hyperuricaemia with total and cardiovascular mortality. *QJM* 2013;106:647–58.
- 58 Lin H-C, Daimon M, Wang C-H, *et al.* Allopurinol, benzbromarone and risk of coronary heart disease in gout patients: a population-based study. *Int J Cardiol* 2017;233:85–90.
- 59 Joo K, Kwon S-R, Lim M-J, *et al.* Prevention of comorbidity and acute attack of gout by uric acid lowering therapy. *J Korean Med Sci* 2014;29:657–61.
- 60 Kok VC, Horng J-T, Chang W-S, *et al.* Allopurinol therapy in gout patients does not associate with beneficial cardiovascular outcomes: a population-based matched-cohort study. *PLoS One* 2014;9:e99102.
- 61 Kang EH, Choi HK, Shin A, *et al.* Comparative cardiovascular risk of allopurinol versus febuxostat in patients with gout: a nation-wide cohort study. *Rheumatology* 2019;58:2122–9.
- 62 White WB, Saag KG, Becker MA, *et al.* Cardiovascular safety of febuxostat or allopurinol in patients with gout. *N Engl J Med* 2018;378:1200–10.
- 63 Zhang M, Solomon DH, Desai RJ, *et al.* Assessment of cardiovascular risk in older patients with gout initiating febuxostat versus allopurinol: population-based cohort study. *Circulation* 2018;138:1116–26.

- 64 Houben E, Mendel A, Carette S, *et al.* Predictors of fatal and non-fatal cardiovascular events in ANCA-associated vasculitis: data from the Toronto CanVasc cohort. *Joint Bone Spine* 2020;87:30006–3.
- 65 Bai YH, ZY L, Chang DY. The BVAS is an independent predictor of cardiovascular events and cardiovascular disease-related mortality in patients with ANCA-associated vasculitis: a study of 504 cases in a single Chinese center. *Semin Arthritis Rheum* 2018;47:524–9.
- 66 Robson J, Doll H, Suppiah R, *et al.* Damage in the ANCA-associated vasculitides: long-term data from the European Vasculitis Study Group (EUVAS) therapeutic trials. *Ann Rheum Dis* 2015;74:177–84.
- 67 de Boissson H, Liozon E, Espitia O, *et al.* Different patterns and specific outcomes of large-vessel involvements in giant cell arteritis. *J Autoimmun* 2019;103:102283–83.
- 68 Gale S, Wilson JC, Chia J, *et al.* Risk associated with cumulative oral glucocorticoid use in patients with giant cell arteritis in real-world databases from the USA and UK. *Rheumatol Ther* 2018;5:327–40.
- 69 Magder LS, Petri M. Incidence of and risk factors for adverse cardiovascular events among patients with systemic lupus erythematosus. *Am J Epidemiol* 2012;176:708–19.
- 70 Bessant R, Hingorani A, Patel L, *et al.* Risk of coronary heart disease and stroke in a large British cohort of patients with systemic lupus erythematosus. *Rheumatology* 2004;43:924–9.
- 71 Haque S, Skeoch S, Rakieh C, *et al.* Progression of subclinical and clinical cardiovascular disease in a UK SLE cohort: the role of classic and SLE-related factors. *Lupus Sci Med* 2018;5:e000267.
- 72 Urowitz MB, Ibañez D, Su J, *et al.* Modified Framingham risk factor score for systemic lupus erythematosus. *J Rheumatol* 2016;43:875–9.
- 73 Gustafsson JT, Simard JF, Gunnarsson I, *et al.* Risk factors for cardiovascular mortality in patients with systemic lupus erythematosus, a prospective cohort study. *Arthritis Res Ther* 2012;14:R46.
- 74 Petri MA, Barr E, Magder LS. Development of a systemic lupus erythematosus cardiovascular risk equation. *Lupus Sci Med* 2019;6:e000346.
- 75 Radin M, Sciascia S, Erkan D, *et al.* The adjusted global antiphospholipid syndrome score (aGAPSS) and the risk of recurrent thrombosis: results from the APS action cohort. *Semin Arthritis Rheum* 2019;49:464–8.
- 76 Di Minno MND, Scalera A, Tufano A, *et al.* The association of adjusted global antiphospholipid syndrome score (aGAPSS) with cardiovascular disease in subjects with antiphospholipid antibodies. *Atherosclerosis* 2018;278:60–5.
- 77 Nikpour M, Urowitz MB, Ibanez D, *et al.* Importance of cumulative exposure to elevated cholesterol and blood pressure in development of atherosclerotic coronary artery disease in systemic lupus erythematosus: a prospective proof-of-concept cohort study. *Arthritis Res Ther* 2011;13:R156.
- 78 Mikdashi J, Handwerker B, Langenberg P, *et al.* Baseline disease activity, hyperlipidemia, and hypertension are predictive factors for ischemic stroke and stroke severity in systemic lupus erythematosus. *Stroke* 2007;38:281–5.
- 79 Becker-Merok A, Nossent J, Prevalence NJC. Prevalence, predictors and outcome of vascular damage in systemic lupus erythematosus. *Lupus* 2009;18:508–15.
- 80 Stojan G, Magder LS, Petri M. Blood pressure variability and age-related blood pressure patterns in systemic lupus erythematosus. *J Rheumatol* 2020;47:387–93.
- 81 Tselios K, Gladman DD, Su J, *et al.* Does renin-angiotensin system blockade protect lupus nephritis patients from atherosclerotic cardiovascular events? A case-control study. *Arthritis Care Res* 2016;68:1497–504.
- 82 Williams B, Mancia G, Spiering W, *et al.* 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J* 2018;39:3021–104.
- 83 Nikpour M, Gladman DD, Ibanez D, *et al.* Assessment of coronary risk based on cumulative exposure to lipids in systemic lupus erythematosus. *J Rheumatol* 2013;40:2006–14.
- 84 Yu H-H, Chen P-C, Yang Y-H, *et al.* Statin reduces mortality and morbidity in systemic lupus erythematosus patients with hyperlipidemia: a nationwide population-based cohort study. *Atherosclerosis* 2015;243:11–18.
- 85 Kao AH, Lertratanakul A, Elliott JR, *et al.* Relation of carotid intima-media thickness and plaque with incident cardiovascular events in women with systemic lupus erythematosus. *Am J Cardiol* 2013;112:1025–32.
- 86 Iudici M, Fasano S, Gabriele Falcone L, *et al.* Low-Dose aspirin as primary prophylaxis for cardiovascular events in systemic lupus erythematosus: a long-term retrospective cohort study. *Rheumatology* 2016;55:1623–30.
- 87 Pons-Estel GJ, González LA, Zhang J, *et al.* Predictors of cardiovascular damage in patients with systemic lupus erythematosus: data from LUMINA (LXVIII), a multiethnic US cohort. *Rheumatology* 2009;48:817–22.
- 88 Fernández-Nebro A, Rúa-Figueroa Íñigo, López-Longo FJ, *et al.* Cardiovascular events in systemic lupus erythematosus: a nationwide study in Spain from the RELESSER registry. *Medicine* 2015;94:e1183.
- 89 Mach F, Baigent C, Catapano AL, *et al.* 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020;41:111–88.
- 90 Fanouriakis A, Kostopoulou M, Alunno A, *et al.* 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:736–45.
- 91 Tektonidou MG, Andreoli L, Limper M, *et al.* EULAR recommendations for the management of antiphospholipid syndrome in adults. *Ann Rheum Dis* 2019;78:1296–304.
- 92 Hinojosa-Azaola A, Romero-Diaz J, Vargas-Ruiz AG, *et al.* Venous and arterial thrombotic events in systemic lupus erythematosus. *J Rheumatol* 2016;43:576–86.
- 93 Ibañez D, Gladman DD, Urowitz MB. Adjusted mean systemic lupus erythematosus disease activity Index-2K is a predictor of outcome in SLE. *J Rheumatol* 2005;32:824–7.
- 94 Fasano S, Margiotta DP, Gualtierotti R, *et al.* The incidence of cardiovascular events in Italian patients with systemic lupus erythematosus is lower than in North European and American cohorts: implication of disease-associated and traditional risk factors as emerged by a 16-year retrospective GIRRCs study: GIRRCs=Gruppo Italiano di Ricerca in Reumatologia clinica E Sperimentale. *Medicine* 2018;97:e0370.
- 95 Tselios K, Gladman DD, Su J, *et al.* Evolution of risk factors for atherosclerotic cardiovascular events in systemic lupus erythematosus: a longterm prospective study. *J Rheumatol* 2017;44:1841–9.
- 96 Goldberg RJ, Urowitz MB, Ibañez D, *et al.* Risk factors for development of coronary artery disease in women with systemic lupus erythematosus. *J Rheumatol* 2009;36:2454–61.
- 97 Urowitz MB, Ibañez D, Gladman DD. Atherosclerotic vascular events in a single large lupus cohort: prevalence and risk factors. *J Rheumatology* 2007;34:70–5.
- 98 Chen H-L, Shen L-J, Hsu P-N, *et al.* Cumulative burden of glucocorticoid-related adverse events in patients with systemic lupus erythematosus: findings from a 12-year longitudinal study. *J Rheumatol* 2018;45:83–9.
- 99 Hanly JG, Li Q, Su L, *et al.* Cerebrovascular events in systemic lupus erythematosus: results from an international inception cohort study. *Arthritis Care Res* 2018;70:1478–87.
- 100 Siricheepchaiyan W, Narongroeknawin P, Pakchotanont R, *et al.* Lupus damage and waist circumference as the independent risk factors for cardiovascular disease in SLE patients from Phramongkutklao Hospital. *J Med Assoc Thai* 2016;99:290–300.
- 101 Fasano S, Margiotta DPE, Piero L, *et al.* Prolonged remission is associated with a reduced risk of cardiovascular disease in patients with systemic lupus erythematosus: a GIRRCs (Gruppo Italiano di Ricerca in Reumatologia clinica E Sperimentale) study. *Clin Rheumatol* 2019;38:457–63.
- 102 Ruiz-Arruza I, Lozano J, Cabezas-Rodriguez I, *et al.* Restrictive use of oral glucocorticoids in systemic lupus erythematosus and prevention of damage without worsening long-term disease control: an observational study. *Arthritis Care Res* 2018;70:582–91.
- 103 Mok CC, Tong KH, To CH, *et al.* Risk and predictors of arterial thrombosis in lupus and non-lupus primary glomerulonephritis: a comparative study. *Medicine* 2007;86:203–9.
- 104 Szalai AJ, Alarcón GS, Calvo-Alén J, *et al.* Systemic lupus erythematosus in a multiethnic US cohort (LUMINA). XXX: association between C-reactive protein (CRP) gene polymorphisms and vascular events. *Rheumatology* 2005;44:864–8.
- 105 Smržová A, Horák P, Skácelová M. Cardiovascular events in patients with systemic lupus erythematosus. *Cor Vasa* 2014;56:e145–52.
- 106 Gustafsson J, Gunnarsson I, Börjesson O, *et al.* Predictors of the first cardiovascular event in patients with systemic lupus erythematosus - a prospective cohort study. *Arthritis Res Ther* 2009;11:R186.
- 107 Martínez-Berriotxa A, Ruiz-Irastorza G, Egurbide M-V, *et al.* Transiently positive anticardiolipin antibodies and risk of thrombosis in patients with systemic lupus erythematosus. *Lupus* 2007;16:810–6.
- 108 Bessant R, Duncan R, Ambler G, *et al.* Prevalence of conventional and lupus-specific risk factors for cardiovascular disease in patients with systemic lupus erythematosus: a case-control study. *Arthritis Rheum* 2006;55:892–9.
- 109 Hsu C-Y, Lin Y-S, Su Y-J, *et al.* Effect of long-term hydroxychloroquine on vascular events in patients with systemic lupus erythematosus: a database prospective cohort study. *Rheumatology* 2017;56:2212–21.
- 110 Arnett DK, Blumenthal RS, Albert MA, *et al.* 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American heart association Task force on clinical practice guidelines. *Circulation* 2019;140:e596–646.
- 111 Ridker PM, Buring JE, Rifai N, *et al.* Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds risk score. *JAMA* 2007;297:611–9.
- 112 Visseren FLJ, Mach F, Smulders YM, *et al.* 2021 ESC guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021;42:3227–337.
- 113 Ballocca F, D'Ascenzo F, Moretti C, *et al.* Predictors of cardiovascular events in patients with systemic lupus erythematosus (SLE): a systematic review and meta-analysis. *Eur J Prev Cardiol* 2015;22:1435–41.
- 114 Mohammad AJ, Bakoush O, Sturfelt G, *et al.* The extent and pattern of organ damage in small vessel vasculitis measured by the vasculitis damage index (VDI). *Scand J Rheumatol* 2009;38:268–75.
- 115 Disveld IJM, Zoakman S, Jansen TLTA, *et al.* Crystal-proven gout patients have an increased mortality due to cardiovascular diseases, cancer, and infectious diseases especially when having tophi and/or high serum uric acid levels: a prospective cohort study. *Clin Rheumatol* 2019;38:1385–91.

- 116 Tektonidou MG, Wang Z, Ward MM. Brief report: trends in hospitalizations due to acute coronary syndromes and stroke in patients with systemic lupus erythematosus, 1996 to 2012. *Arthritis Rheumatol* 2016;68:2680–5.
- 117 Unlu O, Erkan D, Barbhaiya M, et al. The impact of systemic lupus erythematosus on the clinical phenotype of antiphospholipid antibody-positive patients: results from the AntiPhospholipid Syndrome Alliance for Clinical Trials and international Clinical Database and Repository. *Arthritis Care Res* 2019;71:134–41.
- 118 Souverein PC, Berard A, Van Staa TP, et al. Use of oral glucocorticoids and risk of cardiovascular and cerebrovascular disease in a population based case-control study. *Heart* 2004;90:859–65.
- 119 Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol* 2021;73:1108–23.
- 120 Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis* 2020;79:685–99.
- 121 Ramos-Casals M, Brito-Zerón P, Bombardieri S, et al. EULAR recommendations for the management of Sjögren's syndrome with topical and systemic therapies. *Ann Rheum Dis* 2020;79:3–18.
- 122 Tardif J-C, Kouz S, Waters DD, et al. Efficacy and safety of low-dose colchicine after myocardial infarction. *N Engl J Med* 2019;381:2497–505.
- 123 Ridker PM, Everett BM, Thuren T, et al. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 2017;377:1119–31.
- 124 Ulander L, Tolppanen H, Hartman O, et al. Hydroxychloroquine reduces interleukin-6 levels after myocardial infarction: the randomized, double-blind, placebo-controlled OXI pilot trial. *Int J Cardiol* 2021;337:21–7.