

Supplementary Table 4: Number of patients assessed through B- and T- cells analysis and frequency of B-cell, CD4⁺ and CD8⁺ T cell and subsets.

B cell subsets	ACA (n=19)	SCL70 (n=10)
Transitionnal	5.2 (\pm 0.9) %	5.3 (\pm 1.5) %
Naive	62.3 (\pm 4.5) %	53 (\pm 4.2) %
Unswitched memory	11.7 (\pm 3.6) %	14.2 (\pm 2.3) %
Switch memory	13.8 (\pm 2.3) %	19.1 (\pm 2) %
Double negative	6 (\pm 1.1) %	6.5 (\pm 0.7) %
Plasmablast	1(\pm 0.2) %	1.9 (\pm 0.7) %

Mean \pm /-SEM.

	ACA (n=17)	SCL70 (n=10)
CD4 CM	28.6 (\pm 4.1) %	28.8 (\pm 1.9) %
CD4 EM	12.7 (\pm 2.4) %	14 (\pm 2.9) %
CD4 EMRA	3.5 (\pm 1.1) %	6 (\pm 2.1) %
CD4 N	55.2 (\pm 5.8) %	51 (\pm 5.3) %
CD8 CM	8 (\pm 1.2) %	8 (\pm 2.4) %
CD8 EM	10.4 (\pm 1.8) %	7.5 (\pm 0.9) %
CD8 EMRA	41 (\pm 3.9) %	36.4 (\pm 5.2) %
CD8 N	40.6 (\pm 4) %	48.1 (\pm 4.7) %

Mean \pm /-SEM in percentage of cell population.