

*** RESPOND Investigators**

R Nasyrova, Central Rheumatology Institute, Moscow, Russia; E Parsik, North Estonian Regional Hospital, Tallinn, Estonia; K Otsa, East-Tallinn's Central Hospital, Tallinn, Estonia; I Butrimiene, Vilnius University Hospital, Vilnius, Lithuania; M Pileckyte, Kaunas Medical University Hospital, Kaunas, Lithuania; W Tlustochowicz, Centralny Szpital Kliniczny, Warszawa, Poland; E Kucharz, Samodzielny Publiczny Szpital, Katowice-Ochojec, Poland; Z Szekanecz, University of Debrecen, Debrecen, Hungary; G Poor, National Institute of Rheumatology and Physiotherapy, Budapest, Hungary; E Koo, Polyclinic of the Hospitaller Brothers of St. John of God, Budapest, Hungary; L Hodinka, National Institute of Rheumatology and Physiotherapy, Budapest, Hungary; Z Macejova, L. Pasteur's Faculty Hospital, Kosice, Slovak Republic; H Direskeneli, Marmara Universitesi Tip Fakultesi, Istanbul, Turkey; N Akkoc, Dokuz Eylul Universitesi Tip Fakultesi, Izmir, Turkey; Y Kabasakal, Ege Universitesi Tip Fakultesi, Izmir, Turkey; I Ertenli, Hacettepe Universitesi Tip Fakultesi, Ankara, Turkey; F Khatib, Arwyp Medical Suites, Kempton Park, South Africa; EM van Duuren, Jacaranda Hospital, Pretoria, South Africa; C Spargo, Vincent Pallotti Hospital, Pinelands, South Africa; I Louw, Panorama Hospital Medical Centre, Cape Town, South Africa; C Codreanu, Centrul de Boli, Bucharest, Romania; R Ionescu, Sp. Clinic Sf. Maria, Bucharest, Romania; RM Chireac, Sp. Clinic de Recuperare, Iasi, Romania; S Rednic, Sp. Clinic Judetean Cluj, Cluj, Romania; M Hammoudeh, Hamad Medical Corporation, Doha, Qatar; J Szechinski, Prywatny Gabinet Internistyczno-Reumatologiczny, Wroclaw, Poland; M Rell-Bakalarska, Przychodnia Przyklinikzna, Warszawa, Poland; D Avraham, Meir M.C. Dermatology, Kfar Saba, Israel; D Andersone, P. Stradins Clinical University Hospital, Riga, Latvia; P Keszthelyi, Bekes County Pandy Kalman Hospital, Gyula, Hungary; P Suranyi, Hajdu-Bihar County Kenezy Gyula Hospital, Debrecen, Hungary