

Appendix G.

Table S5. Mean (CV) Serum Pharmacokinetic Parameters of Infliximab: Pharmacokinetic population

Parameter		CT-P13 3 mg/kg (N=292)		INX 3 mg/kg (N=289)
Dose 1 (Week 0)				
C _{max} (µg/mL)	n=290	91.50 (35)	n=285	88.53 (37)
C _{min} (µg/mL)	n=284	15.76 (58)	n=278	16.76 (44)
T _{max} (h)	n=290	3.00 (1.83, 4.08)	n=285	2.12 (1.50, 5.17)
Dose 2 (Week 2)				
C _{max} (µg/mL)	n=288	111.88 (32)	n=285	105.07 (35)
C _{min} (µg/mL)	n=278	6.15 (75)	n=278	7.65 (85)
T _{max} (h)	n=288	2.25 (0.25, 4.03)	n=285	3.00 (0.50, 3.75)
Dose 3 (Week 6)				
C _{max} (µg/mL)	n=277	97.91 (40)	n=277	96.20 (33)
C _{min} (µg/mL)	n=267	1.51 (301)	n=262	1.42 (187)
T _{max} (h)	n=277	2.25 (0.08, 7.33)	n=277	2.97 (1.00, 3.33)
Dose 4 (Week 14)				
C _{max} (µg/mL)	n=269	90.25 (36)	n=263	85.25 (40)
C _{min} (µg/mL)	n=261	1.05 (136)	n=251	1.07(140)
T _{max} (h)	n=269	3.00 (1.17, 3.20)	n=263	2.42 (0.22, 4.00)
Dose 5 (Week 22)				
C _{max} (µg/mL)	n=258	90.82 (40)	n=253	86.26 (44)
C _{min} (µg/mL)	n=240	0.99 (390)	n=243	1.02 (297)
T _{max} (h)	n=258	3.00 (2.00, 3.25)	n=253	2.25 (2.00, 4.50)
C _{av,ss} (µg/mL)	n=240	47.10 (40)	n=241	44.37 (44)
PTF	n=240	1.86 (12)	n=241	1.90 (8)
Dose 6 (Week 30)				
C _{max} (µg/mL)	n=243	83.87 (38)	n=245	83.78 (35)
T _{max} (h)	n=243	2.08 (2.00, 3.58)	n=245	2.25 (0.10, 3.33)

Note: Note: C_{max} represents the observed maximum serum drug concentration; C_{min} represents minimum serum drug concentration immediately prior to next dose; T_{max} represents time to reach C_{max} and is reported here as the median (minimum, maximum). CV, coefficient of variation.