

Supplementary table

	Week 24			Week 52		
	ADD-ON (N = 115)	SWITCH (N = 111)	p value	ADD-ON (N = 115)	SWITCH (N = 111)	p value
DAS28 remission rate (%)	69.6	55.0	0.03*	72.2	70.3	0.77
Change in DAS28	-2.9 (1.3)	-2.7 (1.5)	0.41	-3.0 (1.4)	-3.0 (1.6)	0.79
EULAR good/moderate responders (%)	96.5	90.1	0.06	94.8	88.3	0.10
Boolean remission rate (%)	20.9	19.8	0.87	37.1	35.1	0.78
SDAI remission rate (%)	44.1	29.6	0.07	52.2	46.8	0.43
CDAI remission rate (%)	39.6	27.8	0.13	47.8	44.1	0.60
Change in PGA (mm)	-23 (26)	-25 (28)	0.99	-2.6 (29)	-3.0 (28)	0.38
Change in EGA (mm)	-34 (22)	-31 (23)	0.48	-3.5 (23)	-3.4 (22)	0.97
Change in pain VAS (mm)	-26 (26)	-25 (28)	0.36	-2.8 (30)	-3.0 (28)	0.75
Change in HAQ-DI	-0.4 (0.5)	-0.4 (0.6)	0.75	-0.4 (0.6)	-0.5 (0.6)	0.50
Change in ESR (mm/h)	-31.6 (25.1)	-34.5 (29.7)	0.69	-31.0 (25.4)	-34.0 (29.9)	0.70
Change in CRP (mg/dL)	-0.8 (3.8)	-1.7 (2.6)	0.49	-0.7 (3.8)	-1.7 (2.6)	0.25

Change in SJC28	-5.1 (3.8)	-5.4 (5.0)	0.66	-5.2 (4.0)	-5.7 (4.9)	0.32
Change in TJC28	-5.2 (5.0)	-4.9 (5.3)	0.65	-5.2 (5.3)	-5.3(5.6)	1.00

DAS28, disease activity score for 28 joints; EULAR, European League Against Rheumatism; SDAI, simplified disease activity index, CDAI, clinical disease activity index; PGA, patient global assessment, EGA, evaluator global assessment; EGA, evaluator general assessment; VAS, visual analogue scale; HAQ-DI, health assessment questionnaire disability index; ESR, erythrocyte sedimentation rate; CRP, C-reactive protein; SJC, swollen joint count; TJC, tender joint count

*, $p < 0.05$

Supplementary figure. (A) Percentage of patients achieving SDAI remission. (B) Percentage of patients achieving CDAI remission. (C) Results over time for percentage of patients achieving DAS28 remission with results at week 8. DAS28, disease activity score for 28 joints; SDAI, simplified disease activity index; CDAI, clinical disease activity index; .

