

Supplementary Table 1. Approximate increased incidence risk of CD and UC in patients with PsO, PsA, and AS compared to the general population

	Approximate increased incidence compared to the general population*	
	Crohn's Disease	Ulcerative Colitis
PsO	2–4 fold ⁽¹⁻⁴⁾	1–2 fold ⁽²⁻⁴⁾
PsA	2–3 fold ^(4, 5)	2 fold ^(4, 5)
AS	3 fold (IBD) ⁽⁶⁾	
^a Cited studies examined disease rates using different methodologies including incidence rates, relative risks, odds ratios and prevalence ratios		

References

1. Li WQ, Han JL, Chan AT, Qureshi AA. Psoriasis, psoriatic arthritis and increased risk of incident Crohn's disease in US women. *Annals of the rheumatic diseases*. 2013;72(7):1200-5.
2. Augustin M, Reich K, Glaeske G, Schaefer I, Radtke M. Co-morbidity and age-related prevalence of psoriasis: Analysis of health insurance data in Germany. *Acta Derm Venereol*. 2010;90(2):147-51.
3. Cohen AD, Dreiher J, Birkenfeld S. Psoriasis associated with ulcerative colitis and Crohn's disease. *Journal of the European Academy of Dermatology and Venereology : JEADV*. 2009;23(5):561-5.
4. Makredes M, Robinson D, Jr., Bala M, Kimball AB. The burden of autoimmune disease: a comparison of prevalence ratios in patients with psoriatic arthritis and psoriasis. *J Am Acad Dermatol*. 2009;61(3):405-10.
5. Egeberg A, Mallbris L, Warren RB, Bachelez H, Gislason GH, Hansen PR, et al. Association between psoriasis and inflammatory bowel disease: a Danish nationwide cohort study. *Br J Dermatol*. 2016;175(3):487-92.
6. Stolwijk C, Essers I, van Tubergen A, Boonen A, Bazelier MT, De Bruin ML, et al. The epidemiology of extra-articular manifestations in ankylosing spondylitis: a population-based matched cohort study. *Annals of the rheumatic diseases*. 2015;74(7):1373-8.