

1 **Ten-year natural course of early hip osteoarthritis in middle-aged persons with hip pain; a**

2 **CHECK study**

3

4 Online supplementary material

5

6 A.C. van Berkel, MD¹, D. Schiphof, PhD¹, J.H. Waarsing, PhD², J. Runhaar, PhD¹, J.M. van Ochten,

7 PhD¹, P.J.E. Bindels PhD, professor¹, S.M.A. Bierma-Zeinstra, PhD, professor^{1,2}

8 1 Department of General Practice, Erasmus Medical Center, Rotterdam, The Netherlands, 2

9 Department of Orthopaedics, Erasmus MC, University Medical Center, Rotterdam, The

10 Netherlands


11

12 Supplementary Figure S1 Flow-chart indicating the available data per time point.

13 Supplementary Table S1 Course of pain, physical functioning, and radiographic OA features
14 during follow-up, for total study group (n=588) and subdivided for
15 H-group (n=170) and H&K-group (n=418).

16 Supplementary Table S2 Course of pain, physical functioning, and radiographic OA features
17 during follow-up, for the no HR-group (n=518) and HR-group
18 (n=69) and for HR-group with LOCF.

19 Supplementary Table S3 Course of pain, physical functioning, and radiographic OA features
20 during follow-up with LOCF.



Supplementary Figure S1 Flow-chart indicating the available data per time point.

42 Supplementary Table S1. Course of pain, physical functioning, and radiographic OA features
 43 during follow-up, for total study group (n=588) and subdivided for H-group (n=170) and H&K-
 44 group (n=418).

	Baseline	T2	T5	T8	T10
WOMAC, mean (sd)					
- Pain (0-20)	5.4 (±3.4)	5.2 (±3.5)	5.1 (±4.0)	4.5 (±3.6)	4.7 (±3.8)
○ H-group	4.8 (±3.2)	4.3 (±3.5)	4.3 (±3.9)	3.5 (±3.4)	3.6 (±3.4)
○ H&K-group	5.7 (±3.5)	5.5 (±3.5)	5.4 (±4.0)	4.9 (±3.6)	5.1 (±3.9)
- Stiffness (0-8)	2.8 (±1.7)	2.6 (±1.6)	2.8 (±1.8)	2.4 (±1.8)	2.6 (±1.9)
○ H-group	2.5 (±1.7)	2.3 (±1.6)	2.5 (±1.8)	2.1 (±1.6)	2.1 (±1.8)
○ H&K-group	2.9 (±1.7)	2.7 (±1.6)	2.9 (±1.8)	2.5 (±1.8)	2.7 (±1.9)
- Physical function (0-68)	17.2 (±12.0)	16.4 (±12.0)	17.6 (±12.0)	16.3 (±13.0)	16.2 (±13.1)
○ H-group	14.7 (±11.1)	13.4 (±10.9)	14.3 (±12.8)	12.4 (±11.5)	12.2 (±10.7)
○ H&K-group	18.3 (±12.2)	17.6 (±12.3)	18.8 (±13.3)	17.8 (±13.2)	17.8 (±13.6)
NRS past week, mean (sd)	3.7 (±2.1)	3.7 (±2.3)	3.2 (±2.6)	2.7 (±2.5)	2.9 (±2.6)
○ H-group	3.4 (±2.2)	3.0 (±2.4)	3.0 (±2.6)	2.2 (±2.3)	2.5 (±2.5)
○ H&K-group	3.8 (±2.1)	4.0 (±2.2)	3.2 (±2.6)	2.9 (±2.5)	3.1 (±2.6)
Use any pain medication, n (%)	250 (43)	263 (48)	251 (47)	250 (49)	249 (50)
○ H-group	63 (38)	61 (39)	61 (40)	55 (38)	57 (40)
○ H&K-group	187 (46)	202 (52)	190 (50)	195 (53)	192 (54)
Hip pain, n (%)	588 (100)	374 (68)	301 (57)	267 (54)	247 (51)
○ H-group	170 (100)	114 (73)	90 (61)	71 (53)	57 (44)
○ H&K-group	418 (100)	260 (66)	211 (56)	196 (54)	190 (53)
Knee pain, n (%)	418 (71)	361 (65)	327 (62)	297 (58)	264 (53)
○ H-group	0 (0)	58 (37)	60 (40)	56 (39)	50 (37)
○ H&K-group	418 (100)	303 (77)	267 (70)	241 (66)	214 (60)
Morning stiffness (hip) <60 min, n (%)	326 (55)	287 (52)	272 (51)	239 (47)	228 (45)
○ H-group	101 (59)	85 (54)	77 (51)	61 (42)	58 (41)
○ H&K-group	225 (54)	202 (51)	195 (52)	178 (49)	170 (47)
Physically active (>30 min) for ≥3 times a week, n (%)	316 (55)	319 (60)	292 (56)	296 (58)	276 (56)
○ H-group	103 (62)	90 (58)	78 (52)	81 (57)	80 (56)
○ H&K-group	213 (53)	229 (60)	214 (57)	215 (59)	196 (56)
Cumulative sum of HR, n (%)	0 (0)	13 (2)	41 (7)	58 (10)	69 (12)
○ H-group	0 (0)	8 (5)	26 (15)	33 (19)	38 (22)
○ H&K-group	0 (0)	5 (1)	15 (4)	25 (6)	31 (7)
Cumulative sum of KR, n (%)	0 (0)	0 (0)	4 (1)	9 (2)	10 (2)
○ H-group	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
○ H&K-group	0 (0)	0 (0)	4 (1)	9 (2)	10 (2)
K/L grade ≥2 either hip, n (%)	110 (19)	128 (22)	151 (28)	183 (35)	253 (49)
○ H-group	38 (23)	40 (25)	48 (33)	54 (39)	64 (48)
○ H&K-group	72 (17)	88 (21)	103 (26)	129 (33)	189 (49)

K/L grade ≥ 2 either knee, n (%)	76 (13)	119 (20)	178 (37)	226 (51)	294 (69)
○ H-group	14 (8)	22 (14)	39 (26)	59 (42)	88 (63)
○ H&K-group	62 (15)	97 (25)	157 (42)	195 (55)	248 (70)
Clinical OA* either knee, n (%)	206 (35)	278 (47)	323 (55)	349 (59)	366 (62)
○ H-only	0 (0)	16 (9)	37 (22)	46 (27)	51 (30)
○ H&K	206 (50)	262 (63)	286 (68)	303 (73)	315 (75)
Painful internal rotation either hip, n (%)	322 (55)	197 (36)	190 (38)	166 (36)	179 (39)
○ H-only	101 (60)	61 (41)	51 (39)	43 (36)	49 (42)
○ H&K	221 (53)	136 (35)	139 (38)	123 (36)	129 (38)
Painful external rotation either hip, n (%)	160 (35)	86 (17)	115 (20)	86 (15)	89 (20)
○ H-only	45 (34)	28 (20)	31 (25)	26 (23)	22 (20)
○ H&K	115 (35)	58 (16)	84 (17)	60 (17)	67 (20)
Painful flexion either hip, n (%)	315 (54)	227 (42)	192 (39)	149 (32)	159 (35)
○ H-only	94 (56)	65 (43)	48 (38)	33 (29)	39 (35)
○ H&K	221 (54)	162 (41)	144 (39)	116 (33)	120 (35)

45 Values are: mean values \pm the standard deviation, or number (percentages %). *according to
 46 the clinical criteria of the American College of Rheumatology; once those clinical ACR criteria
 47 are satisfied, the case will be seen as clinical hip or knee OA. NRS= Numeric Rating Scale (0-10).
 48 WOMAC = Western Ontario and McMaster osteoarthritis index. K/L= Kellgren and Lawrence
 49 score.

50 Supplementary Table S2. Course of pain, physical functioning, and radiographic OA features
 51 during follow-up, for the no HR-group (n=518) and HR-group (n=69) and for HR-group with
 52 LOCF.

	Baseline	T2	T5	T8	T10
WOMAC, mean (sd)					
- Pain (0-20)					
▪ No HR-group	5.4 (±3.4)	5.1 (±3.5)	5.3 (±4.0)	4.7 (±3.6)	4.9 (±3.8)
▪ HR-group	5.7 (±3.9)	5.5 (±3.7)	4.0 (±4.0)	3.6 (±3.8)	3.1 (±3.6)
▪ ^a HR-group with LOCF	5.7 (±3.9)	6.4 (±3.9)	6.9 (±4.1)	7.1 (±4.3)	7.1 (±4.3)
- Stiffness (0-8)					
▪ No HR-group	2.8 (±1.7)	2.6 (±1.6)	2.9 (±1.8)	2.4 (±1.8)	2.6 (±1.9)
▪ HR-group	2.9 (±1.6)	2.6 (±1.7)	2.5 (±1.8)	2.1 (±1.8)	2.2 (±2.0)
▪ ^a HR-group with LOCF	2.9 (±1.6)	2.9 (±1.7)	3.1 (±1.8)	3.1 (±1.9)	3.1 (±1.9)
- Physical function (0-68)					
▪ No HR-group	16.9 (±11.9)	16.1 (±11.8)	17.8 (±13.3)	16.8 (±12.9)	16.9 (±13.1)
▪ HR-group	20.1 (±12.5)	18.5 (±13.6)	15.8 (±13.2)	13.3 (±13.5)	12.0 (±12.2)
▪ ^a HR-group with LOCF	20.1 (±12.5)	21.1 (±13.7)	23.9 (±13.9)	24.9 (±14.6)	24.9 (±14.6)
NRS past week, mean (sd)					
▪ No HR-group	3.6 (±2.1)	3.6 (±2.2)	3.2 (±2.6)	2.8 (±2.4)	3.0 (±2.7)
▪ HR-group	4.1 (±2.4)	4.1 (±2.6)	2.9 (±2.9)	2.3 (±2.5)	2.2 (±2.4)
▪ ^a HR-group with LOCF	4.1 (±2.4)	4.6 (±2.5)	4.9 (±2.7)	5.1 (±2.7)	5.1 (±2.7)
Use any pain medication, n (%)					
▪ No HR-group	221 (44)	227 (48)	222 (48)	220 (49)	224 (52)
▪ HR-group	29 (43)	35 (52)	28 (41)	29 (43)	24 (36)
▪ ^a HR-group with LOCF	29 (43)	36 (53)	43 (64)	45 (66)	45 (66)
Hip pain, n (%)					
▪ No HR-group	518 (100)	326 (68)	266 (58)	239 (54)	231 (53)
▪ HR-group	69 (100)	47 (71)	35 (56)	27 (51)	15 (29)
▪ ^a HR-group with LOCF	69 (100)	59 (87)	59 (88)	61 (90)	61 (90)
Knee pain, n (%)					
▪ No HR-group	386 (75)	328 (68)	295 (64)	265 (60)	240 (55)
▪ HR-group	31 (45)	32 (47)	32 (47)	31 (47)	23 (35)
▪ ^a HR-group with LOCF	31 (45)	34 (50)	32 (48)	34 (50)	34 (50)
Morning stiffness (hip) <60 min, n (%)					
▪ No HR-group	282 (54)	244 (50)	236 (51)	210 (47)	202 (46)
▪ HR-group	43 (62)	42 (62)	35 (52)	28 (43)	25 (38)
▪ ^a HR-group with LOCF	43 (62)	48 (71)	49 (73)	51 (75)	51 (75)
Physically active (>30 min) for ≥3 times a week, n (%)					
▪ No HR-group	282 (56)	283 (61)	257 (56)	256 (58)	239 (56)
▪ HR-group	34 (52)	35 (52)	35 (52)	40 (61)	37 (55)
▪ ^a HR-group with LOCF	34 (52)	34 (52)	35 (55)	38 (58)	38 (58)
Cumulative sum of HR, n (%)					
▪ No HR-group	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
▪ HR-group	0 (0)	13 (19)	41 (59)	58 (84)	69 (100)

Cumulative sum of KR, n (%)					
▪ No HR-group	0 (0)	0 (0)	4 (1)	8 (2)	9 (2)
▪ HR-group	0 (0)	0 (0)	0 (0)	1 (1)	1 (1)
K/L grade ≥ 2 either hip, n (%)					
▪ No HR-group	73 (14)	92 (20)	134 (30)	174 (39)	252 (59)
▪ HR-group	37 (54)	374 (66)	17 (61)	8 (73)	-
K/L grade ≥ 2 either knee, n (%)					
▪ No HR-group	64 (13)	103 (21)	166 (37)	221 (51)	293 (69)
▪ HR-group	12 (17)	16 (24)	30 (44)	33 (51)	42 (65)
Clinical OA* either hip, n (%)					
▪ No HR-group	125 (24)	171 (33)	205 (40)	226 (43)	243 (47)
▪ HR-group	34 (49)	43 (62)	51 (74)	54 (78)	54 (78)
Clinical OA* either knee, n (%)					
▪ No HR-group	194 (38)	253 (49)	296 (57)	320 (62)	335 (65)
▪ HR-group	12 (17)	24 (35)	26 (38)	28 (41)	30 (44)
Painful internal rotation any hip, n (%)					
▪ No HR-group	271 (53)	166 (34)	160 (35)	151 (34)	163 (37)
▪ HR-group	50 (73)	30 (55)	29 (74)	14 (70)	14 (78)
▪ [*] HR-group with LOCF	50 (73)	41 (60)	50 (75)	53 (79)	53 (79)
Painful external rotation any hip, n (%)					
▪ No HR-group	131 (33)	69 (13)	101 (22)	77 (18)	80 (19)
▪ HR-group	28 (50)	17 (32)	13 (45)	8 (44)	8 (67)
▪ [*] HR-group with LOCF	28 (50)	27 (42)	36 (56)	37 (58)	37 (58)
Painful flexion any hip, n (%)					
▪ No HR-group	265 (52)	199 (41)	174 (38)	138 (31)	146 (34)
▪ HR-group	48 (71)	28 (51)	17 (55)	17 (65)	12 (75)
▪ [*] HR-group with LOCF	48 (71)	39 (57)	42 (63)	45 (67)	45 (67)

53 Values are: mean values \pm the standard deviation, or number (percentages %). *according to

54 the clinical criteria of the American College of Rheumatology; once those clinical ACR criteria

55 are satisfied, the case will be seen as clinical hip or knee OA. NRS= Numeric Rating Scale (0-10).

56 WOMAC = Western Ontario and McMaster osteoarthritis index. K/L= Kellgren and Lawrence

57 score. ^{*}HR-group with LOCF: for the participants of the HR-group we also show the

58 observations with the last observation carried forward (LOCF) up to the last visit prior to HR.

59 Supplementary Table S3. Course of pain, physical functioning, and radiographic OA feature
 60 during follow-up with LOCF.

	Baseline	T2	T5	T8	T10
WOMAC, mean (sd)					
- Pain (0-20)	5.4 (±3.4)	5.3 (±3.6)	5.5 (±4.0)	5.0 (±3.8)	5.2 (±3.9)
○ H-group with LOCF for HR	4.8 (±3.2)	4.6 (±3.6)	5.1 (±3.9)	4.6 (±3.8)	4.8 (±3.7)
○ H&K-group with LOCF for HR	5.7 (±3.5)	5.6 (±3.6)	5.6 (±4.0)	5.1 (±3.7)	5.4 (±4.0)
- Stiffness (0-8)	2.8 (±1.7)	2.6 (±1.6)	2.9 (±1.8)	2.5 (±1.8)	2.7 (±1.9)
○ H-group with LOCF for HR	2.5 (±1.6)	2.4 (±1.6)	2.6 (±1.7)	2.4 (±1.8)	2.4 (±1.8)
○ H&K-group with LOCF for HR	2.9 (±1.7)	2.7 (±1.6)	3.0 (±1.8)	2.6 (±1.8)	2.8 (±1.9)
- Physical function (0-68)	17.2 (±12.0)	16.7 (±12.1)	18.6 (±13.5)	17.8 (±13.4)	18.0 (±13.6)
○ H-group with LOCF for HR	14.7 (±11.1)	14.2 (±11.4)	16.6 (±13.1)	15.8 (±12.9)	15.8 (±12.1)
○ H&K-group with LOCF for HR	18.3 (±12.2)	17.7 (±12.3)	19.3 (±13.6)	18.6 (±13.5)	18.8 (±14.0)
NRS past week, mean (sd)	3.7 (±2.1)	3.7 (±2.3)	3.4 (±2.7)	3.1 (±2.6)	3.3 (±2.8)
○ H-group with LOCF for HR	3.4 (±2.2)	3.2 (±2.3)	3.6 (±2.6)	3.1 (±2.7)	3.3 (±2.8)
○ H&K-group with LOCF for HR	3.8 (±2.1)	4.0 (±2.2)	3.4 (±2.7)	3.1 (±2.6)	3.4 (±2.7)
Use any pain medication, n (%)	250 (43)	264 (48)	266 (50)	266 (52)	270 (54)
○ H-group with LOCF for HR	63(38)	62 (40)	72 (48)	68 (47)	73 (51)
○ H&K-group with LOCF for HR	187 (46)	202 (52)	194 (51)	198 (54)	197 (55)
Hip pain, n (%)	588 (100)	386 (70)	325 (61)	301 (59)	293 (58)
○ H-group with LOCF for HR	170 (100)	121 (77)	108 (72)	96 (66)	85 (59)
○ H&K-group with LOCF for HR	418 (100)	265 (67)	217 (57)	205 (56)	208 (58)
Knee pain, n (%)	418 (71)	363 (66)	327 (62)	300 (59)	275 (55)
○ H-group with LOCF for HR	0 (0)	57 (36)	54 (36)	53 (36)	50 (35)
○ H&K-group with LOCF for HR	418 (100)	306 (77)	273 (72)	247 (68)	225 (63)
Morning stiffness(hip)<60 min, n (%)	326 (55)	293 (53)	286 (54)	262 (51)	254 (50)
○ H-group with LOCF for HR	101 (59)	90 (57)	87 (58)	78 (53)	75 (52)
○ H&K-group with LOCF for HR	225 (54)	203 (51)	199 (53)	184 (50)	179 (49)
Physically active (>30 min) for ≥3 times a week, n (%)	316 (55)	318 (60)	292 (56)	294 (58)	277 (56)
○ H-group with LOCF for HR	103 (62)	89 (58)	80 (54)	82 (57)	82 (58)
○ H&K-group with LOCF for HR	213 (53)	229 (60)	212 (57)	212 (58)	195 (55)
Painful internal rotation either hip, n (%)	322 (55)	208 (38)	211 (40)	205 (40)	217 (43)
○ H-only with LOCF for HR	101 (60)	68 (43)	67 (45)	67 (46)	73 (51)
○ H&K with LOCF for HR	221 (53)	140 (35)	477 (38)	138 (38)	144 (40)
Painful external rotation either hip, n (%)	160 (35)	96 (19)	138 (26)	115 (23)	118 (24)
○ H-group with LOCF for HR	45 (34)	34 (23)	45 (30)	44 (31)	39 (28)
○ H&K-group with LOCF for HR	115 (28)	62 (17)	93 (25)	71 (20)	79 (22)
Painful flexion either hip, n (%)	315 (54)	238 (43)	217 (41)	183 (36)	192 (38)
○ H H-group with LOCF for HR	94 (56)	72 (46)	65 (38)	57 (39)	61 (42)
○ H&K-group with LOCF for HR	221 (54)	166 (42)	152 (40)	126 (34)	131 (36)

61 Values are: mean values ± the standard deviation, or number (percentages %). *according to
 62 the clinical criteria of the American College of Rheumatology; once those clinical ACR criteria

63 are satisfied, the case will be seen as clinical hip or knee OA. NRS= Numeric Rating Scale (0-10).
64 WOMAC = Western Ontario and McMaster osteoarthritis index. K/L= Kellgren and Lawrence
65 score. *With LOCF for HR:* for the participants who received an HR we show the observations
66 with the last observation carried forward (LOCF) up to the last visit prior to HR.