

Supplementary Table 1. Descriptive baseline results for quantitative demographic variables and clinical outcome measures

Age						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	57.1	13.2	337	0.72	55.69	58.51
Non-responders	59.9	12.0	102	1.19	57.57	62.23
Persisting pain	57.1	13.0	128	1.15	54.85	59.35
Disease Duration						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	2.5	2.6	337	0.14	2.22	2.78
Non-responders	2.7	2.8	102	0.28	2.16	3.24
Persisting pain	2.5	2.5	128	0.22	2.07	2.93
DAS28-CRP						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	4.9	0.9	337	0.05	4.80	5.00
Non-responders	4.9	1.0	102	0.10	4.71	5.09
Persisting pain	5.3	1.0	128	0.09	5.13	5.47
PD-Q						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	14.0	6.8	287	0.40	13.21	14.79
Non-responders	15.5	7.1	78	0.80	13.92	17.08
Persisting pain	15.5	6.5	111	0.62	14.29	16.71
RAID						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	5.8	2.0	332	0.11	5.58	6.02
Non-responders	6.0	1.9	100	0.19	5.63	6.37
Persisting pain	6.6	1.7	125	0.15	6.30	6.90
PHQ-9						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	7.3	5.1	328	0.28	6.75	7.85
Non-responders	8.4	5.4	98	0.55	7.33	9.47
Persisting pain	7.9	5.1	123	0.46	7.00	8.80
HAQ-DI						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	1.1	0.7	333	0.04	1.02	1.18
Non-responders	1.2	0.6	101	0.06	1.08	1.32
Persisting pain	1.3	0.6	125	0.05	1.19	1.41

PCS						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	19.7	12.4	328	0.68	18.36	21.04
Non-responders	21.5	12.3	96	1.26	19.04	23.96
Persisting pain	24.1	12.5	123	1.13	21.89	26.31
VAS pain						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	68.3	13.0	337	0.71	66.91	69.69
Non-responders	69.0	12.8	102	1.27	66.52	71.48
Persisting pain	73.8	13.0	128	1.15	71.55	76.05
NRS activity						
	Mean	SD	N	SEM	95%CI(low)	95%CI(hi)
Reference	6.7	1.5	337	0.08	6.54	6.86
Non-responders	6.9	1.4	102	0.14	6.63	7.17
Persisting pain	7.2	1.4	128	0.12	6.96	7.44